
LITERATURA

1. Alexopoulos GS, Meyers BS, et al. The course of geriatric depression with „reversible dementia“: a controlled study. *Am. J. Psychiatry* 1993;150:1693–1699.
2. Blumer D. Antidepressant and double antidepressant treatment for the affective disorder of epilepsy. *J. Clin. Psychiatry* 1997;58:3–11.
3. Carney RM, Rich MW, Freeland KE, et al. Major depressive disorder predicts cardiac events in patients with coronary artery disease. *Psychosom. Med.* 1988;50:627–633.
4. Cassem EH. Depression and anxiety secondary to medical illness. *Psychiatr. North. Am.* 1990;13:597–612.
5. Cohen-Cole SA, Stoudemire A. Major depression and physical illness: special considerations in diagnosis and biologic treatment. *Psychiatr. North. Am.* 1987;10:1–17.
6. Freeland KE, Carney RM, Lustman PJ, et al. Major depression in coronary artery disease: patients with vs without a prior history of depression. *Psychosom. Med.* 1992;54:416–421.
7. Holmgren A, Wise M, et al. Pain management. V: Wise MG, Rundell JR (eds.): *Textbook of consultation-liaison psychiatry*, American Psychiatric Publishing, Washington 2002: s. 989–338–1013.
8. Herman E, Hovorka J, Praško J, Jokl I, Ivanco A, Syrovátka J. *Deprese u somaticky nemocných*. Maxdorf, Praha. 2002;s. 1–112.
9. Hovorka J, Herman E, Němcová I. Treatment of interictal depression in patients with epilepsy. *Epilepsy and Behavior* 2000;6:444–447.
10. Hovorka J, Herman E. Neuropsychiatrické aspekty u epilepsie. *Ces Slov Neurol Neurochir (tematické suppl. březen 2001):*1–16.
11. Janicak PG. *Handbook of psychopharmacotherapy* Baltimore, Lippincott Williams and Wilkins 1999.
12. Katon W. The epidemiology of depression in medical care. *Int. J. Psychiatry Med.* 1987;17:93–112.
13. Kroenke K, Spitzer RL, Williams JB, et al. Physical symptoms in primary care: predictors of psychiatric disorders and functional impairment. *Arch. Fam. Med.* 1994;3:774–779.
14. Roose SP, Laghrissi-Thode F, et al. Comparison of paroxetine and nortriptyline in depressed patients with ischemic heart disease. *JAMA* 1998;279:287–291.
15. Rouchell AM, Pounds R, Tierney JG. Depression. V: Wise MG, Rundell JR (eds.): *Textbook of consultation-liaison psychiatry*, American Psychiatric Publishing, Washington, 2002; s. 307–338.

16. Simon GE, VonKoftt M, Piccinelli M, Fullerton C, Ormel J. An international study of the relation between somatic symptoms and depression. *N Engl J Med* 1999;341:1329–1335.
17. Smith BD, Salzman C. Do benzodiazepines cause depression? *Hospital and Community Psychiatry*. 1996;42:1101–1102.
18. Švestka J. Antidepresiva a bolestivá porucha. Jsou analgeticky stejně účinná noradrenergní, serotonergní a smíšená (noradrenergní + serotonergní) antidepresiva? *Bolest* 3 1999;3:119–126.
19. Watson CP, Vernich L, Chipman M, et al. Nortriptyline versus amitriptyline in postherpetic neuralgia: a randomized trial. *Neurology* 1998;51:1166–1171.
20. Wells KB, Rogers W, Burnam MA. Psychiatric disorder in a sample of general population with and without chronic medical conditions. *Am. J. Psychiatry* 1988;145:976–981.
21. Winokur G. The concept of secondary depression and its relationship to comorbidity. *Psychiatr. Clin. North. Am.* 1990;123:567–583.
22. Yates WR. Epidemiology of Psychiatric Disorders in the Medically Ill. V: Psychiatric treatment of the medically ill. Robinson RG, Yates WR (eds.) Marcel Dekker, New York 1999; s. 41–64.