

References

- Adema, F., Leenhouts, P.W. & van Welzen, P.C. (1994). Sapindaceae. *Flora Malesiana*, Ser. I, 11: 419–768. National Herbarium of the Netherlands, Leiden.
- Agudelo-H., C.A. (2008). Amaranthaceae. *Flora de Colombia* 23. Universidad Nacional de Colombia, Bogotá D.C.
- Airy Shaw, H. K. (1953). Thymelaeaceae—Gonyystyloideae. *Flora Malesiana*, Ser. I, 4: 349–365. National Herbarium of the Netherlands, Leiden.
- Alverson, W. S. (2004). Bombacaceae. In: N. Smith, S.A. Mori, A. Henderson, D.W. Stevenson & S.V. Heald (eds), *Flowering Plants of the Neotropics*, pp. 55–58. Princeton University Press, Princeton.
- Alverson, W. S., Whitlock, B.A., Nyffeler, R., Bayer, C. & Baum, D.A. (1999). Phylogeny of the core Malvales: evidence from *ndhF* sequence data. *Amer. J. Bot.* 86: 1474–1486.
- Anderberg, A.A., Baldwin, B. G., Bayer, R. G., Breitwieser, J., Jeffrey, C., Dillon, M. O., Eldenäs, P., Funk, V.A., Garcia-Jacas, N., Hind, D. J., N., Karis, P. O., Lack, H. W., Nesom, G. L., Nordenstam, B., Oberprieler, Ch., Panero, J. L., Puttock, C., Robinson, H., Stuessy, T. F., Susanna, A., Urtubey, E., Vogt, R., Ward, J. & Watson, L. E. ([2006]2007). Compositae. In: J. Kadereit & C. Jeffrey (eds), *The Families and Genera of Vascular Plants* 8: 561–588. Springer-Verlag, Berlin.
- Anderson, W. R. (2004). Malpighiaceae. In: N. Smith, S.A. Mori, A. Henderson, D.W. Stevenson & S.V. Heald (eds), *Flowering Plants of the Neotropics*, pp. 299–232. Princeton University Press, Princeton.
- Anderson, W. R., Anderson, C. & Davis, C. C. (2006). Malpighiaceae. <http://herbarium.lsa.umich.edu/malpigh/index.html>
- Andersson, L. (1998). Marantaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 4: 278–293. Springer-Verlag, Berlin.
- Andreatta, R. H. P. (1995). Revisão das espécies brasileiras do gênero *Smilax* Linnaeus (Smilacaceae). 2nd edition. São Paulo.
- Angiosperm Phylogeny Group [APG] (1998). An ordinal classification for the families of flowering plants. *Ann. Missouri Bot. Gard.* 85: 531 – 553.
- Angiosperm Phylogeny Group [APG] (2003). An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG II. *Bot. J. Linn. Soc.* 141: 399 – 436.
- Angiosperm Phylogeny Group [APG] (2009). An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Bot. J. Linn. Soc.* 161: 105 – 121.
- Archer, R. H. (2000). Ochnaceae. In: O.A. Leistner (ed.), *Seeds of Southern Africa, Families and Genera*. Strelitzia 10. South African National Biodiversity Institute, Pretoria.
- Ashton, P. S. (1982). Dipterocarpaceae. *Flora Malesiana*, Ser. I, 9: 237–552. National Herbarium of the Netherlands, Leiden.
- Ashton, P. S. (2003). Dipterocarpaceae. In: K. Kubitzki & C. Bayer (eds), *The Families and Genera of Vascular Plants* 4: 182–197. Springer-Verlag, Berlin.
- Austin, D. F. (1975). Flora of Panama: part 9. Family 164: Convolvulaceae. *Ann. Missouri Bot. Gard.* 62 (1): 157–224.
- Austin, D. F. (1982). Convolvulaceae. *Flora of Ecuador* 15. University of Gothenburg, Gothenburg.
- Aymard, C. G. A., Cuello, A. N. L., Berry, P. E., Rudd, V. E., Cowan, R. S., Fantz, P. R., Maxwell, R. H., Stirton, C. H., Poppendieck, H.-H., Lima, H. C. de, Fortunato, R. H., Stergiou, B., de Enrich, N. X., Neill, D. A., Pennington, R. T. & Gil, C. (1999). Fabaceae (Leguminosae-Papilionoideae). In: P. E. Berry, K. Yatskivych & B. K. Holst (eds), *Flora of the Venezuelan Guayana* 5: 231–433. Missouri Botanical Garden Press, St. Louis.
- Aymard, G., Berry, P. E. & Eriksen, B. (2004). Polygalaceae. In: P. E. Berry, K. Yatskivych & B. K. Holst (eds), *Flora of the Venezuelan Guayana* 8: 316–347. Missouri Botanical Garden Press, St. Louis.
- Backeberg, C. (1958–1962). *Die Cactaceae*. 6 vols. G. Fisher, Jena.
- Backer, C. A. (1949). Amaranthaceae. *Flora Malesiana*, Ser. I, 4: 69–98. National Herbarium of the Netherlands, Leiden.
- Backlund, M., Oxelman, B. & Bremer, B. (2000). Phylogenetic relationships within the Gentianales based on *ndhF* and *rbcL* sequences, with particular reference to the Loganiaceae. *Amer. J. Bot.* 87: 1029–1043.
- Bakhuisen van den Brink, R. C. (1943). A contribution to the knowledge of the Melastomataceae occurring in the Malay Archipelago especially in the Netherlands East Indies. *Rec. Trav. Bot. Néerl.* 40: 1–391. [Reprinted in *Meded. Bot. Mus. Herb. Rijks Univ.* 91: 1–391 (1947).]
- Bakker, M. E. (2002). *Annonaceae: Neotropical Genera and Species*. Expert Center for Taxonomic Identification, University of Amsterdam, Amsterdam. [CD-ROM.]
- Ballard, H. E. (2007). Violaceae. In: V. H. Heywood, R. K. Brummitt, A. Culham & O. Seberg (eds), *Flowering Plant Families of the World*, pp. 331–333. Royal Botanic Gardens, Kew.
- Ballard, H. E., de Paula-Souza, J. & Wahlert, G. A. (2014). Violaceae. In K. Kubitzki, *Flowering Plants. Eudicots: Malpighiales*. Springer, Heidelberg.
- Barker, W. R. (1980). Taxonomic revisions in Theaceae in Papuasia. I. *Gordonia*, *Ternstroemia*, *Adinandra* and *Archboldiodendron*. *Brunonia* 3 (1): 1–60.
- Barlow, B. (1997). Loranthaceae. *Flora Malesiana*, Ser. I, 13: 209–401. National Herbarium of the Netherlands, Leiden.
- Barneby, R. C. (1970). Revision of neotropical Menispermaceae tribe Tinosporae. *Mém. New York Bot. Gard.* 20: 81–158.
- Barneby, R. C. & Krukoff, B. A. (1971). Supplementary notes on American Menispermaceae. VIII. A generic survey of the American Triclisieae and Anomospermeae. *Mém. New York Bot. Gard.* 22: 1–89.
- Barneby, R. C., Grimes, J. W., Berry, P. E., Brunner, D., Forero, E., Cárdenas, L., De Martino, G., Hopkins, H. C. F. & de Lamare Occhioni, E. M. (2001). Mimosaceae (Leguminosae-Mimosoideae). In: P. E. Berry, B. K. Holst & K. Yatskivych (eds), *Flora of the Venezuelan Guayana* 6: 580–685. Missouri Botanical Garden Press, St. Louis.
- Barneby, R. C., Stergiou, B., Cowan, R. S., Berry, P. E., Zarucchi, J., Wunderlin, R., Kearns, D., da Silva, M. F., Sprada, A., Velazquez, D., Xena, N. & Aymard, C. G. A. (1998). Caesalpiniaceae (Leguminosae-Caesalpinioidae). In: P. E. Berry, B. K. Holst & K. Yatskivych (eds), *Flora of the Venezuelan Guayana* 4: 1–121. Missouri Botanical Garden Press, St. Louis.
- Barnett, E. C. (1942). The Fagaceae of Thailand and their geographical distribution. *Trans. Bot. Soc. Edinburgh* 33 (3): 327–343.
- Barringer, K. & Nevling, L. I. (2004). Thymelaeaceae. In: N. Smith, S.A. Mori, A. Henderson, D.W. Stevenson, & S.V. Heald (eds), *Flowering Plants of the Neotropics*, pp. 372 – 374. Princeton University Press, New Jersey.
- Barthlott, W. & Hunt, D. R. (2001). Seed diversity in the Cactaceae subfamily Cactoideae. *Succulent Pl. Res.* 5: 1–173.
- Baum, D. A. (1995). The comparative pollination and floral biology of Baobabs (*Adansonia*- Bombacaceae.) *Ann. Missouri Bot. Gard.* 82: 322–348.

- Baum, D.A., Alverson, W.S. & Nyffeler, R. (1998). A durian by any other name: taxonomy and nomenclature of the core Malvales. *Harvard Pap. Bot.* 3: 315–330.
- Bayer, C., Fay, M.F., De Bruijn, A.Y., Savolainen, V., Mortan, C.M., Kubitzki, K.K., Alverson, W.S. & Chang, M.W. (1999). Support for an expanded family concept of Malvaceae within a recircumscribed order Malvales: a combined analysis of plastid *atpB* and *rbcL* DNA sequences. *Bot. J. Linn. Soc.* 129: 267–303.
- Bayer, C. & Kubitzki, K. (2003). Malvaceae. In: K. Kubitzki & C. Bayer (eds), *The Families and Genera of Vascular Plants* 5: 225–311. Springer-Verlag, Berlin. (Subfamily Byttnerioideae, pp. 241–247; subfamily Bombacoideae, pp. 271–277; subfamily Malvoideae, pp. 277–311.)
- Beentje, H.J. (1993). Pandanaceae. *Flora of Tropical East Africa*. Balkema, Rotterdam.
- Beentje, H.J. (2006). Ericaceae. *Flora Tropical East Africa*. Royal Botanic Gardens, Kew.
- Beentje, H.J. (2008a). Hymenophyllaceae. *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Beentje, H.J. (2008b). Aspleniaceae. *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Beentje, H.J. (2015). *The Kew Plant Glossary*, 2nd edition. Royal Botanic Gardens, Kew.
- Beier, B.-A., Chase, M.W. & Thulin, M. (2003). Phylogenetic relationships and taxonomy of subfamily Zygophylloideae (Zygophyllaceae) based on molecular and morphological data. *Pl. Syst. Evol.* 240: 11–39.
- Berg, C.C. (1972). Olmedieae and Brosimeae. *Flora Neotropica, Monograph* 7. New York Botanical Garden, New York.
- Berg, C.C. (1977). Revisions of African Moraceae (excluding *Dorstenia* & *Ficus*). *Bull. Jard. Bot. Natl. Belg.* 47 (3/4): 267–407.
- Berg, C.C. (1992). Urticaceae. In: *Flora of the Guianas*, Ser. A, fasc. 11, fam. 23. Koenigstein.
- Berg, C.C. (2003). Moraceae – *Ficus*. *Flora Malesiana*, Ser. I, vol. 1, part 2. National Herbarium of the Netherlands, Leiden.
- Berg, C.C. & Dahlberg, S.V. (2001). A revision of *Celtis* subgenus *Mertensia*. *Brittonia* 53 (1): 66–81.
- Berg, C.C. & Hijman, M.E.E. (1999). The genus *Dorstenia* (Moraceae). Bergen University.
- Bidgood, S., Verdcourt, B. & Vollesen, K. (2006). Bignoniacae. In: H.J. Beentje & S.A. Ghazanfar (eds), *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Biffin, E., Craven, L.A., Crisp, M.D. & Gadek, P.A. (2006). Molecular systematics of *Syzygium* and allied genera (Myrtaceae): evidence from the chloroplast genome. *Taxon* 55: 79–94.
- Bohs, L. (1994). Cyphomandra (Solanaceae). *Flora Neotropica, Monograph* 63. New York Botanical Garden, New York.
- Bornstein, A.J. (1989). Taxonomic studies in the Piperaceae — I. The pedicellate pipers of México and Central América (Piper subg. *Arctotonia*). *J. Arnold Arbor.* 70: 1–55.
- Bramley, G.L.C., Go, R. & de Kok, R.P.J. with contributions by Wearn, J.A. & Mabberley, D.J. (2011). Lamiaceae. In: E. Soepadmo, L.G. Saw, R.C.K. Chung & R. Kiew (eds), *Tree Flora of Sabah and Sarawak* 7. Forest Research Institute Malaysia; Sabah Forestry Department; Sarawak Forestry Department, Kuala Lumpur.
- Bremer, K. (1994). Asteraceae: Cladistics & Classification. Timber Press, Portland, Oregon.
- Brenan, J.P.M. (1967). Leguminosae Subfamily Caesalpinioideae. *Flora of Tropical East Africa*, Crown Agents, London.
- Bricker, J.S. (1991). A revision of the genus *Crinodendron* (Elaeocarpaceae). *Syst. Bot.* 16 (1): 77–87.
- Bridson, D.M. & Verdcourt, B. (2003). Rubiaceae. In: G.V. Pope (ed.), *Flora Zambesiaca* 5 (2): 379–720. Royal Botanic Gardens, Kew.
- Bruce, E.A. & Lewis, J. (1960). Loganiaceae. In: C.E. Hubbard & E. Milne-Redhead (eds), *Flora of Tropical East Africa*. Crown Agents, London.
- Brummitt, R.K. (2007a). Scrophulariaceae. In: V.H. Heywood, R.K. Brummitt, A. Culham & O. Seberg (eds), *Flowering Plant Families of the World*, p. 311. Royal Botanic Gardens, Kew.
- Brummitt, R.K. (2007b). Simaroubaceae. In: V.H. Heywood, R.K. Brummitt, A. Culham & O. Seberg (eds), *Flowering Plant Families of the World*, p. 303–304. Kew Publishing, Royal Botanic Gardens, Kew.
- Brummitt, R.K. (2007c). Acanthaceae. In: V.H. Heywood, R.K. Brummitt, A. Culham & O. Seberg (eds), *Flowering Plant Families of the World*, pp. 23–24. Royal Botanic Gardens, Kew.
- Burkill, I.H. (1951). Dioscoreaceae. *Flora Malesiana*, Ser. I, 4: 293–335. National Herbarium of the Netherlands, Leiden.
- Callmander, M.W., Lowry, P.P.II, Forest, F., Devey, D.S., Beentje, H.J. & Buerki, S. (2012). Benstonea (Pandanaceae): characterization, circumscription, and distribution of a new genus of screw-pines, with a synopsis of accepted species. *Candollea* 67: 323–345.
- Camus, A. (1936–1938). *Les Chenes* (3 vols). Paul Le Chevalier, Paris.
- Chai, P.P.K. & Yii, P.C. (2002). Sapotaceae. In: E. Soepadmo, L.G. Saw & R.C.K. Chung (eds), (2004), *Tree Flora of Sabah and Sarawak* 4: 203–345. Forest Research Institute Malaysia; Sabah Forestry Department; Sarawak Forestry Department, Kuala Lumpur.
- Chase, M.W., Reveal, J.L. & Fay, M.F. (2009). A subfamilial classification for the expanded asparagalean families Amaryllidaceae, Asparagaceae and Xanthorrhoeaceae. *Bot. J. Linn. Soc.* 161: 132–136.
- Chase, M.W., Zmarzty, S., Lledó, M.D., Wurdack, K.J., Swensen, S.M., & Fay, M.F. (2002). When in doubt, put it in Flacourtiaceae: a molecular phylogenetic analysis based on plastid *rbcL* DNA sequences. *Kew Bull.* 57: 141–181. [Tribal groupings subsequently revised.]
- Chautems, A. (2009). Gesneriaceae. In: W. Milliken, B. Klitgaard & A. Baracat (eds), *Neotropikey — Interactive key and information resources for flowering plants of the Neotropics*. www.kew.org/neotropikey
- Cheek, M.R. (2007). Bombacaceae, p. 65; Brownlowiaceae, p. 71; Byttneriaceae, p. 76; Durionaceae, p. 134; Helicteraceae, p. 165; Malvaceae, pp. 201–202; Pentapetaceae, pp. 247–248; Sparmanniaceae, pp. 307–308; Sterculiaceae, pp. 311–312; Tiliaceae, pp. 321–322. In: V.H. Heywood, R.K. Brummitt, A. Culham & O. Seberg (eds), *Flowering Plant Families of the World*. Royal Botanic Gardens, Kew.
- Cheek, M. & Dorr, L. (2007). Sterculiaceae. In: H.J. Beentje & S.A. Ghazanfar (eds), *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Chen, S.-L., Li, D.Z., G.H., Wu, Z.I., Liu, L., Wang, Z.P., Sun, B.Z., Zhu, Z.D., Xia, N., Jia, L.Z., Guo, Z.H., Chne, W.L., Chen, X., Yang, G., Phillips, S.M., Stapleton, C., Soreng, R.J., Aiken, S.G., Tzvelev, N.N., Peterson, P.M., Renvoise, S.A., Okonova, M.V., Ammann, K., (2000). Poaceae. In: Wu, Z.Y., Okonova, M.V., Ammann, K. (eds). *Flora of China* 22. Beijing and St. Louis: Science Press and Missouri Botanical Garden Press, St. Louis.
- Chen, Y.-L. & Schirarend, C. (2007). Rhamnaceae. *Flora of China* 12: 115–355.
- Chew, W.-L. (1972). The genus *Piper* (Piperaceae) in New Guinea, Solomon Islands and Australia. *J. Arnold Arbor.* 53: 1–25.
- Clayton, W.D. & Renvoize, S.A. (1970–1982). Gramineae vols. 1–3. *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.

- Clayton, W. D. & Renvoize, S. A. (1986). *Genera Graminum*. HMSO, London.
- Coode, M. J. E. (1978a). A conspectus of Elaeocarpaceae in Papuasia. *Brunonia* 1: 131–302.
- Coode, M. J. E. (1978b). Combretaceae. In: J. S. Womersley (ed.), *Handbooks of the Flora of Papua New Guinea* 1: 43–110.
- Coode, M. J. E. (1983). A conspectus of Sloanea (Elaeocarpaceae) in the Old World. *Kew Bull.* 38: 347–427.
- Cornejo, X. (2009). Neotropical Capparaceae. In: W. Milliken, B. Klitgård & A. Baracat (2009 onwards), *Neotropikey – Interactive key and information resources for flowering plants of the Neotropics*. www.kew.org/science/tropamerica/neotropikey/
- Cornejo, X. & Iltis, H. H. (2009). The reinstatement of Beaumepsia (Capparaceae) and a key to the genera of Neotropical Capparaceae with variously stellate or peltate indumenta. *J. Bot. Res. Inst. Texas* 3: 683–689.
- Cowley, E. J. (1989). Smilacaceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Croat, T. B. (1978). *Flora of Barro Colorado Island*. Stanford University Press, Stanford, California.
- Darbyshire, I. (2006). Gesneriaceae. In: H. J. Beentje & S. A. Ghazanfar (eds), *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Davies, F. G. & Verdcourt, B. (1998). Sapindaceae. In: H. J. Beentje & C. M. Whitehouse (eds), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Davies, N., Kirkup, D. & Bramley, G. L. C. (2009). *Interactive Key to the Genera of Lamiaceae*. Royal Botanic Gardens, Kew. www.kew.org/herbarium/keys/lamiales/
- Davis, A. P. & Bridson, D. M. (2007). Rubiaceae. In: V. H. Heywood, R. K. Brummitt, A. Culham & O. Seberg (eds), *Flowering Plant Families of the World*, pp. 284–286. Royal Botanic Gardens, Kew.
- Davis, A. P., Govaerts, R., Bridson, D. M., Ruhsam, M., Moat, J. & Brummitt, N. A. (2009). A global assessment of distribution, diversity, endemism, and taxonomic effort in the Rubiaceae. *Ann. Missouri Bot. Gard.* 96: 68–78.
- Davis, C. C. & Anderson, W. R. (2010). A complete generic phylogeny of Malpighiaceae inferred from nucleotide sequence data and morphology. *Amer. J. Bot.* 97 (12): 2031–2048.
- de Wilde, W. J. J. O. (1972). Passifloraceae. *Flora Malesiana*, Ser. I, 7: 405–434. National Herbarium of the Netherlands, Leiden.
- de Wilde, W. J. J. O. (1975). Passifloraceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. Crown Agents, London.
- de Wilde, W. J. J. O. (2000). Myristicaceae. *Flora Malesiana*, Ser. I, 14. National Herbarium of the Netherlands, Leiden. [Keys and descriptions to all the species in South East Asia region.]
- de Wilde, W. J. J. O. & Duyfjes, B. E. E. (2008). Cucurbitaceae. In: T. Santisuk & K. Larsen (eds), *Flora of Thailand* 9 (4).
- de Wilde, W. J. J. O. & Duyfjes, B. E. E. (2014). Lagerstroemia (Lythraceae) in Malesia. *Blumea* 59: 113–122.
- Demissew, S. (2006). Asparagaceae. In: H. J. Beentje & S. A. Ghazanfar (eds), *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Descoings, B. (1967). Vitacées. I 24 Famille, Flore de Madagascar et des Comores, I–151. Museum National d'Histoire Naturelle Laboratoire de Phanerogamie, Paris, France.
- Ding Hou (1960). Thymelaeaceae. *Flora Malesiana*, Ser. I, 6: 1–48. National Herbarium of the Netherlands, Leiden.
- Ding Hou (1962). Celastraceae – I. *Flora Malesiana*, Ser. I, 6: 227–291. National Herbarium of the Netherlands, Leiden.
- Ding Hou (1964). Celastraceae – II. *Flora Malesiana*, Ser. I, 6: 389–421. National Herbarium of the Netherlands, Leiden.
- Ding Hou (1978). Anacardiaceae. *Flora Malesiana*, Ser. I, 8: 395–548. National Herbarium of the Netherlands, Leiden.
- Ding Hou (1984). Aristolochiaceae. *Flora Malesiana*, Ser. I, 10: 53–108. National Herbarium of the Netherlands, Leiden.
- Dransfield, J. & Beentje, H. (1995). *The Palms of Madagascar*. Royal Botanic Gardens, Kew.
- Dransfield, J., Uhl, N. W., Asmussen-Lange, C. B., Baker, W. J., Harley, M. M. & Lewis, C. E. (2008). *Genera Palmarum. Evolution and Classification of the Palms*. Royal Botanic Gardens, Kew.
- Ebihara, A., Dubuisson, J.-Y., Iwatsuki, K., Hennequin, S. & Ito, M. (2006). A taxonomic revision of Hymenophyllaceae. *Blumea* 51: 221–280.
- Edmonds, J. M. (2012). Solanaceae. In: H. J. Beentje (ed.), *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Edwards, P. J. E. (2005). Cyatheaceae. In: H. J. Beentje (ed.), *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Edwards, S. (2009). Neotropical Aristolochiaceae. In: W. Milliken, B. Klitgaard & A. Baracat (eds), *Neotropikey — Interactive key and Information Resources for Flowering Plants of the Neotropics*. www.kew.org/science/tropamerica/neotropikey/families/Aristolochiaceae.htm.
- Elffers, J., Graham, R. A. & DeWolf, G. P. (1964). Capparidaceae. In: C. E. Hubbard & E. Milne-Redhead (eds), *Flora of Tropical East Africa*. Crown Agents, London.
- El Hadidi, M. N. (1985). Zygophyllaceae. *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Ensermu, K. & Faden, R. W. (1997). Commelinaceae. In: S. Edwards, D. Sebsebe, D. & I. Hedberg (eds), *Flora of Ethiopia and Eritrea* 6: 339–374. Addis Ababa University, Addis Ababa and Uppsala University, Uppsala.
- Eriksen, B. & Persson, C. (2007). Polygalaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 9: 345–363. Springer-Verlag, Berlin.
- Faden, R. W. (1998). Commeliaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 4: 109–128. Springer-Verlag, Berlin.
- Fernández del Valle, Á. (2005). Violaceae. In: P. E. Berry, K. Yatskievych & B. K. Holst (eds), *Flora of the Venezuelan Guayana* 9: 445–465. Missouri Botanical Garden Press, St. Louis.
- Feuillet, C. & MacDougal, J. M. (2007). Passifloraceae. In: K. Kubitzki (ed.), *Families and Genera of Vascular Plants* 9: 270–281. Springer-Verlag, Berlin.
- Fiaschi, P. & Plunkett, G. M. (2009). Neotropical Araliaceae. In: W. Milliken, B. Klitgaard & A. Baracat (eds), *Neotropikey – Interactive Key and Information Resources for Flowering Plants of the Neotropics*. www.kew.org/neotropikey/
- Fischer, E. (1992). Systematic der Afrikanischen Lindernieae (Scrophulariaceae). Franz Steiner Verlag, Stuttgart.
- Fischer, E. (2004). Scrophulariaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 7: 333–432. Springer-Verlag, Berlin.
- Fischer, E., Theisen, I. & Lohman, L. G. (2004). Bignoniacae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 7: 9–38. Springer-Verlag, Berlin.
- Fontella, J. F. (1980). Estudos em Asclepiadaceae, XI. Chave para determinação dos gêneros de Asclepiadaceae Brasileiras e mais cultivadas no Brasil. *Bol. Mus. Bot. Mun. Curitiba* 42: 1–16, figs 1–11.
- Forman, L. L. (1986). Menispermaceae. *Flora Malesiana*, Ser. I, 10: 157–253. National Herbarium of the Netherlands, Leiden.
- Forrest, L. L., Hughes, M. & Hollingsworth, P. M. (2005). A phylogeny of Begonia using nuclear ribosomal sequence data and morphological characters. *Syst. Bot.* 30: 671–682.
- Friis, I. (1991). Urticaceae. In: Launert, E. & Pope, G. V. (eds), *Flora Zambeziaca* 9 (6). London. Royal Botanic Gardens, Kew.

- Friis, I. (1993). Urticaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 2: 612–630. Springer-Verlag, Berlin.
- Frodin, D. G. (2004). Araliaceae. In: N. Smith, S. A. Mori, A. Henderson, D. W. Stevenson & S. V. Heald (eds), *Flowering Plants of the Neotropics*, pp. 28–31. Princeton University Press, Princeton.
- Frodin, D. G. & Govaerts, R. (2003). *World Checklist and Bibliography of Araliaceae*. Kew Publishing, Royal Botanic Gardens, Kew.
- Fryxell, P.A. (1988). Malvaceae of Mexico. *Systematic Botany Monographs* vol. 25, pp. 522. The American Society of Plant Taxonomists, Ann Arbor.
- Fryxell, P.A. (2004a). Malvaceae. In: N. Smith, S. A. Mori, A. Henderson, D. W. Stevenson & S. V. Heald (eds), *Flowering Plants of the Neotropics*, pp. 232–235. Princeton University Press, Princeton.
- Fryxell, P.A. (2004b). Sterculiaceae. In: N. Smith, S. A. Mori, A. Henderson, D. W. Stevenson & S. V. Heald (eds), *Flowering Plants of the Neotropics*, pp. 360–362. Princeton University Press, Princeton.
- Gentian Research Network (online). Newest classification of Gentianaceae. www.rci.rutgers.edu/~gentian/classNEW123.htm.
- Gentry, A. H. (1980). Flora Neotropica: Bignoniacae — Part I (tribes Crescentieae and Tourrettieae). *Flora Neotropica Monograph* 25 (1): 1–150.
- Gentry, A. H. (1992). Flora Neotropica: Bignoniacae — Part II (tribe Tecomeae). *Flora Neotropica Monograph* 25 (2): 1–130.
- Gentry, A. H. (1993). Anacardiaceae, pp. 219–225; Burseraceae, pp. 299–304; Euphorbiaceae, pp. 403–424; Meliaceae, pp. 613–617; Rutaceae, pp. 749–758; Sapindaceae, pp. 762–771; Simaroubaceae, pp. 783–786. In: A. H. Gentry, *A Field Guide to the Families and Genera of Northwest South America (Colombia, Ecuador, Peru)*. The University of Chicago Press, Chicago.
- Gentry, H. S. (1982). *Agaves of Continental North America*. University of Arizona Press, Tucson, Arizona.
- Gibson, A. C. & Nobel, P. S. (1986). *The Cactus Primer*. Harvard University Press, London.
- Gillet, J. B. (1971). Leguminosae subfamily Papilionoideae. E. Milne-Redhead & R. M. Polhill. In: *Flora of Tropical East Africa*. HMSO, London.
- Gillet, J. B. (1991). Burseraceae. In: Polhill, R. M. (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Givnish, T. J., Millam, K. C., Berry, P. E. & Sytsma, K. J. (2007). Phylogeny of adaptive radiation, and historical biogeography of Bromeliaceae inferred from ndhF sequence data. *Also* 23: 3–26.
- Givnish, T. J., Millam, K. C., Evans, T. M., Hall, J. C., Pires, J. C., Berry, P. E. & Sytsma, K. J. (2004). Ancient vicariance or recent long-distance dispersal? Inferences about phylogeny and South American-African disjunctions in Rapateaceae and Bromeliaceae based on ndhF sequence data. *Int. J. Plant Sci.* 165 (4 Suppl.): S35–S54.
- Goetghebeur, P. (1998). Cyperaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 4: 141–190. Springer-Verlag, Berlin.
- Goldblatt, P. (1993). Iridaceae. In: G. V. Pope (ed.), *Flora Zambesiaca* 12 (4). Royal Botanic Gardens, Kew.
- Goldblatt, P. (1996). Iridaceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. Balkema, Rotterdam.
- Goldblatt, P. (1998). Iridaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 3: 295–333. Springer-Verlag, Berlin.
- Goldblatt, P. & Manning, J. C. (2008). *The Iris family: Natural History and Classification*. Timber Press, Portland, Oregon.
- Görts-van Rijn, A. R. A. & Mennega, A. M. W. (1994). Hippocrateaceae. *Fl. Guianas, Ser. A, Phanerogams*, 16: 3–81.
- Görts-van Rijn, A. R. A., Mennega, A. M. W., de Roon, A. C., Welle, B. J. H. & Detienne, P. (1994). *Flora of the Guianas: Series A. Phanerogams*: fascicle 16. I 10. Hippocrateaceae; I 12. Icacinaceae. Koeltz Scientific, Havlickuv Brod.
- Govaerts, R., Fernández Casas, F. J., Barker, C., Carter, S., Davies, S., Esser, H.-J., Gilbert, M., Hoffmann, P., Radcliffe-Smith, A., Steinmann, V., van Welzen, P., Whitmore, T. (2011). *World Checklist of Euphorbiaceae (with Pandaceae)*. Royal Botanic Gardens, Kew. www.kew.org/wcsp
- Govaerts, R., Frodin, D. G. & Pennington, T. D. (2001). *World Checklist and Bibliography of Sapotaceae*. Royal Botanic Gardens, Kew.
- Govaerts, R., Simpson, D. A., Bruhl, J., Egorova, T., Goetghebeur, P. & Wilson, K. (2007). *World Checklist of Cyperaceae*. Royal Botanic Gardens, Kew.
- Goyder, D. J. (2007). Apocynaceae. In: V. H. Heywood, R. K. Brummitt, A. Culham & O. Seberg (eds), *Flowering Plant Families of the World*, pp. 38–40. Royal Botanic Gardens, Kew.
- Goyder, D. J., Masinde, P. S., Meve, U. & Venter, H. J. T. (2012). Apocynaceae 2. *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Graham, S. A. (2007). Lythraceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants*, 9: 226–246. Springer-Verlag, Berlin.
- Graham, S. A. & Graham, A. (2014). Ovary, fruit and seed morphology of the Lythraceae. *Int. J. Plant Sci.* 175: 202–240.
- Graham, S. A., Hall, J., Systma, K. & Shi, S-H. (2005). Phylogenetic analysis of the Lythraceae based on four gene regions and morphology. *Int. J. Plant Sci.* 166: 995–1017.
- Grass Phylogeny Working Group II (2011). New grass phylogeny resolves deep evolutionary relationships and discovers C4 origins. *New Phytol.* 193: 304–312.
- Green, P. S. (1994). A revision of *Chionanthus* (Oleaceae) in S. America and the description of *Priogymnanthus*, gen. nov. *Kew Bull.* 49: 261–286.
- Green, P. S. (2004). Oleaceae. In: J. W. Kadereit (ed.), *The Families and Genera of Vascular Plants* 7: 296–306. Springer-Verlag, Berlin.
- Gu, C.-Z., Peng, C.-I. & Turland, N. J. (2007). Begoniaceae. In: Z.-Y. Wu, P. H. Raven & D.-Y. Hong (eds), *Flora of China* 13: 153–207. Science Press, Beijing; Missouri Botanical Garden Press, St. Louis.
- Hágaster, E., Soto, M., Salazar, G., Jimenez, R., Lopez, M. & Dressler, R. (2005). *Orchids of Mexico*. Productos Farmacéuticos, S. A. de C.V., Mexico City.
- Hallé, N. (1962). Monographie des Hippocrateacées d’Afrique occidentale. *Mém. Inst. Franç. Afrique Noire* 64: 1–246.
- Harley, R. M., Atkins, S., Budantsev, A. L., Cantino, P. D., Conn, B. J., Grayer, R., Harley, M. M., de Kok, R. P. J., Krestovskaja, T., Morales, R., Paton, A. J., Ryding, O. & Upson T. (2004). Labiateae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 7: 167–275. Springer-Verlag, Berlin.
- Haston, E., Richardson, J. E., Stevens, P. F., Chase, M. W. & Harris, D. J. (2009). The Linear Angiosperm Phylogeny Group (LAPG) III: a linear sequence of the families in APG III. *Bot. J. Linn. Soc.* 161: 128–131.
- Hekking, W. H. A. (1988). Violaceae. Part I: *Rinorea* and *Rinoreocarpus*. *Flora Neotropica, Monograph* 46. New York Botanical Garden, New York.
- Henderson, A. (2009). *Palms of Southern Asia*. New York Botanical Garden & Princeton University Press, New Jersey.
- Henderson, A., Galeano, G. & Bernal, R. (1995). *Field Guide to the Palms of the Americas*. Princeton University Press, New Jersey.
- Hennipman, E., Veldhoen, P., Kramer, K. U. & Price, M. G. (1990). Polypodiaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 1: 203–230. Springer-Verlag, Berlin.

- Herber, B. E. (2002[2003]). Thymelaeaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 5: 373–396. Springer-Verlag, Berlin.
- Heywood, V. H., Brummitt, R. K., Culham, A. & Seberg O. (eds) (2007). *Flowering Plant Families of the World*. Kew Publishing, Royal Botanic Gardens, Kew.
- Hind, D. J. N. (2007). Asteraceae (Compositae). In: V. H. Heywood, R. K. Brummitt, A. Culham & O. Seberg (eds), *Flowering Plant Families of the World*, pp. 46–52. Royal Botanic Gardens, Kew.
- Hind, D. J. N. (2009). Neotropical Asteraceae. In: W. Milliken, B. Klitgaard & A. Baracat (eds), *Neotropikey – Interactive key and information resources for flowering plants of the Neotropics*. www.kew.org/neotropikey
- Hoenselaar, K., Verdcourt, B., & Beentje, H. J. (2010). Cyperaceae. *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Hoffmann, P. (2007). Phyllanthaceae. In: V. H. Heywood, R. K. Brummitt, A. Culham & O. Seberg (eds), *Flowering Plant Families of the World*, pp. 250–252. Royal Botanic Gardens, Kew.
- Hoffmann, P., Kathriarachchi, H. & Wurdack, K. J. (2006). A phylogenetic classification of Phyllanthaceae (Malpighiales; Euphorbiaceae s.l.). *Kew Bull.* 61: 37–53.
- Holtum, R. (1981). Thelypteridaceae. *Flora Malesiana*, Ser. II, I: 331–560. National Herbarium of the Netherlands, Leiden.
- Holtum, R. E. (1982). Cyatheaceae. *Flora Malesiana*, Ser. II, I: 65–176. National Herbarium of the Netherlands, Leiden.
- Hong, D. & DeFillipps, R. A. (2000). Commelinaceae. In: Z. Wu & P. H. Raven (eds), *Flora of China* 24: 19–42. Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
- Hou, D. (1958). Rhizophoraceae. *Flora Malesiana*, Ser. I, 5: 429–494. National Herbarium of the Netherlands, Leiden.
- Hou, D., Larsen, K. & Larsen, S. S. (1996). Caesalpiniaceae (Leguminosae–Caesalpinoideae). *Flora Malesiana*, Ser. I, 12: 409–730. National Herbarium of the Netherlands, Leiden.
- Hovenkamp, P. H., Bosman, M. T. M., Hennipman, E., Nooteboom, H. P., Rödl-Linder, G. & Roos, M. C. (1998). Polypodiaceae. *Flora Malesiana*, Ser. II, 3: 1–234. National Herbarium of the Netherlands, Leiden.
- Howard, R. A. & Duno de Stefano, R. (1999). Icacinaceae. In: Berry, P., Yatskievych, K. & Holst, B. K. (eds), *Flora of the Venezuelan Guayana* 5. Missouri Botanical Garden Press, St. Louis.
- Huber, H. (1993). Aristolochiaceae. In: K. Kubitzki, J. G. Rohwer & V. Bittrich (eds), *The Families and Genera of Vascular Plants* 2: 129–137. Springer-Verlag, Berlin.
- Huft, M. J. (1994). Smilacaceae. In: G. Davidse, S. M. Sousa & A. O. Chater (eds), *Flora Mesoamericana* 6: 20–25. UNAM, Mexico.
- Hughes, M. & Pullan, M. (2007). Southeast Asian Begonia Database. <http://elmer.rbge.org.uk/begonia/>
- Hunt, D. R. (1994). Commelinaceae. In: G. Davidse, S. M. Sousa & A. O. Chater (eds), *Flora Mesoamericana* 6: 157–172. UNAM, Mexico.
- Hunt, D., Taylor, N. & Charles, G. (2006). *The New Cactus Lexicon*. 2 vols. dh Publications, Milborne Port.
- Hutchinson, J. & Dalziel, J. M. (1954). In: R. W. J. Keay. *Flora of West Tropical Africa* 1 (1). 2nd Edition. Crown Agents, London.
- Hutchinson, J. & Dalziel, J. M. (1958). Papilionaceae (Leguminosae – Papilionoideae). *Flora of West Tropical Africa*: 505–587.
- Iwatsuki, K. (1990). Hymenophyllaceae. In: K. Kubitzki, J. G. Rohwer & V. Bittrich (eds), *The Families and Genera of Vascular Plants* 1: 157–163. Springer-Verlag, Berlin.
- Jacobs, M. (1955). Malpighiaceae. *Flora Malesiana*, Ser. I, 5: 125–145. National Herbarium of the Netherlands, Leiden.
- Jacobs, M. (1960). Capparaceae. *Flora Malesiana*, Ser. I, 6: 61–105. National Herbarium of the Netherlands, Leiden.
- Jacobs, M. & Moore, D. M. (1971). Violaceae. *Flora Malesiana*, Ser. I, 7: 179–212. National Herbarium of the Netherlands, Leiden.
- Jacques, E. L. & Couto, A. V. S. (2009). Neotropical Begoniaceae. In: W. Milliken, B. Klitgaard & A. Baracat (eds), *Neotropikey – Interactive Key and Information Resources for Flowering Plants of the Neotropics*. www.kew.org/science/tropamerica/neotropikey/families/Begoniaceae.htm.
- Jacques-Felix, H. (1978). Les genres de Memecyleae (Melastomataceae) en Afrique, Madagascar et Mascareignes. *Adansonia* 18 (2): 221–235.
- Jansen-Jacobs, M. J., Lindeman, J. C. (1979). Theaceae. In: A. L. Stoffers & J. C. Lindeman (eds), *Flora of Suriname*, Volume 5, Part 1. Foundation Van Eedenfonds, Amsterdam.
- Janssen, T. & Rakotonrainibe, F. (2008). A revision of the indusiate scaly tree ferns (Cyatheaceae, Cyathea subgen. *Alsophila* sect. *Alsophila*) in Madagascar, the Comoros and the Seychelles. *Adansonia* 30: 221–376.
- Jaramillo, T. S., Muriel, P. & Balslev, H. (2004). Myristicaceae. *Flora of Ecuador* 72. University of Gothenborg, Gothenborg.
- Jarrett, F. M. (1960). Studies in *Artocarpus* and allied genera. *J. Arnold Arbor.* 41: 320–334.
- Jeffrey, C. (1967). Curcurbitaceae. In: *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Jeffrey, C. (1980). A review of the Cucurbitaceae. *Bot. J. Linn. Soc.* 81: 233–247.
- Jeffrey, C. (2005). A new system of Cucurbitaceae. *Bot. Zhurn.* 90: 332–335.
- Ji, Z., Tsai, Z.-W. & Meerow, A. W. (2000). Amaryllidaceae. In: Z. Wu & P. H. Raven (eds), *Flora of China* 24: 322–381. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis.
- Johnston, M. C. (1972). Rhamnaceae. In: E. Milne-Redhead & R. M. Polhill (eds), *Flora of Tropical East Africa*. Crown Agents, London.
- Johnston, M. C. & Johnston, L. V. (1978). *Rhamnus*. *Flora Neotropicana*, Monograph 20. New York Botanical Garden, New York.
- Jones, D. T. (1995). Burseraceae. In: E. Soepadmo & K. M. Wong (eds), *Tree Flora of Sabah and Sarawak* 1: 351–419. Forest Research Institute Malaysia, Kuala Lumpur.
- Jongkind, C. C. H. (1999). Combretaceae. *Flore du Gabon*. Museum National d'Histoire Naturelle, Paris.
- Judd, W. S., Campbell, C. S., Kellogg, E. A. & Stevens, P. F. (1999). *Plant Systematics: A Phylogenetic Approach*. Sinauer Associates, Inc., Sunderland, MA.
- Kalkman, C. (1997). Boraginaceae. *Flora Malesiana*, Ser. I, 13: 43–144. National Herbarium of the Netherlands, Leiden.
- Kanis, A. (1971). Ochnaceae. *Flora Malesiana*, Ser. I, 7: 97–119. National Herbarium of the Netherlands, Leiden.
- Kårehed, J. (2001). Multiple origin of the tropical forest tree family Icacinaceae. *Amer. J. Bot.* 88: 2259–2274.
- Kativu, S., Hoell, G., Björk, C. S. & Nordal, I. (2008). Anthericaceae. In: J. Timberlake & E. S. Martins (eds), *Flora Zambeziaca* 13 (1), Royal Botanic Gardens, Kew.
- Kaul, R. B. (1986). Evolution and reproductive biology of inflorescences in *Lithocarpus*, *Castanopsis*, *Castanea* and *Quercus* (Fagaceae). *Ann. Missouri Bot. Gard.* 73: 284–296.
- Kawasaki, M. L. (2006). Vochysiaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 9: 480–487. Springer-Verlag, Berlin.

- Kearns, D. E., Berry, P. E., Stevens, P. F., Cuello, N. L., Pipoly III, J. J., Robson, N. K. B., Holst, B. K., Kubitzki, K. & Weitzman, A. L. (1998). Clusiaceae. In: P. E. Berry, B. K. Holst & K. Yatskivych (eds), *Flora of the Venezuelan Guayana* 4: 248–329. Missouri Botanical Garden Press, St. Louis.
- Keng, H. (1978). Theaceae. In: F. S. P. Ng (ed.), *Tree Flora of Malaya* 3: 275–296. Longman, Malaysia.
- Kennedy, H., Andersson, L. & Hagberg, M. (1988). Marantaceae. *Flora of Ecuador* 32. University of Gothenburg, Gothenburg.
- Kern, J. H. (1974). Cyperaceae. *Flora Malesiana*, Ser. I, 7: 435–753. National Herbarium of the Netherlands, Leiden.
- Kessler, P. J. A. (1993a). Annonaceae. In: K. Kubitzki, J. G. Rohwer & V. Bittrich (eds), *The Families and Genera of Vascular Plants* 2: 93–129. Springer-Verlag, Berlin.
- Kessler, P. J. A. (1993b). Menispermaceae. In: K. Kubitzki, J. G. Rohwer & V. Bittrich (eds), *The Families and Genera of Vascular Plants* 2: 402–418. Springer-Verlag, Berlin.
- Kiew, R. (2005). *Begonias of Peninsular Malaysia*. Natural History Publications (Borneo) Sdn. Bhd.
- Killip, E. P. (1938). The American species of Passifloraceae. *Publ. Field Mus. Nat. Hist. Bot. Ser* 19: 1–613.
- Knapp, S. (2002). Solanum section Geminata (G. Don) Walpers (Solanaceae). *Flora Neotropica*, Monograph 84. New York Botanical Garden, New York.
- Knox, E. B., Muasya, A. M. & Muchhalo, N. (2008). The predominantly South American clade of Lobeliaceae. *Syst. Bot.* 33: 462–468.
- Kochummen, K. M. (1995). Burseraceae. In: E. Soepadmo & K. M. Wong (eds), *Tree Flora of Sabah and Sarawak* I. Sabah Forestry Department/Sarawak Forestry Department/Forest Research Institute Malaysia (FRIM), Kepong, Malaysia.
- Koechlin, J. (1964). Marantaceae. *Flore du Gabon*. 9: 91–158.
- Kokwaro, J. O. (1986). Anacardiaceae. In: R. M. Polhill, *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Koyama, T. (1978). Smilacaceae. In: H. Li, T. Lui, T. Huang, T. Koyama & C. E. De Vol (eds), *Flora of Taiwan* 5: 110–137.
- Koyama, T. (1984). A taxonomic revision of the genus *Heterosmilax* (Smilacaceae). *Brittonia* 36: 184–205.
- Kramer, K. U. (1990). Cyatheaceae. In: K. Kubitzki, J. G. Rohwer & V. Bittrich (eds), *The Families and Genera of Vascular Plants* 1: 94–99. Springer-Verlag, Berlin.
- Kramer, K. U. & Viane, R. (1990). Aspleniaceae. In: K. Kubitzki, J. G. Rohwer & V. Bittrich (eds), *The Families and Genera of Vascular Plants* 1: 52–56. Springer-Verlag, Berlin.
- Kühn, K. & Kubitzki, K. (1993). Myristicaceae. In: K. Kubitzki, J. G. Rohwer & V. Bittrich (eds), *The Families and Genera of Vascular Plants* 2: 457–467. Springer-Verlag, Berlin.
- Kujit, J. & Hansen, B. (2015). Loranthaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 12. Springer-Verlag, Berlin.
- Kupicha, F. K. (1983). Sapotaceae. *Flora Zambesiaca* 7: 210–247. Royal Botanic Gardens, Kew.
- Lammers, T. G. (2007a). Campanulaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 8: 26–59. Springer-Verlag, Berlin.
- Lammers, T. G. (2007b). *World Checklist and Bibliography of Campanulaceae*. Royal Botanic Gardens, Kew.
- Landrum, L. R. & Kawasaki, M. L. (1997). The genera of Myrtaceae in Brazil: an illustrated synoptic treatment and identification keys. *Brittonia* 49: 508–536.
- Larsen, K., Lock, J. M., Maas, H. & Maas, P. J. M. (1998). Zingiberaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 4: 109–128. Springer-Verlag, Berlin.
- Launert, E. (1968). Malpighiaceae. In: E. Milne-Redhead & R. M. Polhill (eds), *Flora of Tropical East Africa*. Crown Agents, London.
- Leenhouts, P. W. (in collaboration with C. Kalkman & H. J. Lam) (1956). Burseraceae. *Flora Malesiana*, Ser. I, 5: 209–296. National Herbarium of the Netherlands, Leiden.
- Leenhouts, P. W. (1962). Loganiaceae. *Flora Malesiana*, Ser. I, 6: 293–387 (*Fagraea* pp. 299–226). National Herbarium of the Netherlands, Leiden.
- Leeuwenberg, A. J. M. (2001). Series of revisions of Apocynaceae L. Key to Apocynaceae in continental Africa. *Wageningen Univ. Pap.* 2001 (1): 111–122, 126.
- Leeuwenberg, A. J. M. & Kupicha, F. K. (1985). Apocynaceae. *Flora Zambesiaca* 7 (2). Royal Botanic Gardens, Kew.
- Lehnert, M. (2006). The Cyatheaceae and Dicksoniaceae (Pteridophyta) of Bolivia. *Brittonia* 5: 229–244.
- Lewis, G., Schrire, B., Mackinder, B. & Lock, M. (eds) (2005). *Legumes of the World*. Royal Botanic Gardens, Kew.
- Li J.-Q. (1996). The origin and distribution of the family Fagaceae. *Acta Phytotax. Sin.* 34: 376–396.
- Li Ping-tao, Gilbert, M. G. & Stevens, W. D. (1995). Asclepiadaceae. *Flora of China* 16: 189–270.
- Litt, A. J. & Stevenson, D. W. (2003a). Floral development and morphology of Vochysiaceae. I. The structure of the gynoecium. *Amer. J. Bot.* 90: 1533–1547.
- Litt, A. J. & Stevenson, D. W. (2003b). Floral development and morphology of Vochysiaceae. II. The position of the single fertile stamen. *Amer. J. Bot.* 90: 1548–1559.
- Lock, J. M. (2003). Zingiberaceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Lohmann, L. G. (2006). Untangling the phylogeny of neotropical lianas (Bignonieae–Bignoniaceae). *Amer. J. Bot.* 93: 304–318.
- Lombardi, J. A. (2004). Vitaceae. In: N. Smith, S. A. Mori, A. Henderson, D. W. Stevenson & S. V. Heald (eds), *Flowering Plants of the Neotropics*, pp. 394–396. Princeton, Princeton University Press.
- Londoño Vega, A. C., Álvarez, E., Forero, E. & Morton, C. M. (1995). A new genus and species of Dipterocarpaceae from the Neotropics. I. Introduction, taxonomy, ecology, and distribution. *Brittonia* 47: 225–236.
- Louis, J. & Leonard, J. (1948). Olacaceae. In: *Flore du Congo Belge et du Ruanda-Urundi. Spermatophytes* I: 249–278. Institut National pour l'Etude Agronomique du Congo, Yangambi.
- Lucas, E. J., Harris, S. A., Mazine, F. F., Belsham, S. R., Nic Lughadha, E. M., Telford, A., Gasson, P. E. & Chase, M. W. (2007). Suprageneric phylogenetics of Myrtaceae, the generically richest tribe in Myrtaceae (Myrtales). *Taxon* 56: 1105–1128.
- Luteyn, J. L. (ed.) (1995). Ericaceae: part 2. The superior-ovariated genera (Monotropoideae, Pyroloideae, Rhododendroideae and Vaccinioideae p.p.). *Flora Neotropica*, Monograph 66. New York Botanical Garden, New York.
- Luteyn, J. L. & Pedraza, P. Online resource Neotropical Blueberries. <http://www.nybg.org/bsci/res/lut2/main.html>
- Lüttge, U. (ed.) (2007). Clusiaceae: A Woody Neotropical Genus of Remarkable Plasticity and Diversity. Springer, Berlin.
- Maas, P. J. M. (1977). Renalmia, Costoideae (additions). *Flora Neotropica*, Monograph 18. New York Botanical Garden, New York.
- Mabberley, D. J., Pannell, C. M. & Sing, A. M. (1995). Meliaceae. *Flora Malesiana*, Ser. I, 12: 1–407. National Herbarium of the Netherlands, Leiden.

- Macbride, J. F. (1956). *Flora of Peru: Ochnaceae*. Field Museum of Natural History-Botany 13 (3a/2): 686–697.
- Mayo, S. J. (1985). Araceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Mayo, S. J., Bogner, J. & Boyce, P. C. (1997). *The Genera of Araceae*. Royal Botanic Gardens, Kew.
- McDade, L. A., Daniel, T. F. & Kiel, C. A. (2008). Towards a comprehensive understanding of phylogenetic relationships among lineages of Acanthaceae s.l. (*Lamiales*). *Amer. J. Bot.* 95: 1136–1152.
- McMurtry, D., Grobler, L., Grobler, J. & Burns, S. (2008). *Field Guide to the Orchids of Northern South Africa and Swaziland*. Umdaus Press, Hatfield.
- Medan, D. & Schirarend, C. (2004). Rhamnaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 6: 320–338. Springer-Verlag, Berlin.
- Meerow, A. W. (1989). Systematics of the Amazon lilies, *Eucharis* and *Caliphruria* (Amaryllidaceae). *Ann. Missouri Bot. Gard.* 76: 136–220.
- Meerow, A. W. (1990). Amaryllidaceae. *Flora of Ecuador* 41. University of Gothenburg, Gothenburg.
- Meerow, A. W. & Snijman, D. A. (1998). Amaryllidaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 3: 70–78. Springer-Verlag, Berlin.
- Mez, C. (1902). Myrsinaceae. In: A. Engler (ed.), *Das Pflanzenreich* 9 (IV, 236): 1–437. Wilhelm Engelmann, Berlin.
- Middleton, D. J. (2007). Apocynaceae (subfamilies Rauvolfioideae and Apocynoideae). *Flora Malesiana*, Ser. I, 18. National Herbarium of the Netherlands, Leiden.
- Minkin, J. P. (1983). *A Comparative Pollen Morphology of the Orobanchaceae and Rhiananthoid Scrophulariaceae*. Miami University.
- Mori, S. A. & Prance, G. T. (1989). Lecythidaceae, the zygomorphic flowered New World Genera. *Flora Neotropica, Monograph* 21 (II). New York Botanical Garden, New York.
- Morillo, G. N. (1997). Asclepiadaceae. *Flora of the Venezuelan Guayana* 3: 129–177. Missouri Botanical Garden Press, St. Louis.
- Nadaf, A. & Zanan, R. (2012). Indian Pandanaceae — an Overview. Springer-Verlag, Berlin.
- N'Kounkou, J. S. (1993). Les Dioscoreaceae du Congo. *Fragm. Flor. Geobot. Suppl.* 2 (1): 139–182.
- Newman, M., Lhuillier, A. & Poulsen, A. D. (2004). Checklist of the Zingiberaceae of Malesia. *Blumea Suppl.* 16.
- Ng, F. S. P. (1989). Elaeocarpaceae. In: F. S. P. Ng (ed.), *Tree Flora of Malaya* 4: 82–97. Longman, Malaysia.
- Ng, F. S. P. (2004). Ebenaceae. In: E. Soepadmo, L. G. Saw & R. C. K. Chung (eds), *Tree Flora of Sabah and Sarawak*. Vol. 4: 29–100. Malaysian Nature Society, Kuala Lumpur.
- Nicholson, D. H. (1988). Araceae. In: M. D. Dassanayake & F. R. Fosberg (eds), *A Revised Handbook to the Flora of Ceylon* 6: 17–101. A. A. Balkema, Rotterdam.
- Nickrent, D. L., Malecot, V., Vidal-Russell, R. & Der, J. P. (2010). A revised classification of the Santalales. *Taxon* 59: 538–558.
- Nielsen, I. C. (1992). Mimosaceae (Leguminosae—Mimosoideae). *Flora Malesiana*, Ser. I, 11: 1–226. National Herbarium of the Netherlands, Leiden.
- Nixon, K. C. (1989). Origins of Fagaceae. In: P. R. Crane & S. Blackmore (eds), *Evolution, Systematics and the Fossil History of the Hamamelidae* 2: 23–43. Systematics Association Special Volume 40B.
- Nunes, T. (2009). Neotropical Passifloraceae. In: W. Milliken, B. Klitgaard & A. Baracat (eds), *Neotropikey — Interactive Key and Information Resources for Flowering Plants of the Neotropics*. www.kew.org/science/tropamerica/neotropikey/families/Passifloraceae.htm.
- Nyffler, R. & Baum, D. A. (2000). Phylogenetic relationships of the durians (Bombacaceae-Durioneae or /Malvaceae/Helicteroidae/Durioneae) based on chloroplast and nuclear ribosomal DNA sequences. *Plant Syst. Evol.* 224: 55–82.
- Omino, E. A. (2002). Apocynaceae. Part I. In: H. J. Beentje & S. A. Ghazanfar (eds), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Pacheco, L. (1995). Hymenophyllaceae. In: G. Davidse, S. Sousa, C. Mario & S. Knapp (eds), *Flora Mesoamericana* 1: 62–83. Universidad Nacional Autónoma de México, Instituto de Biología, México, D.F.
- Paiva, J. (2007). Polygalaceae. In: H. J. Beentje & S. Ghazanfar (eds), *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Parris, B. S. (1992). *The Plants of Mount Kinabalu I: Ferns and Fern Allies*. Royal Botanic Gardens, Kew.
- Paton, A. J., Bramley, G., Ryding, O., Polhill, R. M., Harvey, Y. B., Iwarson, M., Willis, F., Phillipson, P. B., Balkwill, K., Lukhoba, C. W., Otiend, D. F. & Harley, R. M. (2009). Lamiaceae (Labiatae). In: H. J. Beentje, S. A., Ghazanfar & R. M. Polhill (eds), *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- PBI Solanum Project. (2015). Solanaceae Source. www.solanaceaesource.org/.
- Pennington, T. D. (1991). *The Genera of Sapotaceae*. Royal Botanic Gardens, Kew.
- Pennington, T. D. (2007). 152. Sapotaceae. *Flora of Ecuador* 80. University of Gothenburg, Gothenburg.
- Pennington, T. D. & Styles, B. T. (1975). A generic monograph of the Meliaceae. *Blumea* 22: 419–540.
- Pennington, T. D., Styles, B. T. & Taylor, D. A. H. (1981). Meliaceae. *Flora Neotropica, Monograph* 28. New York Botanical Garden, New York.
- Peralta, I. E., Spooner, D. M. & Knapp, S. (2008). Taxonomy of wild tomatoes and their relatives (*Solanum* sections *Lycopersicoides*, *Juglandifolia*, *Lycopersicon*; Solanaceae). *Systematic Botany Monographs* 84: 1–186.
- Peterson, B. (1978). Thymelaeaceae. *Flora of Tropical East Africa*. Crown Agents, London.
- Philipson, W. R. (1979). Araliaceae. *Flora Malesiana*, Ser. I, 9: 1–105. National Herbarium of the Netherlands, Leiden.
- Phillips, O. & Miller, J. (2002). *Global Patterns of Plant Diversity, Alwyn H. Gentry's Forest Transect Data Set (MSB 89)*.
- Pipoly, J. J. (1998). The genus *Cybianthus* (Myrsinaceae) in Ecuador and Peru. *Sida* 18 (1): 1–160.
- Pires de Godoy, S. A. (2009). Neotropical Campanulaceae. In: W. Milliken, B. Klitgaard & A. Baracat (eds), *Neotropikey — Interactive Key and Information Resources for Flowering Plants of the Neotropics*. www.kew.org/science/tropamerica/neotropikey/families/Campanulaceae.htm.
- Polhill, R. M. (1966). Ulmaceae. *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Polhill, R. M. (1968). Ulmaceae. In: *Flore Cameroun* 8: 1–30.
- Polhill, R. M. & Wiens, D. (1998). Mistletoes of Africa. Royal Botanic Gardens, Kew.
- Polhill, R. M. & Wiens, D. (1999). Loranthaceae. In: H. J. Beentje (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Porter, D. M. (2005). Zygophyllaceae. *Flora of Ecuador* 75. University of Gothenburg, Gothenburg.
- Prance, G. T. (1972). Chrysobalanaceae. *Flora Neotropica, Monograph* 9. New York Botanical Garden, New York.
- Prance, G. T. (1989a). Chrysobalanaceae. *Flora Malesiana* 10: 635–678. National Herbarium of the Netherlands, Leiden.
- Prance, G. T. (1989b). Chrysobalanaceae: Supplement. *Flora Neotropica, Monograph* 9S. New York Botanical Garden, New York.
- Prance, G. T. (2001). Rhizophoraceae. *Flora of Nicaragua*. *Monogr. Syst. Bot. Missouri Bot. Gard.* 85 (3): 2200–2201.

- Prance, G.T. (2009). In: W. Milliken, B. Klitgaard & A. Baracat (eds), *Neotropikey – Interactive key and Information Resources for Flowering Plants of the Neotropics*. www.kew.org/neotropikey
- Prance, G.T. & Mori, S.A. (1979). Lecythidaceae, the actinomorphic flowered New World Genera. *Flora Neotropica, Monograph 21* (I). New York Botanical Garden, New York.
- Prance, G.T. & White, F. (1988). The genera of Chrysobalanaceae. *Phil. Trans. R. Soc. Lond. B* 320: 1–184.
- Prance, G.T., Beentje, H., Dransfield, J. & Johns, R.J. (2000). The tropical flora remains undercollected. *Ann. Missouri Bot. Gard.* 87: 67–71.
- Pridgeon, A.M., Cribb, P.J., Chase, M. & Rasmussen, F.N. (1999). *Genera Orchidacearum*. Volumes 1–6. Oxford University Press, Oxford.
- Prince, L.M. (2007). A brief nomenclatural review of genera and tribes in Theaceae. *Aldo 24*: 105–121.
- Prince, L.M. & Kress, W.J. (2006). Phylogenetic relationships and classification in Marantaceae: insights from plastid DNA sequence data. *Taxon* 55: 281–296.
- Radcliffe-Smith, A. (2001). *Genera Euphorbiacearum*. Royal Botanic Gardens, Kew.
- Rahmanzadeh, R., Müller, K., Fischer, E., Bartels, D. & T. Borsch, T. (2004). The Linderniaceae and Gratiolaceae are further lineages distinct from the Scrophulariaceae (Lamiales). *Plant Biol.* 7: 1–12.
- Rahn, K. & Rudall, P. (1998). Aliiaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants 3*: 70–78. Springer-Verlag, Berlin.
- Renner, S.S., Clausing, G., Cellinese, N. & Meyer, K. (2001). Melastomataceae. In: K. Larsen & I. Nielsen (eds), *Flora of Thailand* 7(3): 412–496.
- Ridsdale, C.E. (1974). A revision of the family Leeaceae. *Blumea* 22: 57–100.
- Robbrecht, E. (1988). Tropical woody Rubiaceae. *Opera Bot. Belg.* 1: 1–271.
- Rohwer, J.G. (1993). Lauraceae. In: K. Kubitzki, J.G. Rohwer & V. Bittrich (eds), *The Families and Genera of Vascular Plants 2*: 366–391. Springer-Verlag, Berlin.
- Sajo, M.G. & Rudall, P.J. (2002). Leaf and stem anatomy of Vochysiaceae in relation to subfamilial and suprafamilial systematics. *Bot. J. Linn. Soc.* 138: 339–364.
- Samain, M.S., Mathieu, G., Wanke, S., Neinhuis, C. & Goetghebeur, P. (2008). *Verhuellia* revisited — unravelling its intricate taxonomic history and a new subfamilial classification of Piperaceae. *Taxon* 57: 583–587.
- Sanchez, C. & Regaldo Gabancho, L. (2003). Aspleniaceae. In: W. Greuter, R. Rankin Rodriguez (eds), *Flora de la Republica de Cuba. Serie A. Fasciculo 8*. Ruggell, Gantner.
- Sanchez, M.N.S. (2010). Boraginaceae. In: W. Milliken, B. Klitgaard & A. Baracat (eds), *Neotropikey – interactive key and information resources for flowering plants of the Neotropics*. www.kew.org/neotropikey
- Sauer, J.D. (1967). The grain amaranths and their relatives: a revised taxonomic and geographic survey. *Ann. Missouri Bot. Gard.* 54: 103–137.
- Sauquet, H. (2004). Systematic revision of Myristicaceae (Magnoliales) in Madagascar, with four new species of *Mauloutchia*. *Bot. J. Linn. Soc.* 146 (3): 351–368.
- Schatz, G.E. (2001). Araliaceae, p. 59; Rutaceae, pp. 345–351; Sapindaceae, pp. 360–375; Simaroubaceae, pp. 387–389. In: G.E. Schatz, *Generic Tree Flora of Madagascar*. Royal Botanic Gardens, Kew.
- Scotland, R.W. & Vollesen, K. (2000). Classification of Acanthaceae. *Kew Bull.* 55: 513–589.
- Seidenfaden, G. & Wood, J.J. (1992). *The Orchids of Peninsular Malaysia and Singapore*. Olsen & Olsen, in association with The Royal Botanic Gardens Kew and Botanic Gardens Singapore.
- Sheahan, M.C. & Cutler, D.F. (1993). Contribution of vegetative anatomy to the systematics of Zygophyllaceae. *Bot. J. Linn. Soc.* 113: 227–262.
- Simmons, M.P. (2004). Celastraceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants 6*: 29–64. Springer-Verlag, Berlin.
- Simpson, D.A. & Inglis, C.A. (2001). Cyperaceae of economic, ethnobotanical and horticultural importance: a checklist. *Kew Bull.* 56: 257–360.
- Sinclair, J. (1955). The revision of the Malayan Annonaceae. *Gard. Bull. Straits Settlem.* 14: 149–516.
- Sleumer, H.O. (1955). Flacourtiaceae. *Flora Malesiana*, Ser. I, 5: 1–106. National Herbarium of the Netherlands, Leiden.
- Sleumer, H.O. (1966). Ericaceae. *Flora Malesiana*, Ser. I, 6: 469–914. National Herbarium of the Netherlands, Leiden.
- Sleumer, H.O. (1971a). Icacinaceae. *Flora Malesiana*, Ser. I, 7: 1–87. National Herbarium of the Netherlands, Leiden.
- Sleumer, H.O. (1975). Flacourtiaceae. In: R.M. Polhill (ed.), *Flora of Tropical East Africa*. Crown Agents, London.
- Sleumer, H.O. (1980a). Flacourtiaceae. *Flora Neotropica, Monograph 22*. New York Botanical Garden, New York.
- Sleumer, H.O. (1980b). A taxonomic account of the Olacaceae of Asia, Malesia, and the adjacent areas. *Blumea* 26: 145–168.
- Sleumer, H.O. (1984). Olacaceae. *Flora Neotropica, Monograph 38*. New York Botanical Garden, New York.
- Smith, A.R. (1983). Polypodiaceae-Thelypteridoideae. *Flora of Ecuador* 18. University of Gothenburg, Gothenburg.
- Smith, A.R. (1990). Thelypteridaceae. In: K. Kubitzki, J.G. Rohwer & V. Bittrich (eds), *The Families and Genera of Vascular Plants 1*: 263–272. Springer-Verlag, Berlin.
- Smith, A.R. (1992). Thelypteridaceae. In: R.M. Tryon & R.G. Stolze (eds), *Pteridophyta of Peru. Part III. Fieldiana Bot. n.s.* 29: 1–80.
- Smith, C.E. (1954). The New World species of Sloanea (Elaeocarpaceae). *Contr. Gray Herb.* 175: 1–114.
- Smith, L.B. & Downs, R.J. (1974–1979). Bromeliaceae. *Flora Neotropica* 14. Hafner Press, New York.
- Smith, L.B. & Till, W. (1998). Bromeliaceae. In: K. Kubitzki, J.G. Rohwer & V. Bittrich (eds), *The Families and Genera of Vascular Plants 4*: 74–99. Springer-Verlag, Berlin.
- Smith, S. & Stansbie, J. (2003). Aliiaceae. In: H.J. Beentje (ed.) *Flora of Tropical East Africa*. A.A. Balkema, Rotterdam.
- Smitinand, T., Santisuk, T. & Phengklai, C. (1980). The manual of Dipterocarpaceae of mainland South East Asia. *Thai Forest Bull.* 12: 1–133.
- Sobral, M., Proença, C., Souza, M., Mazine, F. & Lucas, E. (2010). Myrtaceae. In: R.C. Forzza et al. (eds), *Catálogo de Plantas e Fungos do Brasil 2. Jardim Botânico do Rio de Janeiro*.
- Soepadmo, E. (1972). Fagaceae. *Flora Malesiana*, Ser. I, 7: 265–403. National Herbarium of the Netherlands, Leiden.
- Soepadmo, E. (1977). Ulmaceae. *Flora Malesiana*, Ser. I, 8: 31–76. National Herbarium of the Netherlands, Leiden.
- Spirlet, M.L. (1966). Guttiferae. *Contribution à la Flore du Congo, du Rwanda et du Burundi*.
- Stace, C.A. (2010). Combretaceae. In: *Flora Neotropica* 107, 1–374. New York Botanical Garden Press, New York.
- Ståhl, B. & Anderberg, A.A. (2004). Myrsinaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants 6*: 266–281. Springer-Verlag, Berlin.
- Stannard, B. (2000). Simaroubaceae. In: H.J. Beentje (ed.) *Flora of Tropical East Africa*. A.A. Balkema, Rotterdam.

- Staples, G.W. & Brummitt, R. K. (2007). Convolvulaceae. In: V. H. Heywood, R. K. Brummitt, A. Culham & O. Seberg (eds), *Flowering Plant Families of the World*, pp. 108–110. Royal Botanic Gardens, Kew.
- Stedje, B. (1997). Hyacinthaceae. In: R. Polhill (ed.), *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Stevens, P. F. (1980). A revision of the Old World species of *Calophyllum* (Guttiferae). *J. Arnold Arbor.* 61 (3): 425–699.
- Stevens, P. F. (2001 onwards). Angiosperm Phylogeny Website. Version 9, June 2008 [and more or less continuously updated since]. www.mobot.org/MOBOT/research/APweb/.
- Stevens, P. F. (2006). Clusiaceae—Guttiferae, pp. 48–66; Hypericaceae, pp. 194–201. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 9. Springer-Verlag, Berlin.
- Stevens, P. F., Dressler, S. & Weitzman, A. L. (2004a). Theaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 6: 463–471. Springer-Verlag, Berlin.
- Stevens, P. F., Lutley, J., Oliver, E. G. H., Bell, T. L., Brown, E. A., Crowdus, R. K., George, A. S., Jordan, G. J., Ladd, P., Lemson, K., McLean, C. B., Menadue, Y., Pate, J. S., Stace, H. M. & Weiller, C. M. (2004b). Ericaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 6: 145–194. Springer-Verlag, Berlin.
- St. John, H. (1960). Revision of the genus *Pandanus* Stickman: part I. Key to the sections. *Pacific Science* 14: 224–241.
- Stolze, R. G. (1981). Ferns and fern allies of Guatemala. Part II. Polypodiaceae. *Fieldiana: Bot.*, n.s. 6.
- Stone, B. C. (1970). Malayan climbing Pandans — the genus *Freycinetia* in Malaya. *Malay. Nat. J.* 23: 84–91.
- Stone, B. C. (1989). Myrsinaceae. In: F. S. P. Ng (ed.), *Tree Flora of Malaya* 4: 264–284. Longman, Malaysia.
- Sruwé, L. & Albert, V. A. (2002). *Gentianaceae — Systematics and Natural History*. Cambridge University Press, Cambridge.
- Sruwé, L., Maas, J. P., Pihlar, O. & Albert, V. A. (1999). Gentianaceae. In: P. E. Berry, K. Yatskievych & B. K. Holst (eds), *Flora of the Venezuelan Guayana*, 5: 474–542. Missouri Botanical Garden, St. Louis.
- Styles, B. T. & White, F. (1991). Meliaceae. In R. M. Polhill (ed.), *Flora of Tropical East Africa*, A. A. Balkema, Rotterdam.
- Suessenguth, K. (1953). Rhamnaceae. In: A. Engler & K. Prantl, *Die natürlichen Pflanzenfamilien* 20d: 7–173. Duncker & Humblot, Berlin.
- Suksathan, P., Gustafsson, M. H. & Borchsenius, F. (2009). Phylogeny and generic delimitation of Asian Marantaceae. *Bot. J. Linn. Soc.* 159: 381–395.
- Tagawa, M. & Iwatsuki, K. (1985). Aspleniaceae. In: T. Smitinand & K. Larsen (eds), *Flora of Thailand* 3 (2): 261–291. Phonphan Printing Company, Bangkok.
- Taylor, P. (1963). Gentianaceae. In: F. N. Hepper (ed.), *Flora of West Tropical Africa*, vol. I. Crown Agents, London.
- Taton, A. (1980). Contribution à l'étude des genres *Embelia*, *Maesa* et *Rapanea* (Myrsinaceae) en Afrique tropicale. *Bull. Jard. Bot. Nation. Belg.* 50 (1/2): 201–226.
- Tellez V. O. & B.G. Schubert (1994). Dioscoreaceae. In: G. Davidse, M. Sousa & A. O. Chater (eds), *Flora Mesoamericana* 6, pp. 54–65. UNAM, Mexico.
- Thulin, M. (1976). Campanulaceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. Crown Agents, London.
- Thulin, M. (1984). Lobeliaceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Townsend, C. (1985). Amaranthaceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Townsend, C. C. (1993). Amaranthaceae. In: K. Kubitzki (ed.), *The Families and Genera of Flowering Plants* 2: 70–91. Springer-Verlag, Berlin.
- Troupin, G. (1962). Monographie des Menispermaceae Africaines. *Mém. Acad. Roy. Sci. Outre-Mer, Cl. Sci. Nat. Méd., Collect. 8vo, Nouvelle Série* 13: 1–313.
- Tryon, R. M., Stolze, R. G. & Leon, B. (1993). Pteridophyta of Peru. Part V. 18: Aspleniaceae – 21: Polypodiaceae. *Fieldiana, Bot.* n.s. 32.
- Utley, J. F. & Burt-Utley, K. (1994). Bromeliaceae. *Flora Mesoamericana* 6: 89–156.
- van Balgooy, M. M. J. (1998). *Malesian Seed Plants Vol 2 — Portraits of Tree Families*. p. 25–27.
- van der Meijden, R. (1988). Polygalaceae. *Flora Malesiana*, Ser. I, 10: 455–539.
- van der Werff, H. & Richter, H. G. (1996). Toward an improved classification of Lauraceae. *Ann. Missouri Bot. Gard.* 83: 409–418.
- van der Werff, H. (1991). A key to the genera of Lauraceae in the New World. *Ann. Missouri Bot. Gard.* 78: 377–387.
- van der Werff, H. (2001). An annotated key to the genera of Lauraceae in the Flora Malesiana region. *Blumea* 46: 125–140.
- Van Heusden, E. C. H. (1992). Flowers of Annonaceae: morphology, classification, and evolution. *Blumea suppl.* 7: 1–218.
- van Ooststroom, S. J. (1953). Convolvulaceae. *Flora Malesiana*, Ser. I, 4: 388–512; 5: 558–564. National Herbarium of the Netherlands, Leiden.
- van Welzen, P. C. (ed.) (ongoing). *Flora Malesiana Euphorbiaceae*. Nationaal Herbarium Nederland, Leiden. www.nationalherbarium.nl/euphorbs
- Verdcourt, B. (1963). Convolvulaceae. In: C. E. Hubbard & E. Milne-Redhead (eds), *Flora of Tropical East Africa*. Crown Agents, London.
- Verdcourt, B. (1971). Annonaceae. In: E. Milne-Redhead & R. M. Polhill (eds), *Flora of Tropical East Africa*. Crown Agents, London.
- Verdcourt, B. (1979). A manual of New Guinea legumes. *Office of Forests, Division of Botany, Lae, Papua New Guinea, Botany Bulletin* 11: 1–645.
- Verdcourt, B. (1986). Aristolochiaceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Verdcourt, B. (1989). Dipterocarpaceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Verdcourt, B. (1991). Boraginaceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. CRC Press, London.
- Verdcourt, B. (1993). Vitaceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Verdcourt, B. (1994). Lythraceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Verdcourt, B. (1996). Lauraceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Verdcourt, B. (1997). Myristicaceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Verdcourt, B. (2001). Polypodiaceae. In: R. M. Polhill (ed.), *Flora of Tropical East Africa*. A. A. Balkema, Rotterdam.
- Verdcourt, B. (2005). Ochnaceae. In: H. J. Beentje & S. A. Ghazanfar (eds), *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Verdcourt, B. (2006). Thelypteridaceae. In: H. J. Beentje & S. A. Ghazanfar (eds), *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.
- Vidal-Russell, R. & Nickrent, D. L. (2008). Evolutionary relationships in the showy mistletoe family (Loranthaceae). *Amer. J. Bot.* 95: 1015–1029.
- Villiers J.-F. (1973b). Icacinacees. *Flore du Cameroun* 15: 3–100. Muséum National d'Histoire Naturelle, Paris.
- Villiers, J.-F. (1973a). Olacaceae. In: *Flore du Cameroun* 15: 101–162. Muséum National d'Histoire Naturelle, Paris.
- Vollesen, K. (2008). Acanthaceae (Part I). In: H. J. Beentje & S. A. Ghazanfar (eds), *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.

- Wallander, E. (2014). Oleaceae information site. www.oleaceae.info/
- Wallander, E. & Albert, V.A. (2000). A phylogeny and classification of Oleaceae based on rps16 and trnL-trnF sequence data. *Amer.J. Bot.* 87: 1827–1841.
- Wallnöfer, B. (2001). The biology and systematics of Ebenaceae: a review. *Ann. Naturhist. Mus. Wien* 103B: 485–512.
- Wallnöfer, B. (2007). A revision of neotropical *Diospyros* (Ebenaceae) : part I. *Ann. Naturhist. Mus. Wien*. 108B: 207–247.
- Wanderley, M. G. L. & Martins, S. E. (2007). Bromeliaceae. In: M. G. L. Wanderley, T. S. Melhem, S. E. Martins, S. L. Jung-Mendaçolli, G. J. Shepherd & M. Kirizawa (eds), *Flora Fanerogâmica do Estado de São Paulo* 5: 1–476. Instituto de Botânica, São Paulo.
- Watson, L. & Dallwitz, M. J. (1992 onwards). *The Families of Flowering Plants: Descriptions, Illustrations, Identification, and Information Retrieval*. <http://delta-intkey.com>
- Weber, A. (2004). Gesneriaceae. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 7: 63–158. Springer-Verlag, Berlin.
- Weber, A. & Skog, L. E. (2007 onwards). *The Genera of Gesneriaceae. Basic Information with Illustration of Selected Species*. 2nd edition. <http://www.genera-gesneriaceae.at>
- Weddell, H.A. (1869). Urticaceae. In: A. De Candolle, *Prodromus Systematis Naturalis Regni Vegetabilis*, vol. 16, I. Paris.
- Wen, J. (2006). Leeaceae, pp. 221–225; Vitaceae, pp. 467–479. In: K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 9. Springer-Verlag, Berlin.
- White, F. & Verdcourt, B. (1996). Ebenaceae. *Flora of Tropical East Africa*. A.A. Balkema, Rotterdam.
- Whitlock, B.A., Bayer, C. & Baum, D.A. (2001). Phylogenetic relationships and floral evolution of the Byttnerioideae (Sterculiaceae or Malvaceae s.l.) based on sequences of the chloroplast gene, ndhF. *Syst. Bot.* 26: 420–437.
- Wickens, G. E. (1973). Combretaceae. *Flora of Tropical East Africa*. Crown Agents, London.
- Wickens, G. E. (1975). Melastomataceae. *Flora of Tropical East Africa*. Crown Agents, London.
- Wild, H. & Drummond, R. B. (1963). Vitaceae. In: A. W. Exell, A. Fernandes & H. Wild (eds), *Flora Zambesiaca* 2 (2). Crown Agents, London.
- Wilkie, P., Clark, A., Pennington, R. T., Cheek, M., Bayer, C. & Wilcock, C. C. (2006). Phylogenetic relationships within the subfamily Sterculioideae (Malvaceae/Sterculiaceae-Sterculieae) using the chloroplast gene ndhF. *Syst. Bot.* 31: 160–170.
- Wilkin, P. (2009). Dioscoreaceae. In: J. Timberlake (ed.), *Flora Zambesiaca* 12 (2): 109–140. Royal Botanic Gardens, Kew.
- Wilkin, P. & Thapayai, C. (2009). Dioscoreaceae. In: T. Santisuk & K. Larsen (eds), *Flora of Thailand* 10 (1): 1–140. The Forest Herbarium, National Park, Wildlife and Plant Conservation Department, Bangkok.
- Wilmot-Dear, C. M. (1991). Ulmaceae and Cannabaceae. In: Pope G.V. (ed.), *Flora Zambesiaca* 9 (6): 1–13. Royal Botanic Gardens, Kew.
- Wilmot-Dear, C. M. & Friis, I. (1996). The New World species of Boehmeria & Pouzolzia (Urticaceae, tribus Boehmerieae). *Opera Bot.* 129: 1–103.
- Wilson, P. G. (2011). Myrtaceae. In K. Kubitzki (ed.), *The Families and Genera of Vascular Plants* 10: 212–271. Springer-Verlag, Berlin.
- Wu, D.-L. & Larsen, K. (2000). Zingiberaceae. In: Z. Wu & P. H. Raven (eds), *Flora of China* 24: 322–381. Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
- Wurdack, J. J., Morley, T. & Renner, S. (1993). *Flora of the Guianas, series A: Phanerogams*. 99: Melastomataceae. Koeltz Scientific Books, Konigstein.
- Wurdack, K. J., Hoffmann, P. & Chase, M. W. (2005). Molecular phylogenetic analysis of uniovulate Euphorbiaceae (Euphorbiaceae *sensu stricto*) using plastid rbcL and trnL-F DNA sequences. *Amer.J. Bot.* 92: 1397–1420.
- Yunker, T. G. (1972). The Piperaceae of Brazil I – *Piper*: Group I, II, III, IV. *Hoehnea* 2: 19–366.
- Yunker, T. G. (1974). The Piperaceae of Brazil III – *Peperomia*; taxa of uncertain status. *Hoehnea* 4: 71–413.
- Zappi, D. (2009). Neotropical Loganiaceae. In: W. Milliken, B. Klitgaard & A. Baracat (eds), *Neotropikey — Interactive key and Information Resources for Flowering Plants of the Neotropics*. www.kew.org/science/tropamerica/neotropikey/families/Loganiaceae.htm.
- Zhang, M. & Tucker, G. C. (2008). Capparaceae. In: Z. Y. Wu, P. H. Raven & D. Y. Hong (eds), *Flora of China* 7: 433–450. Science Press, Beijing; Missouri Botanical Garden Press, St. Louis.