

References

- American Psychological Association (2012). *Stress in America: Our health at risk*. Washington, DC: Author.
- Centers for Disease Control and Prevention. Falls Among Older Adults. <http://www.cdc.gov/HomeandRecreationalSafety/Falls/adultfalls.html>
- Chen, K.W. (2004) An analytic review of studies measuring effects of external QI in China. *Altern Ther Health Med*, 10(4), 38-50.
- Hong, Y. (Ed.). (2008). *Tai chi chuan: State of the art in international research*. New York: Karger.
- Martin, Brook I.; Deyo, Richard A.; Mirza, Sohail K; Turner, Judith A.; Comstock, Bryan A.; Hollingworth, William; Sullivan, Sean D. (2008). Expenditures and health status among adults with back and neck problems. *JAMA*, 299(6), 656-664.
- Mayo Clinic. Exercise and stress: Get moving to combat stress. <http://www.mayoclinic.com/health/exercise-and-stress/SR00036>.
- Salthous, T.A. (1984). Effects of age and skill in typing. *Journal of Experimental Psychology: General*, 113(3), 345-371.
- Schmitz, T. J. (2007). Examination of Sensory Function. In S. B. O'Sullivan & T.J. Schmitz. *Physical Rehabilitation* (5th ed.). Philadelphia, PA: F. A. Davis Company, 121-157.
- Stux, G., & Hammerschlag, R. (2001). *Clinical acupuncture: Scientific basis*. New York: Springer.
- Wolf, S.L., Barnhart, H.X., Kutner, N.G., McNeely, E., Coogler, C., & Xu, T. (1996). Reducing frailty and falls in older persons: An investigation of t'ai chi and computerized balance training. *Journal of the American Geriatrics Society*, 44(5), 599-600.
- Zhu, W., Guan, S., & Yang, Y. (2010). Clinical implication of tai chi interventions: A review. *American Journal of Lifestyle Medicine*, 4 (Sept./Oct.), 418-432.