

References

Chapter 1

1. Bouchard C (ed.). *Physical activity and obesity*. Human Kinetics: Champaign, IL, 2000.
2. World Health Organization. *The global burden of disease: 2004 update*. World Health Organization: Geneva, 2008.
3. World Health Organization. *The world health report 2002: reducing risks, promoting healthy life*. World Health Organization: Geneva, 2002.
4. Bouchard C. The biological predisposition to obesity: beyond the thrifty gene scenario [commentary]. *Int J Obes* 2007; 31: 1337-1339.
5. O'Rahilly S, Farooqi IS. Human obesity: a heritable neurobehavioral disorder that is highly sensitive to environmental conditions. *Diabetes* 2008; 57: 2905-2910.
6. Bouchard C, Agurs-Collins T (eds.). Gene-nutrition and gene-physical activity interactions in the etiology of obesity. *Obesity* 2008; 16 suppl. 3: S1-96.
7. Frayling TM, Timpson NJ, Weedon MN, Zeggini E, Freathy RM, Lindgren CM, et al. A common variant in the *FTO* gene is associated with body mass index and predisposes to childhood and adult obesity. *Science* 2007; 316: 889-894.
8. Andreasen CH, Stender-Petersen KL, Mogensen MS, Torekov SS, Wegner L, Andersen G, et al. Low physical activity accentuates the effect of the *FTO* rs9939609 polymorphism on body fat accumulation. *Diabetes* 2008; 57: 95-101.
9. Rumpersaud E, Mitchell BD, Pollin TI, Fu M, Shen H, O'Connell JR, et al. Physical activity and the association of common *FTO* gene variants with body mass index and obesity. *Arch Intern Med* 2008; 168: 1791-1797.
10. Bray GA, Bouchard C (eds.). *Handbook of obesity: clinical applications*. 3rd ed. Informa Healthcare: New York, 2008.

Chapter 2

1. Warburton DER, Katzmarzyk PT, Rhodes RE, Shephard RJ. Evidence-informed physical activity guidelines for Canadian adults. *Appl Physiol Nutr Metab* 2007; 32: S16-S68.
2. Warburton DE, Nicol C, Bredin SS. Health benefits of physical activity: the evidence. *CMAJ* 2006; 174: 801-809.
3. Katzmarzyk PT, Janssen I. The economic costs associated with physical inactivity and obesity in Canada: an update. *Can J Appl Physiol* 2004; 29: 90-115.
4. Bouchard C, Shephard RJ. Physical activity fitness and health: the model and key concepts. In: Bouchard C, Shephard RJ, Stephens T (eds.), *Physical activity, fitness, and health: international proceedings and consensus statement*. Human Kinetics: Champaign, IL, 1994, pp. 77-88.
5. Ross R, Janssen I. Physical activity, fitness, and obesity. In: Bouchard C, Blair SN, Haskell WL (eds.), *Physical activity and health*. Human Kinetics: Champaign, IL, 2007, pp. 173-190.
6. Saris WH, Blair SN, van Baak MA, Eaton SB, Davies PS, Di Pietro L, et al. How much physical activity is enough to prevent unhealthy weight gain? Outcome of the IASO 1st

Stock Conference and consensus statement. *Obes Rev* 2003; 4: 101-114.

7. Esliger D, Tremblay MS. Physical activity and inactivity profiling: the next generation. *Appl Physiol Nutr Metab* 2007; 98: S195-207.
8. Warburton DE, Nicol C, Bredin SS. Prescribing exercise as preventive therapy. *CMAJ* 2006; 174: 961-974.
9. Tudor-Locke CE, Myers AM. Challenges and opportunities for measuring physical activity in sedentary adults. *Sports Med* 2001; 31: 91-100.
10. Hamilton MT, Hamilton DG, Zderic TW. Role of low energy expenditure and sitting in obesity, metabolic syndrome, type 2 diabetes, and cardiovascular disease. *Diabetes* 2007; 56: 2655-2667.

Chapter 3

1. Katzmarzyk PT, Baur LA, Blair SN, Lambert EV, Oppert J-M, Riddoch C. International Conference on Physical Activity and Obesity in Children: summary statement and recommendations. *Appl Physiol Nutr Metab* 2008; 33: 371-388.
2. Eisenmann JC, Barte RT, Smith DT, Welk GJ, Fu Q. Combined influence of physical activity and television viewing on the risk of overweight in US youth. *Med Sci Sports Exerc* 2008; 32: 613-618.
3. Hamilton MT, Hamilton DG, Zderic TW. Role of low energy expenditure and sitting in obesity, metabolic syndrome, type 2 diabetes, and cardiovascular disease. *Diabetes* 2007; 56: 2655-2667.
4. Healy GN, Dunstan DW, Salmon J, Shaw JE, Zimmet PZ, Owen N. Television time and continuous metabolic risk in physically active adults. *Med Sci Sports Exerc* 2008; 40: 639-645.
5. Rosenberg DE, Bull FC, Marshall AL, Sallis JF, Bauman AE. Assessment of sedentary behavior with the International Physical Activity Questionnaire. *J Phys Act Health* 2008; 5(suppl. 1): S30-S44.
6. Tremblay MS, Esliger DW, Tremblay A, Colley R. Incidental movement, lifestyle-embedded activity and sleep: new frontiers in physical activity assessment. *Appl Physiol Nutr Metab* 2007; 32(suppl. 2E): S208-S217.
7. Shields M, Tremblay MS. Sedentary behavior and obesity among Canadian adults. *Health Rep* 2008; 19: 19-30.
8. Shields M, Tremblay MS. Screen time among Canadian adults: a profile. *Health Rep* 2008; 19: 31-43.
9. Canadian Paediatric Society. Impact of media use on children and youth. *Paediatr Child Health* 2003; 8: 301-306.
10. Tudor-Locke C, Bassett DR Jr. How many steps/day are enough? Preliminary pedometer indices for public health. *Sports Med* 2004; 34: 1-8.
11. Tudor-Locke C, Hatano Y, Pangrazi RP, Kang M. Revisiting "how many steps are enough?" *Med Sci Sports Exerc* 2008; 40(suppl. 7): S537-S543.

12. Reilly JJ, Penpraze V, Hislop J, Davies G, Grant S, Paton JY. Objective measure of physical activity and sedentary behavior: review with new data. *Arch Dis Child* 2008; 93: 614-619.
13. Esliger DW, Tremblay MS. Physical activity and inactivity profiling: the next generation. *Appl Physiol Nutr Metab* 2007; 32(suppl. 2E): S195-S207.

Chapter 4

1. U.S. Department of Health and Human Services. *Physical activity and health: a report of the surgeon general*. U.S. Department of Health and Human Services, Public Health Service, CDC, National Center for Chronic Disease Prevention and Health Promotion: Atlanta, 1996.
2. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 2007; 39: 1423-1434.
3. LaMonte MJ, Ainsworth BE. Quantifying energy expenditure and physical activity in the context of dose-response. *Med Sci Sports Exerc* 2001; 33: S370-S378.
4. Physical Activity Guidelines Advisory Committee. *Physical Activity Guidelines Advisory Committee Report*, 2008. Washington, DC: U.S. Department of Health and Human Services, 2008.
5. McKenzie TL. Use of direct observation to assess physical activity. In: Welk GK (ed.), *Physical activity assessments for health-related research*. Human Kinetics: Champaign, IL, 2002, pp. 179-196.
6. Ward DS, Evenson KR, Vaughn A, Rodgers AB, Troiano RP. Accelerometer use in physical activity: best practices and research recommendations. *Med Sci Sports Exerc* 2005; 37: S582-S588.
7. Tudor-Locke C, Bassett DR. How many steps/day are enough? Preliminary pedometer indices for public health. *Sports Med* 2004; 34: 1-8.
8. Ainsworth BE, Haskell WL, Whitt MC, Irwin ML, Swartz AM, Strath SJ. Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc* 2000; 32: S498-S516.
9. Pettee KK, Tudor-Locke C, Ainsworth BE. The measurement of energy expenditure and physical activity. In: Wolinsky I, Driskell JA (eds.), *Sports nutrition*. CRC Press: Boca Raton, FL, 2008, pp. 159-189.
10. Rodriguez DA, Brown AL, Troped PJ. Portable global positioning units to complement accelerometry-based physical activity monitors. *Med Sci Sports Exerc* 2005 Nov; 37(11 suppl.): S572-581.

Chapter 5

1. Dishman RK, Heath GW, Washburn RA. *Physical activity epidemiology*. Human Kinetics: Champaign, IL, 2004.
2. Brown WH, Pfeiffer KA, McIver KL, Dowda M, Almeida MJCA, Pate RR. Assessing preschool children's physical activity: an Observational System for Recording Physical Activity in Children—Preschool Version (OSRAC-P). *Res Q Exerc Sport* 2006; 77: 167-176.
3. Sallis JF. Self-report measures of children's physical activity. *J Sch Health* 1991; 61: 215-219.
4. Dorminy CA, Choi L, Akohoue SA, Chen KY, Buchowski MS. Validity of a multisensor armband in estimating 24-h energy expenditure in children. *Med Sci Sports Exerc* 2008 40: 699-706.

5. Zhang K, Werner P, Sun M, Pi-Sunyer FX, Boozer CN. Measurement of human daily physical activity. *Obes Res* 2003; 11: 33-40.
6. Zhang K, Pi-Sunyer FX, Boozer CN. Improving energy expenditure estimation for physical activity. *Med Sci Sports Exerc* 2004; 36: 883-889.
7. Troped PJ, Oliveira MS, Matthews CE, Cromley EK, Melly SJ, Craig BA. Prediction of activity mode with global positioning system and accelerometer data. *Med Sci Sports Exerc* 2008; 40: 972-978.

Chapter 6

1. Screening and Interventions to Prevent Obesity in Adults, topic page. U.S. Preventive Services Task Force. Agency for Healthcare Research and Quality, Rockville, MD. Dec 2003. www.ahrq.gov/clinic/uspstf/uspsobes.htm. Accessed 26 May 2008.
2. NHLBI Obesity Education Initiative Expert Panel on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults—the evidence report. *Obes Res* 1998; 6: 51S-63S.
3. World Health Organization. *Obesity: preventing and managing the global epidemic*. World Health Organization: Geneva, 1998.
4. Bray GA, Bouchard C, James WP. Definitions and proposed current classification of obesity. In: Bray GA, Bouchard C, James WP (eds.), *Handbook of obesity*. Marcel Dekker: New York, 1997, pp. 31-40.
5. Bray GA. Classification and evaluation of the overweight patient. In: Bray GA, Bouchard C, James WP (eds.), *Handbook of obesity*. Marcel Dekker: New York, 1997, pp. 831-854.
6. U.S. Department of Health and Human Services. *Clinician's handbook of preventive services: put prevention into family practice*. U.S. Government Printing Office: Washington, DC, 1994, pp. 141-146.

Chapter 7

1. Freedman DS, Wang J, Maynard LM, Thornton JC, Mei Z, Pierson RN, et al. Relation of BMI to fat and fat-free mass among children and adolescents. *Int J Obes* 2005; 29: 1-8.
2. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, Flegal KM, Guo SS, Wei R, et al. CDC growth charts: United States. *Adv Data* 2000; 314: 1-27.
3. Krebs NF, Himes JH, Jacobson D, Nicklas TA, Guilday P, Styne D. Assessment of child and adolescent overweight and obesity. *Pediatrics* 2007; 120(suppl. 4): S193-S228.
4. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity world wide: international survey. *Br Med J* 2000; 320: 1240-1243.
5. de Onis M, Garza C, Onyango AW, Martorell R. WHO child growth standards. *Acta Paediatr* 2006; 95(suppl. 450): 1-101.
6. Himes JH, Park K, Styne D. Menarche and assessment of body mass index in adolescent girls. *Journal of Pediatrics* 2009; 155: 393-397.
7. Himes JH. Agreement among anthropometric indicators identifying the fattest adolescents. *Int J Obes* 1999; 23(suppl. 2): S18-S21.
8. Heymsfield SB, Lohman TG, Wang ZM, Going SB (eds.). *Human body composition*. 2nd ed. Human Kinetics: Champaign, IL, 2005.
9. McCarthy HD, Cole TJ, Fry T, Jebb SA, Prentice AM. Body fat reference curves for children. *Int J Obes* 2006; 30: 598-602.

10. Jartti L, Hakanen M, Paakkunainen U, Raittinen P, Rönnemaa T. Comparison of hand-to-leg and leg-to-leg bioelectric impedance devices in the assessment of body adiposity in prepubertal children. The STRIP study. *Acta Paediatr* 2000; 89: 781-786.

Chapter 8

1. Connor Gorber S, Tremblay M, Moher D, Gorber B. A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obes Rev* 2007; 8: 307-326.
2. Behavioral Risk Factor Surveillance System. Nationwide (states and DC) prevalence data for 2004 – overweight and obesity. <http://apps.nccd.cdc.gov/brfss/sex.asp?cat=OB&yr=2004&qkey=4409&state=UB>. Accessed 30 April 2008.
3. Lethbridge-Çejku M, Rose D, Vickerie J. Summary health statistics for U.S. adults: National Health Interview Survey, 2004. *Vital Health Stat* 10(228). National Center for Health Statistics: Hyattsville, MD, 2006, p. 81.
4. Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA* 2006; 295: 1549-1555.
5. Sallis JF, Saelens BE. Assessment of physical activity by self-report: status, limitations, and future directions. *Res Q Exerc Sport* 2000; 71: 1-14.
6. Troiano RP, Berrigan D, Dodd KW, Mâsse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc* 2008; 40: 181-188.
7. Hagströmer M, Oja P, Sjöström M. Physical activity and inactivity in an adult population assessed by accelerometry. *Med Sci Sports Exerc* 2007; 39: 1502-1508.
8. Plankey MW, Stevens J, Flegal KM, Rust PF. Prediction equations do not eliminate systematic error in self-reported body mass index. *Obes Res* 1997; 5: 308-314.
9. Troiano RP, Dodd KW. Differences between objective and self-report measures of physical activity. What do they mean? *J Kor Soc Meas Eval* 2008; 10(2): 31-42.
8. Bassett D, Schneider PL, Huntington GE. Physical activity in an old order Amish community. *Med Sci Sports Exerc* 2004; 36: 79-85.
9. Bull F, Armstrong T, Dixon T, Ham S, Neiman A, Pratt M. Physical inactivity. In: Ezzati M, Lopez A, Rodgers A, Murray C (eds.), *Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors*. World Health Organization: Geneva, 2004, pp. 729-881.
10. Centers for Disease Control and Prevention. Prevalence of regular physical activity among adults—United States, 2001 and 2005. *MMWR* 2007; 56: 1209-1212.
11. Bryan SN, Tremblay MS, Pérez CE, Ardern CI, Katzmarzyk PT. Physical activity and ethnicity: evidence from the Canadian Community Health Survey. *Can J Pub Health* 2006; 97: 271-276.
12. Ross J, Hamlin M. Maori physical activity: a review of an indigenous population's participation. *Health Promot J Aust* 2007; 18: 73-76.

Chapter 10

1. Katzmarzyk PT, Baur LA, Blair SN, Lambert EV, Oppert JM, Riddoch C. International conference on physical activity and obesity in children: summary statement and recommendations. *Appl Physiol Nutr Metab* 2008; 33: 371-388.
2. Wareham NJ, Rennie KL. The assessment of physical activity in individuals and populations: why try to be more precise about how physical activity is assessed? *Int J Obes Relat Metab Disord* 1998; 22: S30-S38.
3. Ness AR, Leary SD, Mattocks C, Blair SN, Reilly JJ, Wells J, et al. Objectively measured physical activity and fat mass in a large cohort of children. *PLoS Med* 2007; 4: e97.
4. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc* 2008; 40: 181-188.
5. Riddoch CJ, Bo Andersen L, Wedderkopp N, Harro M, Klasson-Heggebo L, Sardinha LB, et al. Physical activity levels and patterns of 9- and 15-yr-old European children. *Med Sci Sports Exerc* 2004; 36: 86-92.
6. Riddoch CJ, Mattocks C, Deere K, Saunders J, Kirkby J, Tilling K, et al. Objective measurement of levels and patterns of physical activity. *Arch Dis Child* 2007; 92: 963-969.
7. Webber LS, Catellier DJ, Lytle LA, Murray DM, Pratt CA, Young DR, et al. Promoting physical activity in middle school girls: Trial of Activity for Adolescent Girls. *Am J Prev Med* 2008; 34: 173-184.
8. Andersen LB, Harro M, Sardinha LB, Froberg K, Ekelund U, Brage S, et al. Physical activity and clustered cardiovascular risk in children: a cross-sectional study (The European Youth Heart Study). *Lancet* 2006; 368: 299-304.
9. Pate RR, Stevens J, Pratt C, Sallis JF, Schmitz KH, Webber LS, et al. Objectively measured physical activity in sixth-grade girls. *Arch Pediatr Adolesc Med* 2006; 160: 1262-1268.

Chapter 11

1. WHO expert consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 2004; 363: 157-163.
2. James WPT, Jackson-Leach R, Ni Mhurchu C, Kalmara E, Shayeghi M, Rigby NJ, et al. Overweight and obesity (high body mass index). In: Ezzati M, Lopez AD, Rodgers A, Murray CJL (eds.), *Comparative quantification of health risks. Global and regional burden of disease attributable to selected*

Chapter 9

1. Brown W, Bauman A, Chey T, Trost S, Mummery K. Comparison of surveys used to measure physical activity. *Aust NZ J Pub Health* 2004; 28: 128-134.
2. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Circulation* 2007; 116: 1081-1093.
3. Sjöström M, Oja P, Hagströmer M, Smith BJ, Bauman A. Health-enhancing physical activity across European Union countries: the Eurobarometer study. *J Pub Health* 2006; 14: 291-300.
4. Armstrong T, Bull F. Development of the World Health Organization Global Physical Activity Questionnaire (GPAQ). *J Pub Health* 2006; 14: 66-70.
5. Martinez-Gonzalez MA, Varo JJ, Santos JL, de Irala J, Gibney M, Kearney J, et al. Prevalence of physical activity during leisure time in the European Union. *Med Sci Sports Exerc* 2001; 33: 1142-1146.
6. Rutten A, Abu-Omar K. Prevalence of physical activity in the European Union. *Soz Präventivmed* 2004; 49: 281-289.
7. Guthold R, Ono T, Strong K, Chatterji S, Morabia A. World-wide variability in physical inactivity. A 51 country study. *Am J Prev Med* 2008; 34: 486-494.

major risk factors (chap. 8, vol. 1). World Health Organization: Geneva, 2004.

3. Yoshiike N, Seino F, Tajima S, Arai Y, Kawano M, Furuhashi T, et al. Twenty-year changes in the prevalence of overweight in Japanese adults: the National Nutrition Survey 1976-1995. *Obes Rev* 2002; 3: 183-190.
4. Bauman A, Allman-Farinelli M, Huxley R, James WPT. Leisure-time physical activity alone may not be a sufficient public health approach to prevent obesity—a focus on China. *Obes Rev* 2008; 9 suppl. 1: 119-120.
5. Ezzati M, Hoorn SV, Lawes CMM, Leach R, James WPT, Lopez AD, et al. Rethinking the “diseases of affluence” paradigm: economic development and global patterns of nutritional risks in relation to economic development. *PLoS Med* 2005; 2: E133.
6. Chen CM. Overview of obesity in Mainland China. *Obes Rev* 2008; 9 suppl. 1: 13-20.
7. James WPT, Schofield EC. *Human energy requirements: a manual for planners and nutritionists*. Published by arrangement with the Food and Agriculture Organization of the United Nations. Oxford University Press: Oxford, 1990.
8. Allman-Farinelli MA, Chey T, Bauman AE, Gill T, James WPT. Age, period and birth cohort effects on prevalence of overweight and obesity in Australian adults from 1990 to 2000. *Eur J Clin Nutr* 2007; 62: 898-907.
9. James WPT. The epidemiology of obesity: the size of the problem. *J Int Med* 2008; 263: 336-352.

Chapter 12

1. Katzmarzyk PT, Janssen I. The economic costs associated with physical inactivity and obesity in Canada: an update. *Can J Appl Physiol* 2004; 29: 90-115.
2. Yates J, Murphy C. A cost benefit analysis of weight management strategies. *Asia Pac J Clin Nutr* 2006; 15: 74-79.
3. Great Britain Parliament House of Commons Health Committee. *Obesity: third report of session 2003-04. Report, together with formal minutes* (vol. 1). London: Parliament. HMSO; 2004.
4. Wolf AM, Colditz GA. Current estimates of the economic cost of obesity in the United States. *Obes Res* 1998; 6: 97-106.
5. Zhao W, Zhai Y, Hu J, et al. Economic burden of obesity-related chronic diseases in Mainland China. *Obes Rev* 2008; 9: 62-67.
6. Raebel MA, Malone DC, Conner DA, Xu S, Porter JA, Lant FA. Health services use and health care costs of obese and nonobese individuals. *Arch Intern Med* 2004; 164: 2135-2140.
7. Wang F, McDonald T, Champagne LJ, Edington DW. Relationship of body mass index and physical activity to health care costs among employees. *J Occup Environ Med* 2004; 46: 428-436.
8. Finkelstein MM. Obesity, cigarette smoking and the cost of physicians' services in Ontario. *Can J Pub Health* 2001; 92: 437-440.
9. Trogon JG, Finkelstein EA, Hylands T, Dellea PS, Kamal-Bahl SJ. Indirect costs of obesity: a review of the current literature. *Obes Rev* 2008; 9: 489-500.
10. Allison DB, Zannolli R, Narayan KM. The direct health care costs of obesity in the United States. *Am J Pub Health* 1999; 89: 1194-1199.
11. van Baal PH, Polder JJ, de Wit GA, et al. Lifetime medical costs of obesity: prevention no cure for increasing health expenditure. *PLoS Med* 2008; 5: e29.

Chapter 13

1. Lobstein T, Baur L, Uauy R. IASO International Obesity Task Force. Obesity in children and young people: a crisis in public health. *Obes Rev* 2004; 5: 4-104.
2. Wang Y, Lobstein T. Worldwide trends in childhood overweight and obesity. *Int J Pediatr Obes* 2006; 1: 11-25.
3. Dietz WH, Robinson TN. Use of the body mass index (BMI) as a measure of overweight in children and adolescents. *J Pediatr* 1998; 132: 191-193.
4. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000; 320: 1240-1243.
5. World Health Organization. Growth reference data for 5-19 years. World Health Organization: Geneva, 2007. www.who.int/growthref/. Accessed 4 April 2008.

Chapter 14

1. Rice DP, Hodgson TA, Kopstein AN. The economic costs of illness: a replication and update. *Health Care Financ Rev* 1985; 7: 61-80.
2. Wang F, McDonald T, Champagne LJ, Edington DW. Relationship of body mass index and physical activity to health care costs among employees. *J Occup Environ Med* 2004; 46: 428-436.
3. Wang F, McDonald T, Reffitt B, Edington DW. BMI, physical activity, and health care utilization/costs among Medicare retirees. *Obes Res* 2005; 13: 1450-1457.
4. Pratt M, Macera CA, Wang G. Higher direct medical costs associated with physical inactivity. *Physician Sportsmed* 2000; 28: 63-70.
5. Anderson LH, Martinson BC, Crain AL, et al. Health care charges associated with physical inactivity, overweight, and obesity. *Prev Chronic Dis* 2005; 2: A09.
6. Katzmarzyk PT, Janssen I, Ardern CI. Physical inactivity, excess adiposity and premature mortality. *Obes Rev* 2003; 4: 257-290.
7. Popkin BM, Kim S, Rusev ER, Du S, Zizza C. Measuring the full economic costs of diet, physical activity and obesity-related chronic diseases. *Obes Rev* 2006; 7: 271-293.
8. Katzmarzyk PT, Janssen I. The economic costs associated with physical inactivity and obesity in Canada: an update. *Can J Appl Physiol* 2004; 29: 90-115.
9. Allison DB, Zannolli R, Narayan KM. The direct health care costs of obesity in the United States. *Am J Pub Health* 1999; 89: 1194-1199.
10. van Baal PH, Polder JJ, de Wit GA, et al. Lifetime medical costs of obesity: prevention no cure for increasing health expenditure. *PLoS Med* 2008; 5: e29.

Chapter 15

1. Tjepkema M. Adult obesity. *Health Rep* 2006; 17: 9-25.
2. Colditz GA. Economic costs of obesity and inactivity. *Med Sci Sports Exerc* 1999; 31: S663-S667.
3. World Health Organization. *Preventing chronic diseases: a vital investment*. World Health Organization: Geneva, 2005.
4. Ganz ML. The economic evaluation of obesity interventions: its time has come. *Obes Res* 2003; 11: 1275-1277.
5. Roux L, Kuntz KM, Donaldson C, Goldie SJ. Economic evaluation of weight loss interventions in overweight and obese women. *Obesity (Silver Spring)* 2006; 14: 1093-1106.

6. Haby MM, Vos T, Carter R, Moodie M, Markwick A, Magnus A, et al. A new approach to assessing the health benefit from obesity interventions in children and adolescents: the assessing cost effectiveness in obesity project. *Int J Obes* 2006; 30: 1463-1475.
7. Roux L, Pratt M, Tengs TO, Yore MM, Yanagawa TL, Van Den Bos J, et al. Cost-effectiveness of community-based physical activity interventions. *Am J Prev Med* 2008; 35: 578-588.
8. Centers for Disease Control and Prevention. Guide to community preventive services (community guide). www.thecommunityguide.org/. Accessed 10 Sept 2008.
9. Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System: 2003 survey data. www.cdc.gov/bfss/technical_infodata/surveydata/2003.htm.

Chapter 16

1. Dishman RK, Berthoud HR, Booth FW, Cotman CW, Edgerton VR, Fleshner MR, et al. Neurobiology of exercise. *Obesity (Silver Spring)* 2006; 14: 345-356.
2. Snitker S, Tataranni PA, Ravussin E. Spontaneous physical activity in a respiratory chamber is correlated to habitual physical activity. *Int J Obes Relat Metab Disord* 2001; 25: 1481-1486.
3. Levine JA, Eberhardt NL, Jensen MD. Role of nonexercise activity thermogenesis in resistance to fat gain in humans. *Science* 1999; 283: 212-214.
4. Kotz CM, Teske JA, Billington CJ. Neuroregulation of non-exercise activity thermogenesis and obesity resistance. *Am J Physiol Regul Integr Comp Physiol* 2008; 294: R699-710.
5. Levine JA, Kotz CM. NEAT—non-exercise activity thermogenesis—egocentric & geocentric environmental factors vs. biological regulation. *Acta Physiol Scand* 2005; 184: 309-318.
6. Tou JC, Wade CE. Determinants affecting physical activity levels in animal models. *Exp Biol Med (Maywood)* 2002; 227: 587-600.
7. Seburn KL. Measurements taken using the Comprehensive Lab Animal Monitoring System. MPD:92. Mouse Phenome Database Web Site, The Jackson Laboratory, Bar Harbor, Maine USA. World Wide Web (URL: <http://www.jax.org/phenome>).
8. Nogueiras R, Lopez M, Lage R, Perez-Tilve D, Pfluger P, Mendieta-Zeron H, et al. Bsx, a novel hypothalamic factor linking feeding with locomotor activity, is regulated by energy availability. *Endocrinology* 2008; 149: 3009-3015.
9. Schoeller DA. The importance of clinical research: the role of thermogenesis in human obesity. *Am J Clin Nutr* 2001; 73: 511-516.
10. Speakman J, Hambly C, Mitchell S, Krol E. Animal models of obesity. *Obes Rev* 2007; 8: 55-61.

Chapter 17

1. Rankinen T, Bouchard C. Genetics of physical activity. In: Clement K, Sorensen TIA (eds.), *Obesity genomics and post-genomics*. Informa Healthcare: New York, 2008, pp. 277-286.
2. Stubbe JH, Boomsma DI, Vink JM, Cornes BK, Martin NG, Skytthe A, et al. Genetic influences on exercise participation in 37,051 twin pairs from seven countries. *PLoS ONE* 2006; 20: 1: e22.
3. Cai G, Cole SA, Butte N, Bacino C, Diego V, Tan K, et al. A quantitative trait locus on chromosome 18q for physical activity and dietary intake in Hispanic children. *Obesity* 2006 Sept; 14: 1596-1604.

4. Bray MS, Hagberg JM, Perusse L, Rankinen T, Roth SM, Wolfarth B, et al. The human gene map for performance and health-related fitness phenotypes: the 2005 update. *Med Sci Sports Exerc* 2009; 41: 35-73.
5. Simonen RL, Rankinen T, Perusse L, Leon AS, Skinner JS, Wilmore JH, et al. A dopamine D2 receptor gene polymorphism and physical activity in two family studies. *Physiol Behav* 2003; 78: 751-757.
6. Loos RJ, Rankinen T, Tremblay A, Perusse L, Chagnon Y, Bouchard C. Melanocortin-4 receptor gene and physical activity in the Quebec Family Study. *Int J Obes (London)* 2005; 29: 420-428.
7. Stefan N, Vozarova B, Del Parigi A, Ossowski V, Thompson DB, Hanson RL, et al. The Gln223Arg polymorphism of the leptin receptor in Pima Indians: influence on energy expenditure, physical activity and lipid metabolism. *Int J Obes Relat Metab Disord* 2002; 26: 1629-1632.
8. Winnicki M, Accurso V, Hoffmann M, Pawlowski R, Dorigatti F, Santonastaso M, et al. Physical activity and angiotensin-converting enzyme gene polymorphism in mild hypertensives. *Am J Med Genet A* 2004; 125: 38-44.
9. Simonen RL, Rankinen T, Perusse L, Rice T, Rao DC, Chagnon Y, et al. Genome-wide linkage scan for physical activity levels in the Quebec Family Study. *Med Sci Sports Exerc* 2003; 35: 1355-1359.
10. De Moor MH, Spector TD, Cherkas LF, Falchi M, Hottenga JJ, Boomsma DI, et al. Genome-wide linkage scan for athlete status in 700 British female DZ twin pairs. *Twin Res Hum Genet* 2007; 10: 812-820.
11. De Moor MH, Posthuma D, Hottenga JJ, Willemsen G, Boomsma DI, De Geus EJ. Genome-wide linkage scan for exercise participation in Dutch sibling pairs. *Eur J Hum Genet* 2007; 15: 1252-1259.
12. De Moor MH, Stubbe JH, Boomsma DI, De Geus EJ. Exercise participation and self-rated health: do common genes explain the association? *Eur J Epidemiol* 2007; 22: 27-32.

Chapter 18

1. Gluckman PD, Hanson MA. Living with the past: evolution, development, and patterns of disease. *Science* 2004; 305: 1733-1736.
2. Vickers MH, Breier BH, McCarthy D, Gluckman PD. Sedentary behavior during postnatal life is determined by the prenatal environment and exacerbated by postnatal hypercaloric nutrition. *Am J Physiol Regul Integr Comp Physiol* 2003; 285: R271-273.
3. Watkins AJ, Wilkins A, Cunningham C, Perry VH, Seet MJ, Osmond C, et al. Low protein diet fed exclusively during mouse oocyte maturation leads to behavioural and cardiovascular abnormalities in offspring. *J Physiol* 2008; 586: 2231-2244.
4. Owen D, Matthews SG. Repeated maternal glucocorticoid treatment affects activity and hippocampal NMDA receptor expression in juvenile guinea pigs. *J Physiol* 2007; 578: 249-257.
5. Dahlgren J, Nilsson C, Jennische E, Ho HP, Eriksson E, Niklasson A, et al. Prenatal cytokine exposure results in obesity and gender-specific programming. *Am J Physiol Endocrinol Metab* 2001; 281: E326-334.
6. Meaney MJ, Szyf M. Environmental programming of stress responses through DNA methylation: life at the interface between a dynamic environment and a fixed genome. *Dialog Clin Neurosci* 2005; 7: 103-123.