

grassland weedy species. *Journal of Plant Diseases and Protection*, Special Issue XX: 377-383.

Pavlů V., Gaisler J., Hejcman M., Pavlů L. (2008): Effect of different grazing intensity on weed control under conditions of organic farming. *Journal of Plant Diseases and Protection*, Special Issue XXI: 441-446.

Pavlů V., Hejcman M., Pavlů L., Gaisler J. (2003): Effect of rotational and continuous grazing on vegetation of an upland grassland in the Jizerské hory Mts., Czech Republic. *Folia Geobotanica*, 38: 21-34.

Pavlů V., Hejcman M., Pavlů L., Gaisler J., Nežerková P. (2006): Effect of continuous grazing on forage quality, quantity and animal performance. *Agriculture, Ecosystems and Environment*, 113: 349-355.

Pavlů V., Hejcman M., Pavlů L., Gaisler J., Nežerková P., Guerochich Andaluz M. (2005): Vegetation changes after cessation of grazing management in the Jizerské Mountains (Czech Republic). *Annales Botanici Fennici*, 42: 343-349.

## V. PŘEHLED POUŽITÉ LITERATURY

Bakker J.P., Elzinga J.A., de Vries Z. (2002): Effects of long-term cutting a grassland system: perspectives for restoration of plant communities on nutrient-poor soils. *Applied Vegetation Science*, 5: 107-120.

Begon M., Townsend C.R., Harper J.L. (2005): *Ecology: From Individuals to Ecosystems*. Wiley-Blackwell, Oxford, UK.

Chytrý, M. (ed.) (2007): *Vegetace České republiky. 1. Travná a keříčková vegetace*. Academia, Praha.

Correll O., Isselstein J., Pavlů V. (2003): Studying spatial and temporal dynamics of sward structure at low stocking densities: the use of an extended rising-plate-meter method. *Grass and Forage Science*, 58, 450-454.

Dierschke H., Peppler-Lisbach C. (2009): Conservation and regeneration of structure and floristic biodiversity of montane meadows – 15 years of scientific contribution to conservation management in the Harz Mountains. *Tuexenia*, 29: 145-179.

Duffková R. (2008): Evaluation of management-dependent changes in the water regime of extensive grasslands. *Soil and Water Research*, 3: 1-11.

Duru M., Ansquer P., Jouany C., Theau J.P., Cruz P. (2010): Comparison of methods for assessing the impact of different disturbances and nutrient conditions upon functional characteristics of grassland communities. *Annals of Botany*, 106: 823-831.

Eler K., Vidrih M., Batič F. (2005): Vegetation characteristics in relation to different management regimes of calcareous grassland: A functional analysis using plant traits. *Phyton - Annales Rei Botanicae*, 45: 417-426.

Grime J.P., Hodgson J.G., Hunt R. (1988): *Comparative plant ecology: a functional approach to common British species*. Unwin Hyman, London, UK.

Hansson M., Fogelfors H. (2000): Management of a semi-natural grassland; Results from a 15-year-old experiment in southern Sweden. *Journal of Vegetation Science*, 11: 31-38.

Hejcman M., Auf D., Gaisler J. (2005): Year-round cattle grazing as an alternative management of hay meadows in the Giant Mts. (Krkonoše, Karkonosze), The Czech Republic. *Ekologia – Bratislava*, 24: 419-429.

Hellström K., Huhta A.P., Rautio P., Tuomi J. (2006): Search for optimal mowing regime - slow community change in a restoration trial in northern Finland. *Annales Botanici Fennici*, 43: 338-348.

Huhta A.P., Rautio P., Tuomi J., Laine K. (2001): Restorative mowing on an abandoned semi-natural meadow: short-term and predicted long-term effects. *Journal of Vegetation Science*, 12: 677-686.

Kahmen S., Poschlod P. (2008): Effects of grassland management on plant functional trait composition. *Agriculture, Ecosystems and Environment*, 128: 137-145.

Klimek S., Kemmermann A.R., Hofmann M., Isselstein J. (2007): Plant species richness and composition in managed grasslands: The relative importance of field management and environmental factors. *Biological Conservation*, 134: 559-570.

Klimeš L., Klimešová J. (2002): The effect of mowing and fertilization on carbohydrate reserves and regrowth of grasses: do they promote plant coexistence in species-rich meadows? *Evolutionary Ecology*, 15: 363-382.

Krahulec F., Blažková D., Balátová-Tuláčková E., Štursa J., Pecháčková S., Fabšičová M. (1996): Louky Krkonoše: Rostlinná společenstva a jejich dynamika [Grasslands of the Krkonoše Mountains: Plant communities and their dynamics]. *Opera Concorctica*, 33: 3-250.

Krahulec F., Skálová H., Herben T., Hadincová V., Wildová R., Pecháčková S. (2001): Vegetation changes following sheep grazing in abandoned mountain meadows. *Applied Vegetation Science*, 4: 97-102.

Laser H. (2002): Long-term and short-term effects of undisturbed plant succession, mulching, and meadow utilisation on the botanical diversity

in a moist *Arrhenatherion elatioris*. *Grassland Science in Europe*, 7: 806-807.

Lexa M., Krahulec F. (2000): Vliv mulčování na rozkladné procesy a druhové složení horských luk v Krkonoších. *Opera Concorctica*, 36: 571-577.

Martin G., Hossard L., Theau J.P., Therond O., Josien E., Cruz P., Rellier J.P., Martin-Clouaire R., Duru M. (2009): Characterizing potential flexibility in grassland use. Application to the French Aubrac area. *Agronomy for Sustainable Development*, 29: 381-389.

Mašková Z., Doležal J., Květ J., Zemek F. (2009): Long-term functioning of species-rich mountain meadow under different management regimes. *Agriculture, Ecosystems and Environment*, 132: 192-202.

Mehlich A. (1984): Mehlich no. 3 soil test extractant – a modification of Mehlich no. 2. *Communications in Soil Science and Plant Analysis*, 15: 1409-1416.

Moog D., Poschlod P., Kahmen S., Schreiber K.F. (2002): Comparison of species composition between different grassland management treatments after 25 years. *Applied Vegetation Science*, 5: 99-106.

Pecháčková S., Krahulec F. (1995): Efficient nitrogen economy – key to the success of *Polygonum bistorta* in an abandoned mountain meadow. *Folia Geobotanica*, 30: 211-222.

Römermann Ch., Bernhardt-Römermann M., Kleyer, M., Poschlod P. (2009): Substitutes for grazing in semi-natural grasslands – do mowing or mulching represent valuable alternatives to maintain vegetation structure. *Journal of Vegetation Science*, 20: 1086-1098.

Ryser P., Langenauer R., Gigon A. (1995): Species richness and vegetation structure in a limestone grassland after 15 years management with 6 biomass removal regimes. *Folia Geobotanica et Phytotaxonomica*, 30: 157-167.

Wahlman H., Milberg P. (2002): Management of semi-natural grassland vegetation: evaluation of a long-term experiment in southern Sweden. *Annales Botanici Fennici*, 39: 159-166.