

10. Literatura

1. Abrams, P., Cardozo, L., Fall, M., et al. The standardisation of terminology in lower urinary tract function: report from the standardisation sub-committee of the International Continence Society. *Urology*, 2003, 61, p. 37–49.
2. Krhut, J. *Neurourologie*. 1. vydání, Praha: Galén, 2005, 141 s.
3. Krofta, L. Operační léčba stresové inkontinence moči. *Urolog pro Praxi*, 2005, 6, s. 67–73.
4. Halaška, M. *Urogynekologie*. 1. vydání, Praha: Galén, 2004, 256 s.
5. Zachoval, R., Krhut, J., Zámečník, L., et al. Dotazníky hodnotící kvalitu života u pacientů s inkontinencí moči a hyperaktivním měchýřem. *Urolog pro Praxi*, 2006, 7, s. 286–296.
6. Kolombo, I., Kolombová, J., Poršová, M., et al. Stresová inkontinence u žen – 2. část. *Urolog pro Praxi*, 2009, 10, s. 11–20.
7. Dzvinčuk, P., Muller, O., Látalová, E. Inkontinence moči z pohledu gynekologa. *Interní Med*, 2008, 10, s. 90–95.
8. Vidlář, A., Vrtal, R., Študent, V. Patofyziologie stresové inkontinence u žen. *Urolog pro Praxi*, 2008, 9, s. 133–136.
9. Colombo, M., Scalabrino, S., Maggiony, A. Burch kolposuspension versus modified Marshall-Marchetti-Krantz uretropexy for primary genuine stress urinary incontinence: a prospective, randomized clinical trial. *Am J Obstet Gynecol*, 1994, 171, p. 1573–1579.
10. Ulmsten, U., Falconer, C., Johnson, P., et al. A multicenter study of tension-free vaginal tape (TVT) for surgical treatment of stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*, 1998, 9, p. 210–213.
11. Delorme, E. Transobturator urethral suspension: miniinvasive procedure in the treatment of stress urinary incontinence in women. *Prog Urol*, 2001, 11, p. 1306–1313.
12. De Leval, J. Novel surgical technique for the treatment of female stress urinary incontinence: transobturator vaginal tape: inside-out. *Eur Urol*, 2003, 44, p. 724–730.
13. Huvar, I. Tot – nový „zlatý standard“ v řešení ženské stresové inkontinence? *Urol List*, 2009, 7, s. 37–41.

14. Krofta, L., Feyereisl, J., Velebil, P., et al. TVT-S for surgical treatment of stress urinary incontinence: prospective trial, 1-year follow-up. *Int Urogynecol J Pelvic Floor Dysfunct*, 2010, 21, p. 779–785.
15. Masata, J., Martan, A., Svabik, K., et al. Long-term efficacy of TVT procedure in the treatment of stress urinary incontinence. *Ceska Gynekol*, 2009, 74, p. 118–123.
16. Neuman, M. Tvt and Tvt-Obturator: comparison of two operative procedures. *Eur J Obstet Gynecol Reprod Biol*, 2007, 131, p. 89–92.
17. Irwin, D. E., Abrams, P., Milsom, I., et al. Understanding the elements of overactive bladder: questions raised by the EPIC study. *BJU Int*, 2008, 101, p. 1381–1387.
18. Irwin, D. E., Milsom, I., Hunskaar, S., et al. Population-based survey of urinary incontinence, overactive bladder, and other lower urinary tract symptoms in five countries: results of the EPIC study. *Eur Urol*, 2006, 50, p. 1306–1314; discussion p. 1314–1305.
19. Saleem, A. In women with urinary incontinence how necessary is cystometry? *J Pak Med Assoc*, 2010, 60, p. 356–359.
20. Kraus, S. R., Ruiz-Cerda, J. L., Martire, D., et al. Efficacy and Tolerability of Fesoterodine in Older and Younger Subjects With Overactive Bladder. *Urology*, 2010, Oct. 23 (Epub ahead of print).
21. Krhut, J. *Hyperaktivní močový měchýř*. Praha: Maxdorf, 2007, 148 s.