

References

- ALLEN J. R. L. (1992): Transport-Hydrodynamics. - in: BRIGGS E. G.-CROWTHER P. R.(eds.): Palaeobiology. A synthesis. 227-230. Oxford.
- BARRANDE J. (1852): Systeme silurien du centre de la Boheme. Iere partie. Recherches paleontologiques. I. Trilobites. - Prag, Paris.
- BOUCOT A. J.(1975): Evolution and extinction rates controls. - Elsevier Sci. Publ. Amsterdam.
- (1992): Evolution of communities. - in: BRIGGS G. - CROWTHER P. R. (eds.): Palaeobiology. A synthesis. - 391-394. Oxford.
- BOUCOT A. J. - CARNEY S. R. (1981): Principles of benthic marine paleoecology.- Academic Press. New York.
- BRETT C. E. (1992): Destructive Taphonomic processes and skeletal durability. - in: BRIGGS E. G. - CROWTHER P. R.(eds.): Palaeobiology. A synthesis. 223-226. Oxford.
- BRETT C. E. - BAIRD G. G. (1986): Comparative taphonomy: A key to paleoenvironmental interpretation based on fossil preservation. - Palaios, 1, 207-227. Ann Arbor.
- BRETT C. E. - BORDEAUX Y. L. (1991): Taphonomy of brachiopods from a Middle Devonian shell bed: Implications for the genesis of skeletal accumulations. - in: MACKINNON D. I. - LEE D.E. - CAMPBELL J. D. (eds.): Brachiopods Through Time, 219-226. Balkema. Rotterdam.
- BRETT C. E. - SPEYER S. E. (1992): Taphofacies.- in: BRIGGS E. G.- CROWTHER P. R.(eds.): Palaeobiology. A synthesis. 223-226. Oxford.
- BROOKFIELD M. E. (1973); The life and death of

- Torquirhynchia inconstans (Brachiopoda, Upper Jurassic) in England. - Palaeogeogr. Palaeoclimatol. Palaeoecol., 13, 241-259. Amsterdam.
- CRAME J. A. (1992): Trophic structure. - in: BRIGGS E. G. - CROWTHER P. R. (eds.): Palaeobiology. A synthesis. 258-263. Oxford.
- FRÝDA J. (1988): A new species of Modestopira (Gastropoda) from Ordovician of Bohemia. - Věst. Ústř. Úst. geol., 63, 4, 227-232. Praha.
- FÜRSICH F. T. (1992): Fossil Concentrations and life and death assemblages. - in: BRIGGS E. G. - CROWTHER P. R. (eds.): Palaeocology. A synthesis. 235-239. Oxford.
- FÜRSICH F. T. - HURST J. M. (1974): Environmental factors determining the distribution of brachiopods. - Palaeontology, 17, 4, 879-900. London.
- HAVLÍČEK V. (1949): Orthacea vrstev komárovských (d) (Brachiopoda). - Věst. Ústř. Úst. geol., 14, 249-264. Praha.
- (1977): Brachiopods of the order Orthida in Czechoslovakia. - Rozp. Ústř. Úst. geol., 44, 1-327. Praha.
- (1981): Development of a linear sedimentary depression exemplified by the Prague Basin (Ordovician-Middle Devonian; Barrandian area - central Bohemia). - Sbor. geol. Věd, Geol., 35, 7-48. Praha.
- (1982a): Ordovician in Bohemia: development of the Prague Basin and its benthic communities. - Sbor. geol. Věd, Paleont., 25, 9-82. Praha.
- (1982b): Lingulacea, Paterinacea, and Siphonotretacea (Brachiopoda) in the Lower Ordovician sequence of Bohemia. - Sbor. geol. Věd, Paleont., 25, 9-82. Praha.
- (1989): Climatic changes and development of benthic communities through the Mediterranean Ordovician. - Sbor. geol. Věd, Geol., 44, 79-119. Praha.
- HAVLÍČEK V. - ŠNAJDR M. (1957): Faciální vývoj skidavu, llanvirnu a llandeila v Barrandienu. - Sbor. Ústř. Úst.

- geol. Odd. geol., 24, 540-600. Praha.
- HAVLÍČEK V. - VANĚK J. (1990): Ordovician invertebrate communities in black-shale lithofacies (Prague basin, Czechoslovakia). - Věst. Ústř. Úst. geol., 65, 4, 223-236. Praha.
- HOLLAND S. M. (1988): Taphonomic effect of sea-floor exposure on an Ordovician brachiopod assemblages. - Palaios, 3, 588-597. Ann Arbor.
- HOLMER L. E. (1986): Inarticulate brachiopods around the Middle-Upper Ordovician boundary in Västergötland. - Geol. Forening. Forh., 108, 97-126. Stockholm.
- (1989): Middle Ordovician phosphatic inarticulate brachiopods from Västergötland and Dalarna, Sweden. - Fossils Strata, 26, 1-172. Oslo.
- CHLUPÁČ I. et al. (1992): Paleozoikum Barrandienu. Kambrium-Devon. - Ústř. Úst. geol., Praha.
- JOHNSON J. G. (1990): Lower and Middle Devonian brachiopod dominated communities of Nevada, and their position in biofacies-province-realm model. - J. Paleont., 64, 902-943. Praha.
- KLOUČEK C. (1926): O faunistických obzorech ve vrstvách komárovských (d). - Věst. St. geol. Úst. Čs. Republ., 2, 53-57. Praha.
- KUKAL Z. (1959): Petrografický výzkum vrstev klabavských barrandienského ordoviku. - Sbor. Ústř. Úst. geol., Odd. geol., 25, 7-80. Praha.
- MERGL M. (1979): *Eccoptochile* aff. *clavigera* (Beyrich, 1845) and *Pliomerops lindaueri* (Barrande, 1846). - Věst. Ústř. Úst. geol., 54, 3, 175-177. Praha.
- (1981): The genus *Orbithele* (Brachiopoda, Inarticulata) from the Lower Ordovician of Bohemia and Morocco. - Věst. Ústř. Úst. geol., 56, 5, 287-292. Praha.
- (1991a): Arenig (Lower Ordovician) orthide brachiopods from the Prague Basin, Bohemia. - Čas. Mineral. Geol., 36, 1, 1-12. Praha.
- (1991b): New Lower Ordovician (Arenig) trilobite

- assemblages in Bohemia - Čas. Mineral. geol., 36, 4, 193-203. Praha.
- NOBLE J. P. A. - LOGAN A. (1981): Size-frequency distributions and taphonomy of brachiopods: a recent model - Palaeogeogr. Palaeoclimatol. Palaeoecol., 36, 185-20. Amsterdam.
- PERNER J. (1900): Miscellanea silurica Bohemiae. Příspěvky k poznání českého siluru. I. - Palaeontogr. Bohem., 5, 1-16. Praha.
- PETRÁNEK J. (1975): Ložiska sedimentárních železných rud v Mníšku a Komárově. - Stud. ČSAV, 6, 1-99. Praha.
- PICKERILL R. K. - BRENCHLEY P. J. (1979): Caradoc marine benthic communities of the South Berwyn Hills, North Wales. - Palaeontology, 22, 1, 229-264. London.
- RICHARDS R. P. - Bambach R. K. (1975): Population dynamics of some Palaeozoic brachiopods and their paleontological significance. - J. Paleont., 49, 5, 775-798. Tulsa.
- SCHALLREUTER R. - KRŮTA M. (1988): Ordovician Ostracodes of Bohemia. - Mitt. Geol. Palaont. Inst. Univ. Hamburg, 67, 99-119. Hamburg.
- THAYER C. W. (1975): Size-frequency and population structure of brachiopods. - Palaeogeogr. Palaeoclimatol. Palaeoecol., 17, 139-148. Amsterdam.
- VEEVERS J. J. (1959): The type species of Productella, Emanuella, Crurithyris and Ambocoelia. - J. Palaeont., 33, 5, 902-908. Tulsa.
- VELBEL M. A. - BRANDT D. S. (1989): Differential preservation of brachiopod valves: taphonomic bias in Platystrophia ponderosa. - Palaios, 4, 193-195.
- WANG Y. - BOUCOT A. J. - RONG J. - YANG X. (1987): Community Palaeoecology as Geologic Tool: The Chinese Ashgillian-Eifelian (latest Ordovician through early Middle Devonian) as an example. - Spec. Pap. Geol. Soc. Amer., 211, 1-100. Ann Arber.
- WATKINS R. - BERRY W. B. N. - BOUCOT A. J. (1973): Why "communities" ?. - Geology, 1, 55-58. Boulder.