

LITERATURE

- CETL, I.: Some developmental features in natural populations of *Arabidopsis thaliana* (L.) Heynh. Arabidopsis Research, Rep. Int. Symp. Göttingen, 46–52, 1965.
- CETL, I., and J. DOBROVOLNÁ: Further data on the developmental characters of natural populations of *Arabidopsis thaliana* (L.) Heynh. from Western Moravia. Arabidopsis Inform. Serv. 5:15–16, 1968.
- CETL, I., J. DOBROVOLNÁ, and E. EFFMERTO VÁ: Distribution of spring and winter types in the local populations of *Arabidopsis thaliana* (L.) Heynh. from various localities in Western Moravia. Arabidopsis Inform. Serv. 2:3, 1965.
- CETL, I., J. DOBROVOLNÁ, and E. EFFMERTO VÁ: The role of the temperature factor in the geographical distribution of "summer annual" and "winter annual" natural populations of *Arabidopsis thaliana* (L.) Heynh. Arabidopsis Inform. Serv. 4:9–10, 1967.
- CETL, I., J. DOBROVOLNÁ, and E. EFFMERTO VÁ: The developmental character of natural populations of *Arabidopsis thaliana* (L.) Heynh. in relation to the geographical-climatic conditions of localities. Folia Fac. Sci. Nat. Univ. Purkynianae Brunensis, Biologia IX, 5:37–49, 1968.
- CETL, I., and J. KUČERA: The effect of vernalization and photoperiod on the development of *Arabidopsis thaliana*. Proc. Symp. Differentiation Prague, 295–298, 1964.
- DIERKS, W.: Untersuchungen zum Heterosisproblem. Z. Pflanzenzücht. 40:67–102, 1958.
- DOBROVOLNÁ, J.: The variability of developmental characters in natural populations of *Arabidopsis thaliana* (L.) Heynh. Arabidopsis Inform. Serv. 4:6–7, 1967.
- DOBROVOLNÁ, J.: The heritability of characters "number of days to appearance of the flower primordia" and "number of rosette leaves" in a natural population of *Arabidopsis thaliana* (L.) Heynh. Arabidopsis Inform. Serv. 5:17, 1968.
- EFFMERTO VÁ, E.: The behaviour of "summer annual" "mixed", and "winter annual" natural populations as compared with early and late races in field conditions. Arabidopsis Inform. Serv. 4:8–9, 1967.
- EFFMERTO VÁ, E., and I. CETL: The vernalization requirement of "winter annual" populations from Western Moravia. Arabidopsis Inform. Serv. 3:10, 1966.
- EFFMERTO VÁ, E., and I. CETL: The behaviour of progenies derived from a "winter annual" natural population of *Arabidopsis thaliana* (L.) Heynh. Arabidopsis Inform. Serv. 5:16, 1968.
- FALCONER, D. S.: Introduction to quantitative genetics. New York, 1960.
- FORD, E. B.: Ecological genetics. London–New York, 1964.
- GREGORY, F. G., and G. G. HUSSEY: Photoperiodic responses of *Arabidopsis thaliana*. Proc. Linn. Soc. London 164:137–139, 1953.
- HÄRER, L.: Die Vererbung des Blühalters früher und später sommerjähriger Rassen von *Arabidopsis thaliana* (L.) Heynh. Beitr. Biol. Pflanzen 28:1–35, 1951.
- HUSSEIN, H. A. S., and J. H. van der VEEN: Induced mutations for flowering time. Arabidopsis Inform. Serv. 2:6, 1965.
- HUSSEIN, H. A. S., and J. H. van der VEEN: Genotypic analysis of induced mutations for flowering time and leaf number in *Arabidopsis thaliana*. Arabidopsis Inform. Serv. 5:30, 1968.
- JONES, MARY E.: Variation in flowering time of natural populations of *Arabidopsis thaliana* (L.) Heynh. with special reference to the breeding system. Arabidopsis Inform. Serv. 5:11–13, 1968.
- KEMPTHORNE, O., and O. B. TANDON: The estimation of heritability by regression of offspring on parent. Biometrics 1:90–100, 1953.
- KHOROSHAILOV, N. G.: (Ecological and geographical regularities of clover populations variability.) Bull. Appl. Bot., Gen. Plant Breed. 36:14–58, 1964.
- KISELEVA, S. D.: (Studying population composition of *Vicia villosa* Roth., *Vicia pannonica* Crantz and some other *Vicia* species for revealing possibilities of selection under different agricultural practices.) Bull. Appl. Bot., Gen. Plant Breed. 36:72–139, 1964.

- KUČERA, J., and I. CETL: (Über die Entwicklung der Lokalpopulationen von *Arabidopsis thaliana* [L.] Heynh. in der Abhängigkeit von Jarowisation und Photoperiode.) Publ. Fac. Sci. Univ. J. E. Purkyně, Brno 479:1–13, 1967.
- LAIBACH, F.: *Arabidopsis thaliana* (L.) Heynh. als Objekt für genetische und entwicklungsphysiologische Untersuchungen. Bot. Arch. 44:439–455, 1943.
- LAIBACH, F.: Über sommer- und winterannuelle Rassen von *Arabidopsis thaliana* (L.) Heynh. Beitr. Biol. Pflanzen 28:173–216, 1951.
- LAIBACH, F., and A. ZENKER: Zur Kälteeinflussung der Blütenbildung bei Langtagpflanzen. Planta 43:250–252, 1954.
- LEOKENE, L. V.: (Geographical variability within the Pannonica vetch [*Vicia pannonica* Crantz] population.) Bull. Appl. Bot., Gen. Plant Breed. 36:57–71, 1964.
- MÜLLER, A.: Zur Charakterisierung der Blüten und Infloreszenzen von *Arabidopsis thaliana* (L.) Heynh. Kulturpflanze 9:364–393, 1961.
- NAPP-ZINN, K.: Genetische Grundlagen des Kältebedürfnisses bei *Arabidopsis thaliana* (L.) Heynh. Ber. D. Bot. Ges. 68:369–373, 1965.
- NAPP-ZINN, K.: Untersuchungen zur Genetik des Kältebedürfnisses bei *Arabidopsis thaliana*. Z. ind. Abst. u. Vererbungslehre 88:253–285, 1957.
- NAPP-ZINN, K.: Über die genetischen Grundlagen des Vernalisationsbedürfnisses bei *Arabidopsis thaliana* I. Mitt. Z. Vererbungslehre 93:154–163, 1962.
- NAPP-ZINN, K.: Vernalisation und verwandte Erscheinungen. Handb. der Pflanzenphysiologie B. XVI, 24–75, 1961.
- NAPP-ZINN, K.: Über den Einfluss von Genen und Gibberellinen auf die Blütenbildung von *Arabidopsis thaliana*. Ber. D. Bot. Ges. 76:77–89, 1963.
- NAPP-ZINN, K.: Über genetische und entwicklungsphysiologische Grundlagen jahreszeitlicher Aspekte von Pflanzengesellschaften. Beitr. zur Phytologie, Festschrift H. Walter, Stuttgart 33–49, 1964.
- RATCLIFFE, D.: Observations on the physiology of flowering. *Arabidopsis Inform. Serv.* 1:8–9, 1964.
- RATCLIFFE, D.: The geographical and ecological distribution of *Arabidopsis* and comments on physiological variation. *Arabidopsis Research, Rep. Int. Symp. Göttingen* 37–45, 1965.
- RAZUMOV, V. I.: Sreda i osobennosti razvitija rastenij. Leningrad–Moskva, 1961.
- REEVE, E. C. R.: The variance of the genetic correlation coefficient. *Biometrics* 3:357–374, 1955.
- SINSKAJA, E. N.: (On general regularities of ecological-geographical variability within the populations of wild and cultivated plants.) Bull. Appl. Bot., Gen. Plant Breed. 36:3–13, 1964.
- SOSNA, M.: The origin of radiomutants of the chlorina type in *Arabidopsis thaliana* (L.) Heynh. Biol. Plantarum 4:255–259, 1962.
- SYROVÝ, S. (editor): Atlas podnebí Československé republiky. Praha, 1958.
- van der VEEN, J. H.: Genes for late flowering. *Arabidopsis Inform. Serv.* 2:5–6, 1965a.
- van der VEEN, J. H.: Genes for late flowering in *Arabidopsis*. *Arabidopsis Research, Rep. Int. Symp. Göttingen* 62–69, 1965b.
- VELEMÍNSKÝ, J., and T. GICHNER: Cytological and genetic effects of the insecticide Systox on *Vicia faba* L. and *Arabidopsis thaliana* (L.) Heynh. Biol. Plantarum 5:41–52, 1963.
- ZENKER, A.: Jarowisationsuntersuchungen an sommerannualen *Arabidopsis*-Rassen. Beitr. Biol. Pflanzen 32:135–170, 1955.