

LITERATURA

- [1] Freeman CR, Suissa S, Shenouda G, et al. Clinical experience with a single field rotational total skin electron irradiation technique for cutaneous T-cell lymphoma. *Radiotherapy and Oncology: Journal of the European Society for Therapeutic Radiology and Oncology*. 1992;24(3):155–62.
- [2] Trautinger F, Eder J, Assaf C, Bagot M, Cozzio A, Dummer R et al. European Organisation for Research and Treatment of Cancer consensus recommendations for the treatment of mycosis fungoides/Sézary syndrome – Update 2017. *Eur J Cancer*. 2017;77:57–74.
- [3] Belada D, Trněný M a kolektiv autorů Kooperativní lymfomové skupiny. Diagnostické a léčebné postupy u nemocných s maligními lymfomy. Leden 2023, třinácté (10. tištěné) vydání Czech Lymphoma Study Group, Kooperativní lymfomová skupina. ISBN: 978-80-905210-6-3.
- [4] Polívka J. Primární kožní lymfomy: současný pohled na vzácná hemato-onkologická onemocnění. *Onkologie*. 2016;10(2):66–71.
- [5] Talpur R, Singh L, Daulat S, Liu P, Seyfer S, Trynosky T et al. Long-term outcomes of 1263 patients with mycosis fungoides and Sézary syndrome from 1982 to 2009. *Clin Cancer Res*. 2012;18(18):5051–60.
- [6] Hoppe RT, Cox RS, Fuks Z, Price NM, Bagshaw MA, Farber EM. Electron-beam therapy for mycosis fungoides: the Stanford University experience. *Cancer Treat Rep*. 1979;63(4):691–700.
- [7] Novotvary 2008 ČR, Cancer incidence 2010 in the Czech Republic, ÚZIS 2011, ISSN 1210 857X, ISBN 978-80-7280-918-9 (ISBN 978-80-7472-034-5)
- [8] Novotvary 2013 ČR, ÚZIS 2017, Krejčí D, Svobodová I, Pehalová I, Talábová A, Mužík J, Dušek L, ÚZIS, Praha, 2022. Available from: <http://www.uzis.cz>.
- [9] Harrison C, Young J, Navi D, Riaz N, Lingala B, Kim Y et al. Revisiting low-dose total skin electron beam therapy in mycosis fungoides. *Int J Radiat Oncol Biol Phys*. 2011;81(4):e651–7.
- [10] Wilcox RA. Cutaneous T-cell lymphoma: 2014 update on diagnosis, risk-stratification, and management. *Am J Hematol*. 2014;89(8):837–51.
- [11] Krejsgaard T, Lindahl LM, Mongan NP, Wasik MA, Litvinov IV, Iversen L et al. Malignant inflammation in cutaneous T-cell lymphoma – a hostile takeover. *Semin Immunopathol*. 2017;39(3):269–82.
- [12] Fialová P. Kožní T buněčné lymfomy: mycosis fungoides a Sézaryho syndrom. Diagnostika a terapie. *Česko-slovenská dermatologie* 97. 2022;2:45–59.
- [13] Whittaker S, Hoppe R, Prince HM. How I treat mycosis fungoides and Sézary syndrome. *Blood*. 2016;127(25):3142–53.
- [14] Kamstrup MR, Gniadecki R, Iversen L, Skov L, Petersen PM, Loft A et al. Low-dose (10 Gy) total skin electron beam therapy for cutaneous T-cell lymphoma: an open clinical study and pooled data analysis. *Int J Radiat Oncol Biol Phys*. 2015;92(1):138–43.
- [15] Jawed SI, Myskowski PL, Horwitz S, Moskowitz A, Querfeld C. Primary cutaneous T-cell lymphoma (mycosis fungoides and Sézary syndrome): part II. Prognosis, management, and future directions. *J Am Acad Dermatol*. 2014;70(2):223.e1–17; quiz 240–2.
- [16] Chowdhary M, Chhabra AM, Kharod S, Marwaha, G. Total Skin Electron Beam Therapy in the Treatment of Mycosis Fungoides: A Review of Conventional and Low-Dose Regimens. *Clin Lymphoma Myeloma Leuk*. 2016;16(12):662–71.
- [17] Šlampa P et al. *Radiační onkologie*. Praha: Maxdorf, Jessenius; 2021. p. 771.
- [18] Ettler J, Polívka J, Arenberger P. Primární kožní lymfomy v dermatologické praxi. *Onkologie*. 2018;12/3:111–6.

- [61] Micaily B, Miyamoto C, Kantor G, Lessin S, Rook A, Brady L et al. Radiotherapy for unifocal mycosis fungoides. *Int J Radiat Oncol Biol Phys.* 1998;42(2):361–4.
- [62] Wang P, Gilbert M, Lim HW, McHargue C, Friedman BJ, Veenstra JJ et al. Single-Fraction Radiation Therapy for Localized Cutaneous T-Cell Lymphoma. *Pract Radiat Oncol.* 2023;13(4):346–50.
- [63] Becker M, Hoppe RT, Knox SJ. Multiple courses of high-dose total skin electron beam therapy in the management of mycosis fungoides. *Int J Radiat Oncol Biol Phys.* 1995;30;32(5):1445–9.
- [64] Hoppe RT, Fuks Z, Bagshaw MA. The rationale for curative radiotherapy in mycosis fungoides. *Int J Radiat Oncol Biol Phys.* 1977;2(9–10):843–51.
- [65] Koga K, Nishikawa K, Kuroki Y, Asada Y, Sumiyoshi A, Ogata K et al. Bone marrow suppression caused by whole body electron therapy for mycosis fungoides. *The British journal of radiology.* 1985;58(690):566–9.
- [66] Hoppe RT. Total skin electron beam therapy in the management of mycosis fungoides. *Frontiers of Radiation Therapy and Oncology.* 1991;25:80–9; discussion 132–3.
- [67] Morris SL, McGovern M, Bayne S, Wain M, Child F, Whittaker S. Results of a 5-week schedule of modern total skin electron beam radiation therapy. *Int J Radiat Oncol Biol Phys.* 2013;86(5):936–41.
- [68] Kumar PP, Good RR, Jones EO. Rotational total skin electron beam therapy. *Strahlenther Onkol.* 1988;164(2):73–8.
- [69] Kuten A, Stein M, Mandelzweig Y, Thatcher M, Yaacov G, Epelbaum R et al. Total-skin electron irradiation for cutaneous T-cell lymphoma: the Northern Israel Oncology Center experience. *Strahlenther Onkol.* 1991;167(7):392–6.
- [70] Wilson LD, Quiros PA, Kolenik SA, Heald PW, Braverman IM, Edelson RL et al. Additional courses of total skin electron beam therapy in the treatment of patients with recurrent cutaneous T-cell lymphoma. *J Am Acad Dermatol.* 1996;35(1):69–73.
- [71] Gamsiz H, Beyzadeoglu M, Sager O, Dincoglan F, Uysal B, Demiral S et al. Evaluation of mycosis fungoides management by total skin electron beam therapy with „translational technique“. *J BUON.* 2015;20(4):1124–31.
- [72] Taverniers A, Du-Thanh A, Charissoux M, Kerr C, Dereure O. Low- vs. Middle-dose Total Skin Electron Beam Therapy for Mycosis Fungoides: An Efficiency-based Retrospective Survey of Skin Response. *Acta Derm Venereol.* 2019;99(6):544–50.
- [73] Elsayad K, Kriz J, Moustakis C, Scobioala S, Reinartz G, Haverkamp U et al. Total Skin Electron Beam for Primary Cutaneous T-cell Lymphoma. *Int J Radiat Oncol Biol Phys.* 2015;93(5):1077–86.
- [74] Trautinger F, Knobler R, Willemze R, Peris K, Stadler R, Laroche L et al. EORTC consensus recommendations for the treatment of mycosis fungoides/Sézary syndrome. *Eur J Cancer.* 2006;42(8):1014–30.
- [75] Funk A, Hensley F, Krempien R, Neuhof D, Van Kampen M, Treiber M et al. Palliative total skin electron beam therapy (TSEBT) for advanced cutaneous T-cell lymphoma. *Eur J Dermatol.* 2008;18(3):308–12.
- [76] Klein RS, Dunlop JD, Samimi SS, Morrissey KA, Evans KG, Gardner JM et al. Improvement in peripheral blood disease burden in patients with Sézary syndrome and leukemic mycosis fungoides after total skin electron beam therapy. *J Am Acad Dermatol.* 2013;68(6):972–7.
- [77] Ysebaert L, Truc G, Dalac S, Lambert D, Petrella T, Barillot I et al. Ultimate results of radiation therapy for T1-T2 mycosis fungoides (including reirradiation). *Int J Radiat Oncol Biol Phys.* 2004;58(4):1128–34.
- [78] Ulutin HC, Beyan C, Pak Y. Total skin electron beam therapy for cutaneous T-cell lymphoma: Turkish experience with translational technique. *Haematologia (Budap).* 2002;32(4):397–403.
- [79] Anacak Y, Arican Z, Drumea K, Rosenblatt E, Tamir A, Chetver L et al. Total skin electron irradiation in mycosis fungoides: comparison between a modified Christie Hospital translation technique and the Stanford technique. *Leukemia@Lymphoma.* 2002;43(11):2093–7.
- [80] Wu JM, Leung SW, Wang CJ, Chui CS. Lying-on position of total skin electron therapy. *Int J Radiat Oncol Biol Phys.* 1997;39(2):521–8.

- [81] Gerbi BJ, Khan FM, Deibel FC, Kim TH. Total skin electron arc irradiation using a reclined patient position. *International journal of radiation oncology, biology, physics*. 1989;17(2):397–404.
- [82] Schaff EM, Rosenberg SA, Olson SJ, Howard SP, Bradley KA. Bone marrow suppression as a complication of total skin helical tomotherapy in the treatment of mycosis fungoides. *Radiat Oncol*. 2018;13(1):67.
- [83] Buglione M, Spiazzi L, Urpis M, Baushi L, Avitabile R, Pasinetti N et al. Light and shadows of a new technique: is ph.on total-skin irradiation using helical IMRT feasible, less complex and as toxic as the electrons one? *Radiat Oncol*. 2018;13(1):158.
- [84] Haraldsson A, Engleson J, Bäck SÅJ, Engelholm S, Engström, PE. A Helical tomotherapy as a robust low-dose treatment alternative for total skin irradiation. *J Appl Clin Med Phys*. 2019;20(5):44–54.
- [85] Yonekura K, Ichiki M, Takeda K, Uchiyama N, Nishida H, Dokiya T. Successful treatment of tumor stage mycosis fungoides with total skin helical tomotherapy. *J Dermatol*. 2022;49(2):289–93.
- [86] Antolak JA, Cundiff JH, Ha CS. Utilization of thermoluminescent dosimetry in total skin electron beam radiotherapy of mycosis fungoides. *Int J Radiat Oncol Biol Phys*. 1998;40(1):101–8.
- [87] Anacak Y, Arican Z, Bar-Deroma R, Tamir A, Kuten A. Total skin electron irradiation: evaluation of dose uniformity throughout the skin surface. *Med Dosim*. 2003;28(1):31–4.
- [88] Gamble LM, Farrell TJ, Jones GW, Hayward JE. Two-dimensional mapping of underdosed areas using radiochromic film for patients undergoing total skin electron beam radiotherapy, *International Journal of Radiation Oncology Biology Physics*. 2005;62(3):920–4.
- [89] Schüttrumpf L, Neumaier K, Maihoefer C, Niyazi M, Ganswindt U, Li M et al. Dose optimization of total or partial skin electron irradiation by thermoluminescent dosimetry. *Strahlenther Onkol*. 2018;194(5):444–53. English.
- [90] Gaspar-Olvera JC, Torres-García E, Oros-Pantoja R, Vieyra-Reyes P. Dosimetry of Rotational Total Skin Electron Irradiation in the Treatment of Mycosis Fungoides. *International Journal of Advanced Research in Physical Science (IJARPS)*. 2016;3(7):41–7.
- [91] Hoppe RT, Harrison C, Tavallae M, Bashey S, Sundram U, Li S, et al. Low-dose total skin electron beam therapy as an effective modality to reduce disease burden in patients with mycosis fungoides: results of a pooled analysis from 3 phase-II clinical trials. *J Am Acad Dermatol*. 2015;72(2):286–92.
- [92] Tandberg DJ, Craciunescu O, Kelsey CR. Radiation Therapy for Cutaneous T-Cell Lymphomas. *Dermatol Clin*. 2015;33(4):703–13.
- [93] Reddy S, Parker CM, Shidnia H, Gaffney MG, Blyton BD, Pruner LM et al. Total skin electron beam radiation therapy for mycosis fungoides. *Am J Clin Oncol*. 1992;15(2):119–24.
- [94] Rosenblatt E, Kuten A, Leviov M, Cederbaum M. Total skin electron irradiation in mycosis fungoides dose and fractionation considerations. *Leuk Lymphoma*. 1998;30(1–2):143–51.
- [95] Parida DK, Verma KK, Rath GK. Total skin electron irradiation treatment for mycosis fungoides with a new alternate daily treatment schedule to minimize radiation-associated toxicity: a preliminary experience. *Clinical and Experimental Dermatology*. 2009;34(5):e37–9.
- [96] Heumann TR, Esiashvili N, Parker SR, Switchenko JM, Dhabbaan A, Goodman M et al. Total skin electron therapy for cutaneous T-cell lymphoma using a modern dual-field rotational technique. *International journal of radiation oncology, biology, physics*. 2015;92(1):183–91.
- [97] Quiros PA, Kacinski BM, Wilson LD. Extent of skin involvement as a prognostic indicator of disease free and overall survival of patients with T3 cutaneous T-cell lymphoma treated with total skin electron beam radiation therapy. *Cancer*. 1996;77(9):1912–17.
- [98] Kamstrup MR, Specht L, Skovgaard GL, Gniadecki R. (2008) A prospective, open-label study of low-dose total skin electron beam therapy in mycosis fungoides. *International journal of radiation oncology, biology, physics*. 2008;71(4):1204–7.
- [99] Morris S, Scarisbrick J, Frew J, Irwin C, Grieve R, Humber C et al. The Results of Low-Dose Total Skin Electron Beam Radiation Therapy (TSEB) in Patients With Mycosis Fungoides From the UK Cutaneous Lymphoma Group. *Int J Radiat Oncol Biol Phys*. 2017;99(3):627–33.

- [100] Chowdhary M, Song A, Zaorsky NG, Shi W. Total skin electron beam therapy in mycosis fungoides-a shift towards lower dose? *Chin Clin Oncol*. 2019;8(1):9.
- [101] Dault JB, Slade AN, Zhao S, Song S. Comparison of low-dose and standard-dose total skin electron beam therapy in mycosis fungoides. *Leuk Lymphoma*. 2019;60(9):2334–6.
- [102] Elsayad K, Kroeger K, Greve B, Moustakis C, Assaf C, Stadler R et al. Low-dose total skin electron beam therapy: Quality of life improvement and clinical impact of maintenance and adjuvant treatment in patients with mycosis fungoides or Sezary syndrome. *Strahlenther Onkol*. 2020;196(1):77–84. English.
- [103] Newman NB, Patel CG, Ding GX, Zic JA, Zwerner J, Osmundson EC et al. N. Prospective observational trial of low-dose skin electron beam therapy in mycosis fungoides using a rotational technique. *J Am Acad Dermatol*. 2021;85(1):121–7.
- [104] Song A, Gochoco A, Zhan T, Sahu J, Alpdogan O, Porcu P et al. A prospective cohort study of condensed low-dose total skin electron beam therapy for mycosis fungoides: Reduction of disease burden and improvement in quality of life. *J Am Acad Dermatol*. 2020;83(1):78–85.
- [105] Smits K, Quint KD, Vermeer MH, Daniëls LA, Willemze R, Jansen PM et al. Total skin electron beam therapy for cutaneous T-cell lymphomas in the Netherlands: A retrospective analysis of treatment outcomes and selection for high or low dose schedule. *Clin Transl Radiat Oncol*. 2022;33:77–82.
- [106] Elsayad K, Rolf D, Sunderkötter C, Weishaupt C, Müller EC, Nawar T et al. Low-dose total skin electron beam therapy plus oral bexarotene maintenance therapy for cutaneous T-cell lymphoma. *J Dtsch Dermatol Ges*. 2022;20(3):279–85.
- [107] Rolf D, Eich HT. Radiotherapie kutaner Lymphome [Radiotherapy for cutaneous lymphomas]. *Dermatologie (Heidelb)*. 2022;73(10):772–81. German.
- [108] Ibanez-Rosello B, Bautista JA, Bonaque J. et al. Failure modes and effects analysis of total skin electronirradiation technique. *Clin Transl Oncol*. 2018;20:330–65.
- [109] Jeans EB, Hu YH, Stish BJ, King B, Davis M, Harmsen WS et al. Low-Dose Hypofractionated Total Skin Electron Beam Therapy for Adult Cutaneous T-Cell Lymphoma. *Pract Radiat Oncol*. 2020;10(6):e529–e537.
- [110] Zic JA, Ai W, Akilov OE, Carter JB, Duvic M, Foss F et al. United States Cutaneous Lymphoma Consortium recommendations for treatment of cutaneous lymphomas during the COVID-19 pandemic. *J Am Acad Dermatol*. 2020;83(2):703–4.
- [111] Duvic M, Lemak NA, Redman JR, Eifel PJ, Tucker SL, Cabanillas FF et al. Combined modality therapy for cutaneous T-cell lymphoma, *Journal of the American Academy of Dermatology*. 1996;34(6):1022–9.
- [112] Elsayad K, Stadler R, Steinbrink K, Eich HT. Combined total skin radiotherapy and immune checkpoint inhibitors: A promising potential treatment for mycosis fungoides and Sezary syndrome. *J Dtsch Dermatol Ges*. 2020;18(3):193–7.
- [113] Lutsyk M, Ben-Yosef R, Bergman R, Kuten A, Bar-Sela G. Total Skin Electron Irradiation and Sequential Malignancies in Mycosis Fungoides Patients: Longitudinal Study. *Clin Oncol (R Coll Radiol)*. 2018;30(10):618–24.
- [114] Šlampa P, Šlampa J. Sekundární nádory a radioterapie. *Klinická onkologie*. 2020;33(Suppl. 1):S48–51.
- [115] Lamoureux C, Gerard E, Ouhabache N, Toukal F, Pham-Ledard A, Beylot-Barry M. A case of potential radiation recall dermatitis induced by chemotherapy, following low-dose total-skin electron beam therapy. *Eur J Dermatol*. 2019;29(6):671–2.
- [116] Parida D, Verma K, Chander S, Joshi R, Rath G. Cutaneous T-cell lymphoma treated with electron beam irradiation in Indian patients. *International journal of dermatology*. 2001;40:295–7.
- [117] Desai KR, Pezner RD, Lipsett JA, Vora NL, Luk KH, Wong JY et al. Total skin electron irradiation for mycosis fungoides: relationship between acute toxicities and measured dose at different anatomic sites. *Int J Radiat Oncol Biol Phys*. 1988;15(3):641–5.
- [118] Samie FH. JAAD Game Changers: Acute toxicity and risk of infection during total skin electron beam therapy for mycosis fungoides. *J Am Acad Dermatol*. 2021;85(2):537.

- [119] Quinlan KE, Janiga JJ, Baran R, Lim HW. Transverse melanonychia secondary to total skin electron beam therapy: a report of 3 cases. *J Am Acad Dermatol*. 2005;53(2 Suppl 1):S112–4.
- [120] Martin MT, Lefaix JL, Delanian S. TGF-beta1 and radiation fibrosis: a master switch and a specific therapeutic target? *International journal of radiation oncology, biology, physics*. 2000;47(2):277–90.
- [121] Peter RU, Gottlöber P, Nadeshina N, Krähn G, Braun-Falco O, Plewig G. Interferon gamma in survivors of the Chernobyl power plant accident: new therapeutic option for radiation-induced fibrosis. *Int J Radiat Oncol Biol Phys*. 1999;45(1):147–52.
- [122] Gottlöber P, Steinert M, Bähren W, Weber L, Gerngroß H, Peter RU. Interferon-gamma in 5 patients with cutaneous radiation syndrome after radiation therapy, *International Journal of Radiation Oncology Biology Physics*. 2001;50(1):159–66.
- [123] Lloyd S, Chen Z, Foss FM, Girardi M, Wilson LD. Acute toxicity and risk of infection during total skin electron beam therapy for mycosis fungoides. *J Am Acad Dermatol*. 2013;69(4):537–43.
- [124] Korekawa A, Kaneko T, Nakajima K, Rokunohe D, Akasaka E, Nakano H et al. Mycosis fungoides bullosa associated with bullous pemphigoid. *Int J Dermatol*. 2015;54(9):e366–8.
- [125] Terezakis SA, Bohle GC, Lo YC, Berry SL, Yahalom J. Innovative Approaches to Radiation Treatment for Mycosis Fungoides in the Setting of Collagen Vascular Disease. *Case Rep Oncol Med*. 2015;2015 853823. doi:10.1155/2015/853823.
- [126] Kim YH, Bishop K, Varghese A, Hoppe RT. Prognostic factors in erythrodermic mycosis fungoides and the Sézary syndrome. *Arch Dermatol*. 1995;131(9):1003–8.
- [127] Rampino M, Ragona R, Monetti U, Mussano A, Guarneri A, Sannazzari GL. Total skin electron beam therapy in mycosis fungoides. Our experience from 1985 to 1999. *Radiol Med*. 2002;103(1–2):108–14. English, Italian.
- [128] Agar NS, Wedgeworth E, Crichton S, Mitchell TJ, Cox M, Ferreira S et al. Survival outcomes and prognostic factors in mycosis fungoides/Sézary syndrome: validation of the revised International Society for Cutaneous Lymphomas/European Organisation for Research and Treatment of Cancer staging proposal. *J Clin Oncol*. 2010;28(31):4730–9.
- [129] Hughes CF, Newland K, McCormack C, Lade S, Prince HM. Mycosis fungoides and Sézary syndrome: Current challenges in assessment, management and prognostic markers. *Australas J Dermatol*. 2016;57(3):182–91.
- [130] An Y, Jiang W, Andraos TY, Reddy JP, Yehia ZA, Lloyd S et al. Multi-institutional Investigation: Circulating CD4:CD8 ratio is a prognosticator of response to total skin electron beam radiation in mycosis fungoides, Radiotherapy and Oncology. 2019;131:88–92.
- [131] Herschel S, Zackheim H, Koh, Weinstock MA. Assessing Clinical Outcomes in Cutaneous T-cell Lymphoma, Hematology/Oncology Clinics of North America. 1995;9(5):1021–9.
- [132] Tadros AA, Tepperman BS, Hryniuk WM, Peters VG, Rosenthal D, Roberts JT et al. Total skin electron irradiation for mycosis fungoides: failure analysis and prognostic factors. *Int J Radiat Oncol Biol Phys*. 1983;9(9):1279–87.
- [133] Wilson LD, Quiros PA, Kolenik SA, Heald PW, Braverman IM, Edelson RL et al. Additional courses of total skin electron beam therapy in the treatment of patients with recurrent cutaneous T-cell lymphoma. *J Am Acad Dermatol*. 1996;35(1):69–73.
- [134] Boström A, Lindman H, Swartling C, Berne B, Bergh J. Potent corticosteroid cream (mometasone furoate) significantly reduces acute radiation dermatitis: results from a double-blind, randomized study. *Radiother Oncol*. 2001;59(3):257–65.
- [135] Lovgren ML, Scarisbrick JJ. Update on skin directed therapies in mycosis fungoides. *Chin Clin Oncol*. 2019;8(1):7.
- [136] Piccinno R, Caccialanz M, Percivalle S. Minimal stage IA mycosis fungoides. Results of radiotherapy in 15 patients. *J Dermatolog Treat*. 2009;20(3):165–8.
- [137] Esposito L, Piccinno R, Marchese L, Berti E. Results of radiotherapy in minimal stage mycosis fungoides: a reappraisal after ten years. *G Ital Dermatol Venereol*. 2020;155(4):483–6.
- [138] Fuks Z, Bagshaw MA. Total-skin electron treatment of mycosis fungoides. *Radiology*. 1971;100(1):145–50.

- [139] Wilson LD, Cooper DL, Goodrich AL, Friedman ND, Feldman AM, Braverman IM. Impact of non-CTCL dermatologic diagnoses and adjuvant therapies on cutaneous T-cell lymphoma patients treated with total skin electron beam radiation therapy. *Int J Radiat Oncol Biol Phys*. 1994;28(4):829–37.
- [140] Jones GW, Rosenthal D, Wilson LD. Total skin electron radiation for patients with erythrodermic cutaneous T-cell lymphoma (mycosis fungoides and the Sézary syndrome). *Cancer*. 1999;85(9):1985–95.
- [141] Yu JB, Khan AM, Jones GW, Reavely MM, Wilson LD. Patient perspectives regarding the value of total skin electron beam therapy for cutaneous T-cell lymphoma/mycosis fungoides: a pilot study. *Am J Clin Oncol*. 2009;32(2):142–4.
- [142] Kamstrup MR, Lindahl LM, Gniadecki R, Iversen L, Skov L, Petersen PM et al. Low-dose total skin electron beam therapy as a debulking agent for cutaneous T-cell lymphoma: an open-label prospective phase II study. *Br J Dermatol*. 2012;166(2):399–404.
- [143] Hinds GA, Alhariri J, Klein RQ, Wilson LD. Treatment of mycosis fungoides with total skin electron beam: response and relapse by ethnicity and sex. *Am J Clin Oncol*. 2013;36(5):481–5.
- [144] Piotrowski T, Fundowicz M, Pawlaczyk M. Total Skin Electron Beam Therapy with Rotary Dual Technique as Palliative Treatment for Mycosis Fungoides. *In Vivo*. 2018;32(3):517–22.
- [145] Georgakopoulos I, Papadavid E, Platoni K, Kougioumtzopoulou A, Patatoukas G, Kypraiou, E et al. Low dose total skin electron beam therapy for the management of T cell cutaneous lymphomas. *Dermatol Ther*. 2020;33(4):e13478.
- [146] Canu D, Pham-Ledard A, Ouhabache N, Beylot-Barry M. Changes in total skin electron beam therapy modalities for mycosis fungoides: A single-centre study. *Ann Dermatol Venereol*. 2022;149(3):180–4.
- [147] Muniesa C, Estrach T, Pujol RM, Garcia-Muret P, Gallardo F, Servitje O, Climent J. Folliculotropic mycosis fungoides: Clinicopathological features and outcome in a series of 20 cases, *Journal of the American Academy of Dermatology*. 2010;62(3):418–26.
- [148] Campbell BA, Ryan G, McCormack C, Tangas E, Bressel M, Twigger R et al. Lack of Durable Remission with Conventional-Dose Total Skin Electron Therapy for the Management of Sezary Syndrome and Multiply Relapsed Mycosis Fungoides. *Cancers (Basel)*. 2019;8;11(11):1758.
- [149] O'Malley JT, de Masson A, Lowry EL, Giobbie-Hurder A, LeBoeuf NR, Larocca C et al. Radiotherapy Eradicates Malignant T Cells and Is Associated with Improved Survival in Early-Stage Mycosis Fungoides. *Clin Cancer Res*. 2020;26(2):408–18.
- [150] Sklenář Z. Nová emolienca s obsahem močoviny v magistraliter přípravcích. *Česko-slovenská dermatologie*. 2010;85(5):274–9.
- [151] Drahotuská Z. Postavení emoliencií v současné léčbě atopické dermatitidy. *Dermatologie v praxi*. 2017;11(3):120–5.