

Bibliography

- [1] M. Shatruk, ThCr₂Si₂ structure type: The “perovskite” of intermetallics, *J. Solid State Chem.* **272**, 198 (2019).
- [2] R. Nath, Y. Singh, and D. C. Johnston, “Magnetic, Thermal and Transport Properties of Layered Arsenides BaRu₂As₂ and SrRu₂As₂”, *Phys. Rev. B* **79**, 174513 (2009).
- [3] D. C. Johnston, “Superconductivity above 10 K in Non-Cuprate Oxides”, in *100 Years of Superconductivity*, edited by H. Rogalla and P. Kes (CRC Press, Boca Raton, FL, 2012), pp. 239–244.
- [4] For a comprehensive early review of the Fe-based superconductors and parent compounds, see D. C. Johnston, *Adv. Phys.* **59**, 803 (2010).
- [5] C. Kittel and H. Kroemer, *Thermal Physics*, 2nd Edition (W. H. Freeman, New York, 1980).
- [6] A. Einstein, Die Plancksche Theorie der Strahlung und die Theorie der spezifischen Wärme, *Ann. Physik* **22**, 180 (1907).
- [7] A. C. Victor, Heat Capacity of Diamond at High Temperatures, *J. Chem. Phys.* **36**, 1903 (1962).
- [8] O. O. Vasil’ev, V. B. Muratov, and T. I. Duda, The Study of Low-Temperature Heat Capacity of Diamond: Calculation and Experiment, *J. Superhard Mater.* **32**, 375 (2010).
- [9] D. C. Johnston, Magnetic Susceptibility of Collinear and Noncollinear Heisenberg Antiferromagnets, *Phys. Rev. Lett.* **109**, 077201 (2012).
- [10] D. C. Johnston, Unified Molecular Field Theory for Collinear and Noncollinear Heisenberg Antiferromagnets, *Phys. Rev. B* **91**, 064427 (2015).
- [11] G. R. Stewart, Heavy-fermion systems, *Rev. Mod. Phys.* **56**, 755 (1984).
- [12] R. Stevens and J. Boerio-Goates, Heat capacity of copper on the ITS-90 temperature scale using adiabatic calorimetry, *J. Chem. Thermodyn.* **36**, 857 (2004).
- [13] C. M. Hurd, A Magnetic Susceptibility Apparatus for Weakly Magnetic Metals, and the Susceptibility of Pure Copper in the Range 6–300 °K, *Cryogenics* **6**, 264 (1966).
- [14] M. Shiroishi and M. Takahashi, Integral Equation Generates High-Temperature Expansion of the Heisenberg Chain, *Phys. Rev. Lett.* **89**, 117201 (2002).
- [15] F. J. Blatt, *Physics of Electronic Conduction in Solids* (McGraw-Hill, New York, 1968).
- [16] R. L. Powell, H. M. Roder, and W. J. Hall, Low-Temperature Transport Properties of Copper and Its Dilute Alloys: Pure Copper Annealed and Cold-Drawn, *Phys. Rev.* **115**, 314 (1959).
- [17] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Theory of Superconductivity, *Phys. Rev.* **108**, 1175 (1957).
- [18] B. Mühlischlegel, Die thermodynamischen Funktionen des Supraleiters, *Z. Phys.* **155**, 313 (1959). The heading of the fifth column of the table on page 322 should be $H_c^2/8\pi C_n T_c$ instead of $H_c^2/4\pi C_n T_c$. In our notation, $C_n = \gamma_n T_c$.
- [19] H. Padamsee, J. E. Neighbor, and C. A. Shiffman, Quasiparticle Phenomenology for Thermodynamics of Strong-Coupling Superconductors, *J. Low Temp. Phys.* **12**, 387 (1973).
- [20] J. P. Carbotte, Properties of boson-exchange superconductors, *Rev. Mod. Phys.* **62**, 1027 (1990).
- [21] J. D. Jackson, *Classical Electrodynamics* (Wiley, New York, 1962).
- [22] R. Combescot, Inequalities for $\Delta(T)/k_B T_c$: A test for weak-coupling theory applied to high- T_c superconductors, *Phys. Rev. B* **37**, 2235 (1988).
- [23] T. M. Mishonov, E. S. Penev, and J. O. Indekeu, Comment on “Anisotropic s-wave superconductivity in MgB₂”, *Phys. Rev. B* **66**, 066501 (2002); *Europhys. Lett.* **61**, 577 (2003).

- [24] D. C. Johnston, Magnetic Dipole Interactions in Crystals, *Phys. Rev. B* **93**, 014421 (2016).
- [25] L. A. Openov, Effect of nonmagnetic and magnetic impurities on the heat capacity jump in anisotropic superconductors, *Phys. Rev. B* **69**, 224516 (2004).
- [26] J. A. Osborn, Demagnetizing Factors of the General Ellipsoid, *Phys. Rev.* **67**, 351 (1945).
- [27] D.-X. Chen, J. A. Brug, and R. B. Goldfarb, Demagnetizing Factors for Cylinders, *IEEE Trans. Magn.* **27**, 3601 (1991).
- [28] A. Aharoni, Demagnetizing factors for rectangular ferromagnetic prisms, *J. Appl. Phys.* **83**, 3432 (1998).
- [29] D. C. Johnston, Elaboration of the α -Model Derived from the BCS Theory of Superconductivity, *Supercond. Sci. Technol.* **26**, 115011 (2013).
- [30] A. P. Ramirez, Strongly Geometrically Frustrated Magnets, *Annu. Rev. Mater. Sci.* **24**, 453 (1994).
- [31] A. P. Ramirez, in *Handbook of Magnetic Materials*, Vol. 13, edited by K. H. J. Buschow (North-Holland, Amsterdam, 2001), pp. 423–520.
- [32] R. Moessner and A. P. Ramirez, Geometrical Frustration, *Phys. Today* **59**(2), 24 (2006).
- [33] R. de L. Kronig and W. G. Penney, Quantum Mechanics of Electrons in Crystal Lattices, *Proc. Roy. Soc. London A* **130**, 499 (1931).
- [34] Wikipedia, https://en.wikipedia.org/wiki/Particle_in_a_one-dimensional_lattice
- [35] J. C. Wolfe, Summary of the Kronig-Penney electron, *Am. J. Phys.* **46**, 1012 (1978).
- [36] F. Bloch, Über die Quantenmechanik der Elektronen in Kristallgittern, *Z. Phys.* **52**, 555 (1929).
- [37] E. D. Mun, S. L. Bud'ko, H. Ko, G. J. Miller, and P. C. Canfield, Physical properties and anisotropies of the $R\text{NiGe}_3$ series ($R = \text{Y, Ce-Nd, Sm, Gd-Lu}$), *J. Magn. Magn. Mater.* **322**, 3527 (2010).
- [38] J. W. Stout and S. A. Reed, The Crystal Structure of MnF_2 , FeF_2 , CoF_2 , NiF_2 , and ZnF_2 , *J. Am. Chem. Soc.* **76**, 5279 (1954).
- [39] Charles A. Trapp, The Magnetic Susceptibility of Ferrous-Chloride and Manganous-Fluoride, Ph.D. Thesis, U. Chicago (1963).
- [40] C. Trapp and J. W. Stout, Magnetic Susceptibility of MnF_2 , *Phys. Rev. Lett.* **10**, 157 (1963).
- [41] Wikipedia, <https://en.wikipedia.org/wiki/Diamagnetism>.
- [42] *Metallic Shifts in NMR*, Part I, Editors B. Chalmers, J. W. Christian, and T. B. Massalski, *Prog. Mater. Sci.* **20** (Pergamon, Oxford, 1977), p. 5.
- [43] L. Corliss, Y. Delabarre, and N. Elliott, Paramagnetic Susceptibilities of Solid Solutions of MnF_2 and ZnF_2 , *J. Chem. Phys.* **18**, 1256 (1950).
- [44] X. Lin, B. Fauqué, and K. Behnia, Scalable T^2 resistivity in a small single-component Fermi surface, *Science* **349**, 945 (2015).
- [45] K. Kadowaki and S. B. Woods, Universal Relationship of the Resistivity and heat capacity in Heavy-Fermion Compounds, *Solid State Commun.* **58**, 507 (1986).
- [46] R. J. Goetsch, V. K. Anand, A. Pandey, and D. C. Johnston, Structural, Thermal, Magnetic and Electronic Transport Properties of the $\text{LaNi}_2(\text{Ge}_{1-x}\text{P}_x)_2$ System, *Phys. Rev. B* **85**, 054517 (2012).
- [47] F. Bloch, Nuclear Induction, *Phys. Rev.* **70**(7,8), 460 (1946).
- [48] A. Abragam, *Principles of Nuclear Magnetism* (Clarendon, Oxford, 1961).
- [49] C. P. Slichter, *Principles of Magnetic Resonance* (Harper & Row, New York, 1963).
- [50] A. Abragam and B. Bleaney, *Electron Paramagnetic Resonance of Transition Ions* (Dover, New York, 1970).
- [51] G. E. Pake and T. L. Estle, *The Physical Principles of Electron Paramagnetic Resonance*, 2nd edition (Benjamin, Reading, MA, 1973).
- [52] R. H. Taylor, Electron spin resonance of magnetic ions in metals—An experimental review, *Adv. Phys.* **24**, 681 (1975).
- [53] S. E. Barnes, Theory of electron spin resonance of magnetic ions in metals, *Adv. Phys.* **30**, 801 (1981).
- [54] H. Goldstein, C. P. Polle, and J. Safko, 3rd *Classical Mechanics* (Dorling Kindersley, India, 2011), p. 173.
- [55] D. C. Johnston, Cycloidal Paths in Physics as Superpositions of Translational and Rotational Motions, *Am. J. Phys.* (in press 2019); arXiv:1809.03871. In this paper $\mathbf{H}_1$ is taken to be in the opposite direction to that used here in Eq. (7.47).

- [56] N. S. Sangeetha, E. Cuervo-Reyes, A. Pandey, and D. C. Johnston, EuCo₂P₂: A model molecular-field helical Heisenberg antiferromagnet, *Phys. Rev. B* **94**, 014422 (2016).
- [57] M. Reehuis, W. Jeitschko, M. H. Möller, and P. J. Brown, A neutron diffraction study of the magnetic structure of EuCo₂P₂, *J. Phys. Chem. Solids* **53**, 687 (1992).
- [58] D. C. Johnston, Influence of uniaxial single-ion anisotropy on the magnetic and thermal properties of Heisenberg antiferromagnets within unified molecular field theory, *Phys. Rev. B* **95**, 094421 (2017).
- [59] H. Ibach and H. Lüth, *Solid-State Physics—An Introduction to Principles of Materials Science*, 3rd Edition (Springer-Verlag, Berlin, 2003).
- [60] D. C. Johnston, Influence of classical anisotropy fields on the properties of Heisenberg antiferromagnets within unified molecular field theory, *Phys. Rev. B* **96**, 224428 (2017).
- [61] E. Fawcett, Spin-density-wave antiferromagnetism in chromium, *Rev. Mod. Phys.* **60**, 209 (1988).
- [62] P. Mohn, *Magnetism in the Solid State* (Springer-Verlag, Berlin, 2003), p. 78.
- [63] Y. Takahashi, *Spin Fluctuation Theory of Itinerant Electron Magnetism* (Springer-Verlag, Berlin, 2013).
- [64] N. S. Sangeetha, L.-L. Wang, A. V. Smirnov, V. Smetana, A.-V. Mudring, D. D. Johnson, M. A. Tanatar, R. Prozorov, and D. C. Johnston, Non-Fermi-liquid behaviors associated with a magnetic quantum-critical point in Sr(Co_{1-x}Ni_x)₂As₂ single crystals, arXiv:1907.08238.
- [65] P. Mohn, *Magnetism in the Solid State—An Introduction* (Springer-Verlag, Berlin, 2003), Ch. 12–14.
- [66] Y. Li, Z. Liu, Z. Xu, Y. Song, Y. Huang, D. Shen, N. Ma, A. Li, S. Chi, M. Frontzek, H. Cao, Q. Huang, W. Wang, Y. Xie, Y. Rong, D. P. Young, J. F. DiTusa, and P. Dai, Flat band magnetism and helical magnetic order in Ni-doped SrCo₂As₂, arXiv:1907.09342.
- [67] J. M. Wilde, A. Kreyssig, D. Vaknin, N. S. Sangeetha, B. Li, W. Tian, P. P. Orth, A. I. Goldman, D. C. Johnston, B. G. Ueland, and R. J. McQueeney, Helical magnetic ordering in Sr(Co_{1-x}Ni_x)₂As₂, J. M. Wilde, A. Kreyssig, D. Vaknin, N. S. Sangeetha, B. Li, W. Tian, P. P. Orth, A. I. Goldman, D. C. Johnston, B. G. Ueland, and R. J. McQueeney, arXiv:1907.11676.
- [68] A. Arrott, Criterion for Ferromagnetism from Observations of Magnetic Isotherms, *Phys. Rev.* **108**, 1394 (1957); A. S. Arrott, Equations of state along the road to Arrott's last plot, *J. Magn. Magn. Mater.* **322**, 1047 (2010).
- [69] N. S. Sangeetha, V. Smetana, A.-V. Mudring, and D. C. Johnston, Anomalous Composition-Induced Crossover in the Magnetic Properties of the Itinerant-Electron Antiferromagnet $\text{Ca}_{1-x}\text{Sr}_x\text{Co}_{2-y}\text{As}_2$, *Phys. Rev. Lett.* **119**, 257203 (2017).
- [70] W. Jayasekara, Y. Lee, A. Pandey, G. S. Tucker, A. Sapkota, J. Lamsal, S. Calder, D. L. Abernathy, J. L. Niedziela, B. N. Harmon, A. Kreyssig, D. Vaknin, D. C. Johnston, A. I. Goldman, and R. J. McQueeney, Stripe Antiferromagnetic Spin Fluctuations in SrCo₂As₂, *Phys. Rev. Lett.* **111**, 157001 (2013).
- [71] P. Rhodes and E. P. Wohlfarth, The Effective Curie-Weiss Constant of Ferromagnetic Metals and Alloys, *Proc. Roy. Soc. London, Ser. A* **273**, 247 (1963).
- [72] E. P. Wohlfarth, Very Weak Itinerant Ferromagnets; Application to ZrZn₂, *J. Appl. Phys.* **39**, 1061 (1968).
- [73] J. Schmalian, Failed Theories of Superconductivity, *Mod. Phys. Lett. B* **24**, 2679 (2010).
- [74] Y. Singh, A. Ellern, and D. C. Johnston, Magnetic, transport, and thermal properties of single crystals of the layered arsenide BaMn₂As₂, *Phys. Rev. B* **79**, 094519 (2009).
- [75] M. E. Fisher, Relation between the Specific Heat and Susceptibility of an Antiferromagnet, *Phil. Mag.* **7**, 1731 (1962).
- [76] D. C. Johnston, Normal-State Magnetic Properties of Single-Layer Cuprate High-Temperature Superconductors and Related Materials, in *Handbook of Magnetic Materials*, Vol. 10, ed. K. H. J. Buschow (Elsevier, Amsterdam, 1997), Ch. 1, pp. 1–237.
- [77] Reference [51], pp. 14–25.
- [78] D. C. Johnston, Magnetic structure and magnetization of *z*-axis helical Heisenberg antiferromagnets with XY anisotropy in high magnetic fields transverse to the helix axis at zero temperature, *Phys. Rev. B* **99**, 214438 (2019).
- [79] A. P. Drozdov, M. I. Erements, I. A. Troyan, V. Ksenofontov, and S. I. Shylin, Conventional superconductivity at 203 kelvin at high pressures in the sulfur hydride system, *Nature* **525**, 73 (2015).

- [80] A. P. Drozdov, P. P. Kong, V. S. Minkov, S. P. Besedin, M. A. Kuzovnikov, S. Mozaffari, L. Balicas, F. F. Balakirev, D. E. Graf, V. B. Prakapenka, E. Greenberg, D. A. Knyazev, M. Tkacz, and M. I. Erements, *Nature* **569**, 528 (2019).
- [81] N. W. Ashcroft, Metallic Hydrogen: A High-Temperature Superconductor?, *Phys. Rev. Lett.* **21**, 1748 (1968).
- [82] N. W. Ashcroft, Hydrogen Dominant Metallic Alloys: High Temperature Superconductors?, *Phys. Rev. Lett.* **92**, 187002 (2004).
- [83] G. W. Webb, F. Marsiglio, and J. E. Hirsch, Superconductivity in the elements, alloys and simple compounds, *Physica C* **514**, 17 (2015).
- [84] F. London and H. London, The Electromagnetic Equations of the Supraconductor, *Proc. Roy. Soc. London A* **149**, 71 (1935).
- [85] M. Tinkham, *Introduction to Superconductivity*, 2nd Edition (Dover, Mineola, NY, 1996).
- [86] B. T. Matthias, T. H. Geballe, S. Geller, and E. Corenzwit, Superconductivity of Nb₃Sn, *Phys. Rev.* **95**, 1435 (1954).
- [87] J. R. Gavaler, Superconductivity in Nb-Ge films above 22 K, *Appl. Phys. Lett.* **23**, 480 (1973).
- [88] J. R. Gavaler, M. A. Janocko, and C. K. Jones, Preparation and properties of high- T_c Nb-Ge films, *J. Appl. Phys.* **45**, 3009 (1974).
- [89] P. Drude, *Zur Elektronentheorie der Metalle*, *Annalen der Physik* **306**, 566 (1900).
- [90] L. Hoddeson and G. Baym, The development of the quantum-mechanical electron theory of metals: 1928–1933, *Rev. Mod. Phys.* **59**, 287 (1987).
- [91] G. Grosso and G. P. Parravicini, *Solid State Physics*, 2nd Edition (Elsevier, Amsterdam, 2014), pp. 769–771.
- [92] S. Borowitz, *Fundamentals of Quantum Mechanics* (Benjamin, New York, 1967), p. 198.
- [93] J. K. Hulm, J. E. Kunzler, and B. T. Matthias, *Physics Today* **34**, 1 (1981).
- [94] J. Nagamatsu, N. Nakagawa, T. Muranaka, Y. Zenitani, and J. Akimitsu, Superconductivity at 39 K in magnetism diboride, *Nature* **410**, 63 (2001).
- [95] S. L. Bud'ko, G. Lapertot, C. Petrovic, C. E. Cunningham, N. Anderson, and P. C. Canfield, Boron Isotope Effect in Superconducting MgB₂, *Phys. Rev. Lett.* **86**, 1877 (2001).
- [96] J. G. Bednora and K. A. Müller, Possible High T_c Superconductivity in the Ba-La-Cu-O System, *Z. Phys. B* **64**, 189 (1986).
- [97] D. C. Johnston, H. Prakash, W. H. Zachariasen, and R. Viswanathan, High Temperature Superconductivity in the Li-Ti-O Ternary System, *Mater. Res. Bull.* **8**, 777 (1973).
- [98] D. C. Johnston, Superconducting and Normal State Properties of Li_{1+x}Ti_{2-x}O₄ Spinel Compounds. I. Preparation, Crystallography, Superconducting Properties, Electrical Resistivity, Dielectric Behavior, Magnetic Susceptibility, *J. Low Temp. Phys.* **25**, 145 (1976).
- [99] R. W. McCallum, D. C. Johnston, C. A. Luengo, and M. B. Maple, Superconducting and Normal State Properties of Li_{1+x}Ti_{2-x}O₄ Spinel Compounds. II. Low Temperature Heat Capacity, *J. Low Temp. Phys.* **25**, 177 (1976).
- [100] A. W. Sleight, J. L. Gillson, and P. E. Bierstedt, High-Temperature Superconductivity in the BaPb_{1-x}Bi_xO₃ System, *Solid State Commun.* **17**, 27 (1975).
- [101] Y. Kamihara, T. Watanabe, M. Hirano, and H. Hosono, Iron-Based Layered Superconductor La[O_{1-x}F_x]FeAs ($x = 0.05-0.12$), *J. Am. Chem. Soc.* **130**, 3296 (2008).
- [102] D. J. Griffiths, *Introduction to Quantum Mechanics*, 2nd Edition (Pearson, India, 2015).
- [103] C. D. Cantell, Bound-State Energies of a Particle in a Finite Square Well: An Improved Graphical Solution, *Am. J. Phys.* **39**, 107 (1971).
- [104] D. W. L. Sprung, H. Wu, and J. Martorell, A new look at the square well potential, *Eur. J. Phys.* **13**, 21 (1992).
- [105] D. L. Aronstein and C. R. Stroud, Jr., General series solution for finite square-well energy levels for use in wave-packet studies, *Am. J. Phys.* **68**, 943 (2000).