

SEZNAM POUŽITÉ LITERATURY

- 1 Ainsworth BE, Haskell WL, Leon AS et al. Compendium of physical activities of energy costs of human physical activities. *Med Sci Sports Exerc* 1993; 25: 71–80.
- 2 American College of Sports Medicine. Position Stand: Exercise and Fluid Replacement. *Med Sci Sports Exerc* 1996; 28: i-vii.
- 3 American College of Sports Medicine. Roundtable: The physiological and health effects of oral creatine supplementation. *Med Sci Sports Exerc* 2000; 32: 706–717.
- 4 Anderson RA, Bryden NA, Polansky MM, Deuster PA. Exercise effects on chromium excretion of trained and untrained men consuming a constant diet. *J Appl Physiol* 1988; 64: 249.
- 5 Anderson RA. Chromium as an essential nutrient or humans. *Regul Toxicol Pharmacol* 1997; 26: S35.
- 6 Anthony JC, Anthony TG, Kimball SR. Signalling pathways involved in the translational control of protein synthesis in skeletal muscle by leucine. *J Nutri* 2001; 131: 856–860.
- 7 Antonio J, Sanders MS, Van Gammeren D. The effects of bovine colostrum supplementation on body composition and exercise performance in active men and women. *Nutrition* 2001; 17: 243–247.
- 8 Antonio J, Street C. Glutamine a potentially useful supplement for athletes. *Can J Appl Phys* 1999; 24: 1–14.
- 9 Applegate L. Booster Club. *Runner's World* 1991; 26: 29–30.
- 10 Applegate L. Fad diets and supplement use in athletics. *Sports Sci Exchange* 1988; 1: 1–4.
- 11 Applegate L. Protein Power. *Runner's World* 1992; 27: 22–23.
- 12 Arenas J, Huertas R, Campos Y et al. Effect of L-carnitine and the pyruvate dehydrogenase complex and carnitine palmitoyl transferase activities in muscle of endurance athletes. *FEB Letters* 1994; 341: 91–93. (Feder Europ Biochem Soc).
- 13 Armstrong L, Maresh C. Vitamin and mineral supplements as nutritional aids to exercise performance and health. *Nutrition Reviews* 1996; 54(Suppl.): S148–S158.
- 14 Aronson V. Protein and miscellaneous ergogenic aids. *Physician Sportsmed* 1986; 14: 199–202.

- 15 Åstrand PO. Something old and something new... very new. *Nutr Today* 1968; 3: 9–11.
- 16 Baguet A, Koppo K, Pottier A et al. Beta-alanine supplementation reduces acidosis but not oxygen uptake response during high-intensity cycling exercise. *Eur J Appl Physiol* 2010; 108: 495–503.
- 17 Balasse EO, Fery F, Neef MA. Changes induced by exercise in rates of turnover and oxidation of ketone bodies in fasting man. *J Appl Physiol* 1978; 44: 5–11.
- 18 Banister EW, Cameron BJ. Exercise-induced hyperammonemia: peripheral and central effects. *Int J Sports Med* 1990; 11(Suppl 2): S129–S142.
- 19 Barrarre TL, Scarpino A, Sigmon R et al. Vitamin-mineral supplement use and nutritional status of athletes. *J Am Coll Nutr* 1993; 12: 162–169.
- 20 Benardot D et al. Can vitamin supplements improve sport performance? *Sports Science Exchange Roundtable* 2001; 12: 1–4.
- 21 Bennett WM, Connacher AA, Scrimgeour CM et al. The effect of amino-acid infusion on leg protein turnover assessed by l-[15N]phenylalanine and l-[1-13C]leucine exchange. *Eur J Clin Invest* 1990; 20: 37–46.
- 22 Bergström J, Hermansen L, Hultman E et al. Diet, muscle glycogen and physical performance. *Acta Physiol Scand* 1967; 71: 140–150.
- 23 Biolo G, Tipton KD, Klein S, Wolfe RR. An abundant supply of amino acids enhances the metabolic effect of exercise on muscle protein synthesis. *Am J Physiol* 1997; 273: E122–E129.
- 24 Bishop D, Edge J, Davis C et al. Induced metabolic alkalosis affects muscle metabolism and repeated-sprint ability. *Med Sci Sports Exerc* 2004; 36: 807–813.
- 25 Bohe J, Low A, Wolfe RR et al. Human muscle protein synthesis is modulated by extracellular not intramuscular amino acid availability: a dose response study. *J Physiol* 2003; 552: 315–324.
- 26 Borsheim E, Tipton KD, Wolf SE, Wolfe RR. Essential amino acids and muscle protein recovery from resistance exercise. *Am J Physiol Endocrinol Metab* 2002; 283: E648–E657.
- 27 Brass PE, Scarrow AM, Ruff JL et al. Carnitine delays rat skeletal muscle fatigue in vitro. *J Appl Physiol* 1993; 75: 1595–1600.
- 28 Braun B, Clarkson PM, Freedson PS et al. Effects of coenzyme Q10 supplementation on exercise performance, VO₂max, and lipid peroxidation in trained cyclists. *Int J Sport Nutr* 1991; 1: 353–365.
- 29 Brilla LR, Gunther KB. Effect of magnesium supplementation on exercise time to exhaustion. *Med Exerc Nutr Health* 1995; 4: 230.
- 30 Brilla LR, Haley TF. Effect of magnesium supplementation on strength training in humans. *J Am Coll Nutr* 1992; 1: 326.
- 31 Brinkworth GD, Buckley JD, Slavotinek JP et al. Effect of bovine colostrum supplementation on the composition of resistance trained and untrained limbs in healthy young men. *Eur J Appl Physiol* 2004; 91: 53–60.
- 32 Brounsa F, van der Vusse GJ. Utilization of lipids during exercise in human subjects: metabolic and dietary constraints. *Brit J Nutr* 1998; 79: 117–128.

- 33 Brutsaert TD, Hernandez-Cordero S, Rivera J et al. Iron supplementation improves progressive fatigue resistance during dynamic knee extensor exercise in iron-depleted, nonanemic women. *Am J Clin Nutr* 2003; 77: 441.
- 34 Bucci LR, Unlu L. Proteins and amino acids in exercise and sport. In Driskell J, Wolinsky I (ed.), *Energy-Yielding Macronutrients and Energy Metabolism in Sports Nutrition*. Boca Raton : CRC Press, 2000, p. 197–200.
- 35 Buckley JD, Abbott MJ, Brinkworth GD et al. Bovine colostrum supplementation during endurance running training improves recovery, but not performance. *J Sci Med Sport* 2002; 5: 65–79.
- 36 Buckley JD, Brinkworth GD, Abbott MJ. Effect of bovine colostrum on anaerobic exercise performance and plasma IGF-I. *J Sports Sci* 2003; 21: 577–588.
- 37 Buckspan R, Hoxworth B, Cersosimo E et al. Alpha-Ketoisocaproate is superior to leucine in sparing glucose utilization in humans. *Am J Physiol* 1986; 251: E648–E653.
- 38 Bucsi L, Poór G. Efficacy and tolerability of oral chondroitin sulfate as a symptomatic slow-acting drug for osteoarthritis (SYSADOA) in the treatment of knee osteoarthritis. *Osteoarthritis Cartilage* 1998; 6(Suppl A): 31–36.
- 39 Budgett R et al. The overtraining syndrome. In Harries M et al. (ed.), *Oxford Textbook of Sports Medicine*. Oxford : Oxford University Press, 1998.
- 40 Buford BN, Koch AJ. Glycine-arginine-alfa-ketoisocaproic acid improves performance of repeated cycling sprints. *Med Sci Sports Exerc* 2004; 36: 583–587.
- 41 Buford TW, Kreider RB, Stout JR et al. International Society of Sports Nutrition position stand: creatine supplementation and exercise. *Journal of the International Society of Sports Nutrition* 2007; 4: 6. <http://www.jissn.com/content/4/1/6>.
- 42 Bulow J. Lipid metabolism and utilization. In Poortmans J. *Principles of Exercise Biochemistry*. Switzerland, Basel : Karger, 1993.
- 43 Burke DG, Chilibeck PD, Davidson KS et al. The effect of whey protein supplementation with and without creatine monohydrate combined with resistance training on lean tissue mass and muscle strength. *Int J Sport Nutr Exerc Metab* 2001; 11: 349–364.
- 44 Burke L, Desbrow B, Minehan M. Dietary supplements and nutritional ergogenic aids in sport. In Burke L, Deakin V. *Clinical Sports Nutrition*. Australia, Sydney : McGraw-Hill, 2000, p. 455–553.
- 45 Buse MG, Biggers JF, Friderici KH et al. Oxidation of branched chain amino acids by isolated hearts and diaphragms of the rat. The effect of fatty acids, glucose, and pyruvate respiration. *J Biol Chem* 1972; 247: 8085–8096.
- 46 Byars A, Keith S, Simpson W, Mooneyhan A et al. The influence of a pre-exercise sports drink (PRX) on factors related to maximal aerobic performance. *J Int Soc Sports Nutr* 2010; 7: 12.
- 47 Cairns SP. Lactic acid and exercise performance. *Sports Med* 2006; 36: 279–289.
- 48 Campbell B, Roberts M, Kerksick Ch et al. Pharmacokinetics, safety, and effects on exercise performance of L-arginine alfa-ketoglutarate in trained adult men. *Nutrition* 2006; 22: 872–881.

- 49 Casey A, Constantin-Teodosiu E, Howell S et al. Creatine ingestion favorably affects performance and muscle metabolism during maximal exercise in humans. *Am J Physiol* 1996; 271: E31–E37.
- 50 Castell L et al. Does glutamine have a role in reducing infections in athletes? *Eur J Appl Physiol* 1996; 73: 488–490.
- 51 Castell L. Glutamine supplementation *vitro* and *in vivo*, in exercise and in immunodepression. *Sports Med* 2003; 33: 323–345.
- 52 Centers for Disease Control and Prevention. Recommendations to prevent and control iron deficiency in the United States. *MMWR* 1998; 47: 1.
- 53 Clancy SP, Clarkson PM, DeCheke ME et al. Effects of chromium supplementation on body composition, strength and urinary chromium loss in football players. *Int J Sports Nutr* 1994; 4: 142.
- 54 Clarkson PM. Antioxidants and physical performance. *Crit Rev Food Sci Nutr* 1995; 35: 131–141.
- 55 Clegg DO, Reda DJ, Harris CL et al. Glucosamine, chondroitin sulfate, and the two in combination for painful knee osteoarthritis. *N Engl J Med* 2006; 354: 795–808.
- 56 Colakoglu S, Colakoglu M, Taneli F et al. Cumulative effects of conjugated linoleic acid and exercise on endurance development, body composition, serum leptin and insulin levels. *J Sports Med Phys Fitness* 2006; 46: 570–577.
- 57 Cooke M, Iosia M, Buford T et al. Effects of acute and 14-day coenzyme Q10 supplementation on exercise performance in both trained and untrained individuals. *J Int Soc Sports Nutr* 2008; 5: 14.
- 58 Coombes J, Conacher M, Austen S et al. Dose effects of oral bovine colostrum supplementation on physical work capacity in cyclists. *Med Sci Sports Exerc* 2002; 34: 1184–1188.
- 59 Coombes JS, McNaughton LR. Effects of branched-chain amino acid supplementation on serum creatine kinase and lactate dehydrogenase after prolonged exercise. *J Sports Med Phys Fitness* 2000; 40: 240–246.
- 60 Cornish SM, Candow DG, Jantz NT et al. Conjugated linoleic acid combined with creatine monohydrate and whey protein supplementation during strength training. *Int J Sport Nutr Exerc Metab* 2009; 19: 79–96.
- 61 Cribb PJ, Williams AD, Hayes A et al. The effect of whey isolate on strength, body composition and plasma glutamine. *Med Sci Sports Exerc* 2002; 34: A1688.
- 62 Cribb PJ, Williams AD, Hayes A, Carey MF. The effects of whey isolate and creatine on muscular strength, body composition and muscle fiber characteristics. *FASEB J* 2003; 17: 592. <http://www.the-aps.org/press/conference/eb03/12.htm>.
- 63 Crittenden R, Buckley J, Cameron-Smith D et al. Functional dairy protein supplements for elite athletes. *Aust J Dairy Technol* 2009; 64: 133–137.
- 64 Crowe MJ, Weatherson JN, Bowden BF. Effects of dietary leucine supplementation on exercise performance. *Eur J Appl Physiol* 2006; 97: 664–672.

- 65 Dalbo VJ, Roberts MD, Stout JR, Kerksick CM. Putting to rest the myth of creatine supplementation leading to muscle cramps and dehydration. *Br J Sports Med* 2008; 42: 567–573.
- 66 Dangin M, Guillet C, Garcia-Rodenas C et al. The rate of protein digestion affects protein gain differently during aging in humans. *J Physiol* 2003; 549: 635–644.
- 67 Dangott B, Schultz E, Mozdziak PE. Dietary creatine monohydrate supplementation increases satellite cell mitotic activity during compensatory hypertrophy. *Int J Sports Med* 2000; 21: 13–16.
- 68 Darden E. Olympic athletes view vitamins and victories. *J Home Economics* 1973; 65: 8–11.
- 69 Davis JM et al. Serotonin and central nervous system fatigue: Nutritional Considerations. *Am J Clin Nutr* 2000; 72: 573S–578S.
- 70 Dawson B, Cutler M, Moody A et al. Effects of oral creatine loading on single and repeated maximal short sprints. *Aust J Sci Med Sport* 1995; 27: 56–61.
- 71 de Deckere EA, van Amelsvoort JM, McNeill GP et al. Effects of conjugated linoleic acid (CLA) isomers on lipid levels and peroxisome proliferation in the hamster. *Br J Nutr* 1999; 82: 309–317.
- 72 De Palo EF, Gatti R, Cappellin E et al. Plasma lactate, GH and GH-binding protein levels in exercise following BCAA supplementation in athletes. *Amino Acids* 2001; 20: 1–11.
- 73 Dekkers J et al. The role of antioxidant vitamins and enzymes in the prevention of exercise-induced muscled damage. *Sports Med* 1996; 21: 213–238.
- 74 Dempsey RL, Mazzone MF, Meurer LN. Does oral creatine supplementation improve strength? A meta-analysis. *J Fam Prac* 2002; 51: 11.
- 75 Demura S, Yamada T, Yamaji S et al. The effect of L-ornithine hydrochloride ingestion on human growth hormone secretion after strength training. *Adv Biosci Biotechnol* 2010; 1: 7–11.
- 76 Derave W, Everaert I, Beeckman S et al. Muscle carnosine metabolism and beta-alanine supplementation in relation to exercise and training. *Sports Med* 2010; 40: 247–263.
- 77 Deuster PA, Day BA, Singh A et al. Zinc status of highly trained women runners and untrained women. *Am J Clin Nutr* 1989; 49: 1295.
- 78 Dragan AM, Vasiliu D, Eremia NM et al. Studies concerning some acute biological changes after endovenous administration of 1 g L-carnitine, in elite athletes. *Physiologie* 1987; 24: 231.
- 79 Duquet W, Carter JEL. Somatotyping. In Eston R, Reilly T (ed.), *Kinanthropometry and exercise physiology laboratory manual*. London : Chapman and Hall, 1996.
- 80 Evans GW. The effect of chromium picolinate on insulin-controlled parameters in humans. *Int J Biosoci Med Res* 1989; 11: 163.
- 81 Evans W. Vitamin E, vitamin C, and exercise. *Am J Clin Nutr* 2000; 72(Suppl): 647–652.
- 82 Foehrenbach R, Maerz W, Lohrerer H et al. Der Einfluss von L-Carnitin auf den Lipidstoffwechsel von Hochleistungssportlern. *Dtsch Z Sportmed* 1993; 44: 349–356.

- 83 Food and Nutrition Board, Institute of Medicine. Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. Washington, DC : National Academy Press, 2001.
- 84 Fořt P. Sport a správná výživa. Praha : Euromedia Group – Ikar, 2002.
- 85 Fořt P. Výživa v otázkách a odpovědích. Praha : Svět kulturistiky, 2003.
- 86 Fořt P. Zdraví a potravní doplňky. Praha : Euromedia Group – Ikar, 2005.
- 87 Friedman S, Fraenkel G. Reversible enzymatic acetylation of carnitine. *Arch Biochem Biophys* 1955; 59: 491–501.
- 88 Fryburg DA, Jahn LA, Hill SA et al. Insulin and insulin-like growth factor-I enhance human skeletal muscle protein anabolism during hyperaminoacidemia by different mechanisms. *J Clin Invest* 1995; 96: 1722–1729.
- 89 Fushiki T, Matsumoto K, Inoue K et al. Swimming capacity of mice is increased by chronic consumption of medium-chain triacylglycerols. *J Nutr* 1995; 125: 531–539.
- 90 Gao J, Costill D, Horswill C, Park SH. Sodium bicarbonate ingestion improves performance in interval swimming. *Eur J Appl Physiol Occup Physiol* 1988; 58: 171–174.
- 91 Gaullier JM, Halse J, Høivik HO et al. Six months supplementation with conjugated linoleic acid (CLA) induces regional-specific fat mass decreases in overweight and obese. *Br J Nutr* 2007; 97: 50–60.
- 92 Gaullier JM, Halse J, Høye K et al. Conjugated linoleic acid supplementation for 1 year reduces body fat mass in healthy overweight humans. *Am J Clin Nutr* 2004; 79: 1118–1125.
- 93 Gavino VC, Gavino G, Leblanc MJ et al. An isomeric mixture of conjugated linoleic acids but not pure cis-9, trans-11-octadecadienoic acid affects body weight gain and plasma lipids in hamsters. *J Nutr* 2000; 130: 27–29.
- 94 Giamberardino MA, Dragani L, Valente R et al. Effects of prolonged L-carnitine administration on delayed muscle pain and CK release after eccentric effort. *Int J Sports Med* 1996; 17: 320.
- 95 Gibala M. Dietary protein, amino acid supplements, and recovery from exercise. *Sports Sci Exchange* 2002; 15: 1–4.
- 96 Goedecke JH, Elmer-English R, Dennis SC et al. Effects of medium-chain triacylglycerol ingested with carbohydrate on metabolism and exercise performance. *Int J Sport Nutr* 1999; 9: 35–47.
- 97 Gökbel H, Gül I, Belviranl M et al. The effects of coenzyme Q10 supplementation on performance during repeated bouts of supramaximal exercise in sedentary men. *J Strength Cond Res* 2010; 24: 97–102.
- 98 Goldfinch J, McNaughton L, Davies P. Induced metabolic alkalosis and its effects on 400 m racing time. *Eur J Appl Physiol Occup* 1988; 57: 45–48.
- 99 Golf SW, Bohmer D, Nowacki PE. Is magnesium a limiting factor in competitive exercise? A summary of relevant scientific data. In Golf S, Dralle D, Vecchiet L (ed.), *Magnesium*. London : John Libbey, 1993.

- 100 Gordon B, Kohn LA, Levine SA et al. Sugar content of the blood in runners following a marathon race. *J Am Med Assoc* 1925; 85: 508–509.
- 101 Gorostiaga E, Maurer C, Eklache J. Decrease in respiratory quotient during exercise following L-carnitine supplementation. *Int J Sports Med* 1989; 10: 169–174.
- 102 Graham TE. Caffeine and exercise: metabolism, endurance, performance. *Sports Med* 2001; 31: 765–807.
- 103 Grant KE, Chandler RM, Castle AL, Ivy JL. Chromium and exercise training: effect on obese women. *Med Sci Sports Exerc* 1997; 29: 992.
- 104 Greenhaff PL, Bodin K, Söderlund K et al. Effect of oral creatine supplementation on skeletal muscle phosphocreatine resynthesis. *Am J Physiol* 1994; 266: E725–E730.
- 105 Greenwood M, Kreider RB, Melton C et al. Creatine supplementation during college football training does not increase the incidence of cramping or injury. *Mol Cell Biochem* 2003; 244: 83–88.
- 106 Greenwood M, Kreider RB, Greenwood L, Byars A. Cramping and injury incidence in collegiate football players are reduced by creatine supplementation. *J Athl Train* 2003; 38: 216–219.
- 107 Greenwood M, Kreider RB, Greenwood L, Byars A. The effects of creatine supplementation on cramping and injury occurrence during college baseball training and competition. *J Exerc Physiol Online* 2003; 6: 16–23.
- 108 Groneberg DA, Kindermann B, Althammer M et al. Coenzyme Q10 affects expression of genes involved in cell signalling, metabolism and transport in human CaCo-2 cells. *Int J Biochem Cell Biol* 2005; 37: 1208–1218.
- 109 Grunewald KK, Bailey RS. Commercially marketed supplements for bodybuilding athletes. *Sports Med* 1993; 15: 90–103.
- 110 Gudmundsen O. Conjugated linoleic acid reduces body fat mass in overweight and obese humans. *J Nutr* 2000; 130: 2943–2948.
- 111 Ha E, Zemel MB. Functional properties of whey, whey components, and essential amino acids: mechanisms underlying health benefits for active people. *J Nutr Biochem* 2003; 14: 251–258.
- 112 Hallmark MA, Reynolds TH, DeSouza CA et al. Effects of chromium and resistive training on muscle strength and body composition. *Med Sci Sports Exerc* 1996; 28: 139.
- 113 Hargreaves M, Snow R. Amino acids and endurance exercise. *Int J Sport Nutr Exerc Metabol* 2001; 11: 133–145.
- 114 Harris RC, Söderlund K, Hultman E. Elevation of creatine in resting and exercised muscle of normal subjects by creatine supplementation. *Clin Sci (Colch.)* 1992; 83: 367–374.
- 115 Harris RC, Tallon MJ, Dunnett M et al. The absorption of orally supplied beta-alanine and its effect on muscle carnosine synthesis in human vastus lateralis. *Amino Acids* 2006; 30: 279–289.
- 116 Harrison A, Thompson K. Ergogenic aids: sodium bicarbonate. *Peak Performance* 2005; 219: 9–10.

- 117 Haskell WL. The influence of exercise training on plasma lipids and lipoproteins in health and disease. *Acta Med Scand* 1986; 711(Suppl): 25–37.
- 118 Hasten DL, Rome EP, Franks DB, Hegsted M. Effects of chromium picolinate on beginning weight training students. *Int J Sports Nutr* 1990; 2: 343.
- 119 Häussinger D, Roth E, Lang F et al. Cellular hydration state: An important determinant of protein catabolism in health and disease. *Lancet* 1993; 341: 1330–1332.
- 120 Haymes EM, Spillman DM. Iron status of women distance runners. *Int J Sports Med* 1989; 10: 430.
- 121 Haymes EM. Trace minerals and exercise. In Wolinsky I (ed.), *Nutrition in exercise and spor.* 3rd ed. Boca Raton : CRC Press, 1998.
- 122 Heath BH, Carter JEL. A modified somatotype method. *Am J Physical Anthropol* 1967; 27: 57–74.
- 123 Hendler SS, Rorvik D. *PDR for nutritional supplements.* Montvale : PDR Thompson, 2001.
- 124 Herrero-Beaumont G, Ivorra JA, Del Carmen Trabada M et al. Glucosamine sulfate in the treatment of knee osteoarthritis symptoms: A randomized, double-blind, placebo-controlled study using acetaminophen as a side comparator. *Arthritis Rheum* 2007; 56: 555–567.
- 125 Hespel P, Opteijnde B, Van Leemputte M et al. Oral creatine supplementation facilitates the rehabilitation of disuse atrophy and alters the expression of muscle myogenic factors in humans. *J Physiol* 2001; 536: 625–633.
- 126 Hinton PS, Giordano C, Brownlie T, Haas JD. Iron supplementation improves endurance after training in iron-depleted, nonanemic women. *J Appl Physiol* 2000; 88: 1103.
- 127 Hofman Z, Smeets R, Verlaan G et al. The effect of bovine colostrum supplementation on exercise performance in elite field hockey players. *Int J Sport Nutr Exerc Metab* 2002; 12: 461–469.
- 128 Hogervorst EEF, Bandelow S, Schmitt J et al. Caffeine Improves Physical and Cognitive Performance during Exhaustive Exercise *Medicine Science in Sports Exercise* 2008; 40: 1841–1851.
- 129 Hochberg MC, Zhan M, Langenberg P. The rate of decline of joint space width in patients with osteoarthritis of the knee: a systematic review and meta-analysis of randomized placebo-controlled trials of chondroitinsulfate. *Curr Med Res Opin* 2008; 24: 3029–3035.
- 130 Hochberg MC. Structure-modifying effects of chondroitinsulfate in knee osteoarthritis: an updated meta-analysis of randomized placebo-controlled trials of 2-year duration. *Osteoarthritis and Cartilage* 2010; 18: S28–S31.
- 131 Horstman DH. Nutrition. In Morgan WP (ed.), *Ergogenic Aids and Muscular Performance.* New York: Academic Press, 1972, p. 343–365.
- 132 Hotell MD, Faria IE. Carbohydrate paste ingestion during prolonged treadmill running. *Med Sci Sports Exercise* 1996; 28: S38 (abstr.).
- 133 Huertas R, Campos Y, Diaz E et al. Respiratory chain enzymes in muscle of endurance athletes: effect of L-carnitine. *Biochem Biophys Res Commun* 1992; 188: 102.

- 134 Chen S, Kim W, Henning SM et al. Arginine and antioxidant supplement on performance in elderly male cyclists: a randomized controlled trial. *J Int Soc Sports Nutr* 2010; 7: 13.
- 135 Cheng W, Phillips B, Abumrad N. Effect of HMB on fuel utilization, membrane stability and creatine kinase content of cultured muscle cells. *FASEB J* 1998; 12: A950 (abstr.).
- 136 Chevront S et al. Branched-chain amino acid supplementation and human performance when hypohydrated in the heat. *J Appl Physiol* 2004; 97: 1275–1282.
- 137 Chromiak J, Antonio J. Use of amino acids as growth-hormone releasing agents by athletes. *Nutrition* 2002; 18: 657–661.
- 138 Chytráčková J. Rozdělení somatografu na oblasti podle výkonnosti. In Riegerová J, Ulbrichová M. *Aplikace fyzické antropologie v tělesné výchově a sportu*. Olomouc : Univerzita Palackého, 1993.
- 139 Ichihara A, Noda C, Ogawa K. Control of leucine metabolism with special reference to branched-chain amino acid transaminase isozymes. *Adv Enzyme Regul* 1973; 11: 155–166.
- 140 Isidori A, Monaco AL, Cappa L. A study of growth hormone release in man after oral administration of amino acids. *Curr Med Res Opin* 1981; 7: 475–481.
- 141 Jacobs PL, Goldstein ER. Long-term glycine propionyl-L-carnitine supplementation and paradoxical effects on repeated anaerobic sprint performance. *J Int Soc Sports Nutr* 2010; 7: 35.
- 142 Jacobson B et al. Nutrition practices and knowledge of college varsity athletes: A follow-up. *J Strength Conditioning Res* 2001; 15: 63–68.
- 143 Jefferson LS, Kimball SR. Amino acid regulation of gene expression. *J Nutr* 2001; 131: 2460S–2466S.
- 144 Jeukendrup AE, Thielen JJ, Wagenmakers AJM et al. Effects of medium-chain triacylglycerol and carbohydrate ingestion during exercise on substrate utilization and subsequent cycling performance. *Am J Clin Nutr* 1998; 67: 397–404.
- 145 Ji L. Antioxidants and oxidative stress in exercise. *Proceedings of the Society for Experimental Biology and Medicine* 1999; 222: 283–292.
- 146 Simopoulos AS. The importance of the Omega-6/omega-3 fatty acid ratio in cardiovascular disease and other chronic diseases. *Exp Biol Med* 2000; 233: 674–688.
- 147 Jones AM, Atter T, Georg KP. Oral creatine supplementation improves multiple sprint performance in elite ice-hockey players. *J Sports Med Phys Fitness* 1999; 39: 189–196.
- 148 Jowko E, Ostaszewski P, Jank M. Creatine and beta-hydroxy-beta-methylbutyrate (HMB) additively increase lean body mass and muscle strength during a weighttraining program. *Nutrition* 2001; 17: 558–566.
- 149 Joyner MJ. Glutamine and arginine: immunonutrients and metabolic modulators? *Exerc Sport Sci Rev* 2005; 33: 105–106.
- 150 Kaats GB, Blum K, Fisher JA, Adelman JA. Effects of chromium picolinate supplementation on body composition: a randomized, double-masked, placebo controlled study. *Curr Ther Res* 1996; 57: 747.

- 151 Kahan A, Uebelhart D, DeVathaire F et al. Long-term effects of chondroitins 4 and 6 sulfate on knee osteoarthritis: the study on osteoarthritis progression prevention, a two-year, randomized, double-blind, placebo-controlled trial. *Arthritis Rheum* 2009; 60: 24–33.
- 152 Kajaba I, Budlovský J, Dvorský I et al. Nové odporúčané výživové dávky pre obyvateľstvo ČSFR. *Čas Lék čes* 1992; 131: 198–204.
- 153 Karlic H, Lohninger A. Supplementation of L-carnitine in athletes: does it make sense? *Nutrition* 2004; 20: 709–715.
- 154 Karlsson HK, Nilsson PA, Nilsson J et al. Branched-chain amino acids increase p70S6K phosphorylation in human skeletal muscle after resistance exercise. *Am J Physiol Endocrinol Metab* 2004; 287: E1–E7.
- 155 Kerksick CM, Rasmussen C, Lancaster S et al. Impact of differing protein sources and a creatine containing nutritional formula after 12 weeks of resistance training. *Nutr* 2007; 23: 647–656.
- 156 Kimball SR, Jefferson LS. Regulation of global and specific mRNA translation by oral administration of branched-chain amino acids. *Biochem Biophys Res Commun* 2004; 313: 423–427.
- 157 Kleine MV. Evidence of the efficacy of an enzyme combination preparation using the method of artificial hematomas in combination with a pressure meter: A placebo controlled, randomised, prospective, double-blind study. *J Clin Res* 1998; 1: 87–102.
- 158 Kon M, Tanabe K, Akimoto T et al. Reducing exercise-induced muscular injury in kendo athletes with supplementation of coenzyme Q10. *Br J Nutr* 2008; 100: 903–909.
- 159 Koopman R, Wagenmakers AJM, Ralph JF et al. Combined ingestion of protein and free leucine with carbohydrate increases postexercise muscle protein synthesis in vivo in male subjects. *Am Physiol Endocrinol Metab* 2005; 288: E645–E653.
- 160 Kormasio R, Riederer I, Butler-Browne G et al. Beta-hydroxy-beta-methylbutyrate (HMB) stimulates myogenic cell proliferation, differentiation and survival via the MAPK/ERK and PI3K/Akt pathways. *Biochim Biophys Acta* 2009; 1793: 755–763.
- 161 Kraemer WJ, Volek JS, French DN et al. The effects of L-carnitine L-tartrate supplementation on hormonal responses to resistance exercise and recovery. *J Strength Cond Res* 2003; 17: 455.
- 162 Kraut H, Müller EA, Müller-Wecker H. Die Abhängigkeit des Muskeltrainings und Eiweiss-bestand des Körpers. *Biochem Z* 1953; 324: 280–294.
- 163 Kreider RB, Ferreira MP, Greenwood M et al. Effects of conjugated linoleic acid supplementation during resistance training on body composition, bone density, strength, and selected hematological markers. *J Strength Cond Res* 2002; 16: 325–334.
- 164 Kreider RB, Melton C, Rasmussen CJ et al. Long-term creatine supplementation does not significantly affect clinical markers of health in athletes. *Mol Cell Biochem* 2003; 244: 95–104.

- 165 Lambert EV, Goedecke JH, Bluett K et al. Conjugated linoleic acid versus high oleic acid sunflower oil: effects on energy metabolism, glucose tolerance, blood lipids, appetite and body composition in regularly exercising individuals. *Br J Nutr* 2007; 97: 1001–1011.
- 166 Lambert EV, Goedecke JH, van Zyl C et al. High-fat diet versus habitual diet prior to carbohydrate loading: effects on exercise metabolism and cycling performance. *Internat J Sport Nutr Exerc Metabol* 2001; 11: 209–225.
- 167 Matiegka, J. The testing of physical efficiency. *Am J Phys Anthropol*, 1921, 4: 223–230.
- 168 Lambert EV, Speechly DP, Dennis SC, Noakes TD. Enhanced endurance in trained cyclists during moderate intensity exercise following 2 weeks adaptation to a high fat diet. *Eur J Appl Physiol* 1994; 69: 287–293.
- 169 Lambert M et al. Failure of commercial oral amino acid supplements to increase serum growth hormone concentrations in male body builders. *Int J Sport Nutr* 1993; 3: 298–305.
- 170 Lands LC, Grey VL, Smountas AA. Effect of supplementation with a cysteine donor on muscular performance. *J Appl Physiol* 1999; 87: 1381–1385.
- 171 Lang CH. Elevated plasma free fatty acids decrease basal protein synthesis, but not the anabolic effect of leucine, in skeletal muscle. *Am J Physiol Endocrin Metab* 2006; 291: E666–E674.
- 172 Layman DK. Role of leucine in protein metabolism during exercise and recovery. *Can J Appl Physiol* 2002; 27: 646–663.
- 173 Leeb BF, Schweizer H, Montag K, Smolen JS. A meta-analysis of chondroitin sulphate in the treatment of osteoarthritis. *J Rheumatol* 2000; 27: 205–211.
- 174 Leelarungrayub D, Sawattikanon N, Klaphajone J et al. Coenzyme Q10 Supplementation Decreases Oxidative Stress and Improves Physical Performance in Young Swimmers: A Pilot Study. *Open Sports Medicine Journal* 2010; 4: 1–8.
- 175 Leibowitz BE. L-Carnitine. Basel : Lonza, 1993.
- 176 Levine SA, Gordon B, Derick CL. Some changes in the chemical constituents of the blood following a marathon race. *J Am Med Assoc* 1924; 82: 1778–1779.
- 177 Littarru GP, Tiano L. Clinical aspects of coenzyme Q10: An update. *Nutrition* 2010; 26: 250–254.
- 178 Little JP, Forbes SC, Candow DG et al. Creatine, arginine α -ketoglutarate, amino acids, and medium-chain triglycerides and endurance performance. *Int J Sport Nutr Exerc Metabol* 2008; 18: 493–508.
- 179 Liu Z, Jahn LA, Long W et al. Branched chain amino acids activate messenger ribonucleic acid translation regulatory proteins in human skeletal muscle, and glucocorticoids blunt this action. *J Clin Endocrinol Metab* 2001; 86: 2136–2143.
- 180 Liu Z, Jahn LA, Wei L et al. Amino Acids Stimulate Translation Initiation and Protein Synthesis through an Akt-Independent Pathway in Human Skeletal Muscle. *J Clin Endocrinol & Metabol* 2002; 87: 5553–5558.

- 181 Lukaski HC. Chromium as a supplement. In McCormick DB, Bier DM, Goodridge AG, Parmer R (ed.), *Annual Review of Nutrition*. Palo Alto : Annual Reviews, 1999.
- 182 Lukaski HC. Prevention and treatment of magnesium deficiency in athletes. In Vecchiet L (ed.), *Magnesium and Physical Activity*. London : Parthenon, 1995.
- 183 Lukaski HC. Vitamin and mineral status: effects on physical performance. *Nutrition* 2004; 20: 632–644.
- 184 Lukaski HC. Zinc. In Wolinsky I, Driskell JA (ed.), *Sports Nutrition: Vitamins and Trace Elements*. Boca Raton : CRC Press, 1997.
- 185 Máček M, Radvanský J et al. *Fyziologie a klinické aspekty pohybové aktivity*. Praha : Galén, 2011, 245 s.
- 186 Malm C, Svensson M, Ekblom B et al. Effects of ubiquinone-10 supplementation and high intensity training on physical performance in humans. *Acta Physiol Scand* 1997; 161: 379–384.
- 187 Mandelová L, Hrnčíříková I. *Základy výživy ve sportu*. Brno : Masarykova Univerzita, 2007.
- 188 Manore M. Vitamins and minerals: Part I. How much do I need? *ACSM's Health & Fitness Journal* 2001; 5: 30–33.
- 189 Manore M. Vitamins and minerals: Part II. Who needs to supplement? *ACSM's Health & Fitness Journal* 2001; 5: 34–35.
- 190 Marconi C, Sassi G, Carpinelli A et al. Effect of carnitine loading on aerobic and anaerobic performance of endurance athletes. *Eur J Appl Physiol* 1985; 54: 131–135.
- 191 Mariotti F, Simbelie KL, Makarios-Lahham L et al. Acute ingestion of dietary proteins improves post-exercise liver glutathione in rats in a dose-dependent relationship with their cysteine content. *J Nutr* 2004; 134: 128–131.
- 192 Matsumoto K, Koba T, Hamada K et al. Branched-chain amino acid supplementation increases the lactate threshold during an incremental exercise test in trained individuals. *J Nutr Sci Vitaminol* 2009; 55: 55–58.
- 193 Matthews DE, Bier DM, Rennie MJ et al. Regulation of leucine metabolism in man: a stable isotope study. *Science* 1981; 214: 1129–1131.
- 194 Matthews DE, Harkin R, Battezzati A, Brillon DJ. Splanchnic bed utilization of enteral alpha-ketoisocaproate in humans. *Metabolism* 1999; 48: 1555–1563.
- 195 Maughan J, Burke LM. *Výživa ve sportu*. Praha : Galen, 2006.
- 196 Max SR. Glucocorticoid-mediated induction of glutamine synthetase in skeletal muscle. *Med Sci Sports Exerc* 1990; 22: 325–330.
- 197 Mayer J, Bullen B. Nutrition and athletic performance. *Physiol Rev* 1960; 40: 369–397.
- 198 Mazieres B, Hucher M, Zaim M, Garnerio P. Effect of chondroitin sulfate in symptomatic knee osteoarthritis: A multicentre, randomised, double-blind, placebo-controlled study. *Ann Rheum Dis* 2007; 66: 639–645.
- 199 McAlindon TE, LaValley MP, Gulin JP, Felson DT. Glucosamine and chondroitin for treatment of osteoarthritis: a systematic quality assessment and meta-analysis. *JAMA* 2000; 283: 1469–1475.

- 200 McNaughton LR, Cedaro R. The effect of sodium bicarbonate on rowing ergometer performance in elite rowers. *Austr J Sci Med Sport* 1991; 23: 66–69.
- 201 McNaughton LR, Dalton B, Palmer G. Sodium bicarbonate can be used as an ergogenic aid in high intensity, competitive cycle ergometry of 1h duration. *Eur J Appl Physiol* 1999; 80: 64–69.
- 202 McNaughton LR, Lovell RJ, Siegler JC et al. The effects of caffeine ingestion on time trial cycling performance. *J Sports Med Phys Fitness* 2008; 48: 320–325.
- 203 McNaughton LR. Bicarbonate and citrate. In Maughan RJ. *Nutrition in Sports*. Oxford : Blackwell Science, 2000, p. 379–392.
- 204 Mero A, Nykänen T, Keinänen O. Protein metabolism and strength performance after bovine colostrum supplementation. *Amino Acids* 2005; 28: 327–335.
- 205 Michel BA, Stucki G, Frey D et al. Chondroitin s4 and s6 sulfate in osteoarthritis of the knee: a randomized, controlled trial. *Arthritis Rheum* 2005; 52: 779–786.
- 206 Miller-Ihli NJ. Graphite furnace atomic absorption spectrometry for the determination of the chromium content of selected U.S. foods. *J Food Comp Anal* 1996; 9: 290.
- 207 Mizuno K, Tanaka M, Nozaki S et al. Antifatigue effects of coenzyme Q10 during physical fatigue. *Nutrition* 2008; 24: 293–299.
- 208 Monfort J, Martel-Pelletier J, Pelletier J-P. Chondroitinsulphate for symptomatic osteoarthritis: critical appraisal of meta-analyses. *Curr Med Res Opin* 2008; 24: 1303–1308.
- 209 Monfort J, Pelletier J-P, Garcia-Giralt N, Martel-Pelletier J. Biochemical basis of the effect of chondroitin sulphate on osteoarthritis articular tissues. *Ann Rheum Dis* 2008; 67: 735–740.
- 210 Monteiro GA, Aoki MS, Santos CB et al. Effects of branched-chain amino acid (BCAA) supplementation on endurance exercise performance of pregnant rats. *Science & Sports* 2009; 24: 102–107.
- 211 Morens C, Bos C, Pueyo ME et al. Increasing habitual protein intake accentuates differences in postprandial dietary nitrogen utilization between protein sources in humans. *J Nutr* 2003; 133: 2733–2740.
- 212 Mougios V, Matsakas A, Petridou A et al. Effect of supplementation with conjugated linoleic acid on human serum lipids and body fat. *J Nutr Biochem* 2001; 12: 585–594.
- 213 Muazzezaneh A, Keshavarz SA, Yaraghi SA et al. Effect of L-Arginine supplementation on blood lactate level and VO₂max at anaerobic threshold performance. *KAUMS Journal (FEYZ)* 2010; 14: 200–208.
- 214 Mujika I, Padilla S, Ibanez J et al. Creatine supplementation and sprint performance in soccer players. *Med Sci Sports Exerc* 2000; 32: 518–525.
- 215 Müller DM, Seim H, Kiess W et al. Effects of oral L-carnitine supplementation on in vivo long-chain fatty acid oxidation in healthy adults. *Metabolism* 2002; 51: 1389.
- 216 Müller-Fassbender H. Double-blind clinical trial multicenter study comparing S-adenosyl-methionine, naproxen, and placebo in the treatment of osteoarthritis. *Am J Med* 1987; 83: 1–4

- 217 Muoio DH, Leddy JJ, Horvath PJ et al. Effects of dietary fat on metabolic adjustments to maximal VO_2 and endurance in runners. *Med Sci Sports Exerc* 1992; 26: 81–88.
- 218 Musil J, Nováková O, Kunz K. *Biochemie v obrazech a schématech*. Praha : Avicenum, 1976.
- 219 Nair KS, Schwartz RG, Welle S. Leucine as a regulator of whole body and skeletal muscle protein metabolism in humans. *Am J Physiol Endocrinol Metab* 1992; 263: E928–E934.
- 220 Newhouse IJ, Clement DB. The efficacy of iron supplementation in iron depleted women. In Kies CV, Driskell JA (ed.), *Sports nutrition: minerals and electrolytes*. Boca Raton : CRC Press, 1995.
- 221 Newsholme E et al. Physical and mental fatigue: Metabolic mechanisms and importance of plasma amino acids. *Brit Med Bull* 1992; 48: 477–495.
- 222 Nieman D. Exercise immunology: Nutrition countermeasures. *Can J Appl Physiol* 2001; 26: S45–S55.
- 223 Nicholas C. Legal nutritional supplements during a sporting event. *Essays Biochem* 2008; 44: 45–61.
- 224 Nissen S, Sharp R, Rathmacher JA et al. The effect of leucine metabolite beta-hydroxy beta-methylbutyrate (HMB) on muscle metabolism during resistance-exercise training. *J Appl Physiol* 1996; 81: 2095–2104.
- 225 Nissen SL, Abumrad NN. Nutritional role of the leucine metabolite beta-hydroxy-beta-methylbutyrate (HMB). *J Nutr Biochem* 1997; 8: 300–311.
- 226 Nissen SL, Sharp RL. Effect of dietary supplements on lean mass and strength gains with resistance exercise: a meta-analysis. *J Appl Physiol* 2003; 94: 651–659.
- 227 Norton LE, Layman DK. Leucine regulates translation initiation of protein synthesis in skeletal muscle after exercise. *J Nutr* 2006; 136: S533–S537.
- 228 Nosaka N, Suzuki Y, Nagatoshi A et al. Effect of Ingestion of Medium-Chain Triacylglycerols on Moderate- and High-Intensity Exercise in Recreational Athletes. *J Nutr Sci Vitaminol* 2009; 55, 120–125.
- 229 Novotný V. IIInd Anthropological Congress of Aleš Hrdlička: proceedings of the IIInd Anthropological Congress dedicated to Dr. Aleš Hrdlička (held in Prague and Humpolec, September 3–7, 1979). Praha : Univerzita Karlova, 1982. 504 s.
- 230 Paddon-Jones D, Sheffield-Moore M, Zhang X-J et al. Amino acid ingestion improves muscle protein synthesis in the young and elderly. *Am J Physiol Endocrinol Metab* 2004; 286: E321–E328.
- 231 Paffenbarger RS Jr, Hyde RT, Wing AL et al. The association of changes in physical activity level and other lifestyle characteristics with mortality among men. *New J Engl Med* 1993; 328: 538–545.
- 232 Parise G, Mihic S, MacLennan KE et al. Effects of acute creatine monohydrate supplementation on leucine kinetics and mixed-muscle protein synthesis. *J Appl Physiol* 2001; 91: 1041–1047.
- 233 Pariza MW, Park Y, Cook ME. The biologically active isomers of conjugated linoleic acid. *Prog Lipid Res* 2001; 40: 283–298.

- 234 Park Y, Storkson JM, Albright KJ et al. Evidence that the trans-10, cis-12 isomer of conjugated linoleic acid induces body composition changes in mice. *Lipids* 1999; 34: 235–241.
- 235 Parkhouse WS, McKenzie DC. Possible contribution of skeletal muscle buffers to enhanced anaerobic performance: a brief review. *Med Sci Sports Exerc* 1984; 16: 328–338.
- 236 Pařízková J. *Body Fat and Physical Fitness: Body Composition and Lipid Metabolism in Different Regimes of Physical Activity*. Hague : Nijhoff, 1977. 279 s.
- 237 Pate RR, Miller BJ, Davis JM et al. Iron status of female runners. *Int J Sports Nutr* 1993; 3: 222.
- 238 Patti ME, Brambilla E, Luzi L et al. Bidirectional modulation of insulin action by amino acids. *J Clin Invest* 1998;101: 1519–1529.
- 239 Pavelka K, Gatterová J, Olejarová M et al. Glucosamine sulfate use and delay of progression of knee osteoarthritis. *Arch Int Med* 2002; 162: 2113–2123.
- 240 Phillips SM. Protein requirements and supplementation in strength sports. *Nutrition* 2004; 20: 689–695.
- 241 Pilegaard H, Domino K, Noland T et al. Effect of high intensity training on lactate/H⁺ transport capacity in human skeletal muscle. *Am J Physiol* 1999; 276: 255–261.
- 242 Pitts GC, Consolazio FC, Robinson PF. Dietary protein and physical fitness in temperate and hot environments. *J Nutr* 1944; 27: 497–508.
- 243 Pollock ML, Miller HS Jr, Linnerud AC et al. Frequency of training as a determinant for improvement in cardiovascular function and body composition of middle-aged men. *Arch Phys Med Rehabil* 1975; 56: 141–145.
- 244 Pollock ML, Wilmore JH, Fox SM. *Exercise in Health and Disease*. Philadelphia : WB Saunders, 1984.
- 245 Pollock ML. The quantification of endurance training programs. In Wilmore JH (ed.), *Exercise and Sport Sciences Reviews*. New York : Academic Press, 1973, p. 155–188.
- 246 Portier H, Chatard JC, Filaire E et al. Effects of branched-chain amino acids supplementation on physiological and psychological performance during an offshore sailing race. *Eur J Appl Physiol* 2008; 104: 787–794.
- 247 Poortmans JR, Francaux M. Long-term oral creatine supplementation does not impair renal function in healthy athletes. *Med Sci Sports Exerc* 1999; 31:1108–1110.
- 248 Pozefsky T, Walser M. Effect of intraarterial infusion of the ketoanalogue of leucine on amino acid release by forearm muscle. *Metabolism* 1977; 26: 807–815.
- 249 Proud CG. mTOR-mediated regulation of translation factors by amino acids. *Biochem Biophys Res Commun* 2004; 313: 429–436.
- 250 Rasmussen BB, Tipton KD, Miller SL et al. An oral essential amino acid-carbohydrate supplement enhances muscle protein anabolism after resistance exercise. *J Appl Physiol* 2000; 88: 386–392.
- 251 Rawson ES, Volek JS. Effects of creatine supplementation and resistance training on muscle strength and weightlifting performance. *J Strength Cond Res* 2003; 17: 822–831.

- 252 Reed JD. They hunger for success. *Sports Illustrated* 1977; 46: 65–68, 71–72.
- 253 Reichenbach S, Sterchl R, Scherer M et al. Meta-analysis: chondroitin for osteoarthritis of the knee or hip. *Ann Intern Med* 2007; 146: 580–590.
- 254 Richy F, Bruyere O, Ethgen O et al. Structural and symptomatic efficacy of glucosamine and chondroitin in knee osteoarthritis: a comprehensive meta-analysis. *Arch Intern Med* 2003; 163: 1514–1522.
- 255 Rohde T et al. Effect of glutamine supplementation on changes in the immune system induced by repeated exercise. *Med Sci Sports Exerc* 1998; 30: 856–862.
- 256 Rowland TW, Deisroth MB, Green GM et al. The effect of iron therapy on the exercise capacity of nonanemic iron-deficient adolescent runners. *Am J Dis Child* 1988; 142: 165.
- 257 Santos RS, Pacheco MTT, Martins RABL. Study of the effect of oral administration of L-arginine on muscular performance in healthy volunteers: An isokinetic study. *Isokinetics and Exercise Science* 2002; 10: 153–158.
- 258 Sawitzke AD, Shi H, Finco MF et al. The effect of glucosamine and/or chondroitin sulfate on the progression of knee osteoarthritis: a report from the Glucosamine/Chondroitin Arthritis Intervention Trial. *Arthritis Rheum* 2008; 58: 3183–3191.
- 259 Sen C. Antioxidants in exercise nutrition. *Sports Med* 2001; 31: 891–908.
- 260 Shah OJ, Anthony JC, Kimball SR et al. 4E-BP1 and S6K1: translational integration sites for nutritional and hormonal information in muscle. *Am J Physiol* 2000; 279: E715–E729.
- 261 Sherman WM, Costill DL. The marathon. Dietary manipulation to optimize performance. *Am J Sports Med* 1984; 12: 44–51.
- 262 Schoenekess BO, Lopaschuk GD. The effect of carnitine on myocardial carbohydrate metabolism. In DeJong JW, Ferrari R (ed.), *The Carnitine System*. Amsterdam : Kluwer Academic, 1995, s. 39–52.
- 263 Siliprandi N, Vecchiet L, Di Lisa F. Carnitine administration in physical exercise. In Benzi G (ed.), *Advances in myochemistry*. Paris : J Libbey, 1989, s. 273–278.
- 264 Singh A et al. Chronic multivitamin-mineral supplementation does not enhance physical performance. *Med Sci Sports Exerc* 1992; 24: 726–732.
- 265 Smedman A, Vessby B. Conjugated linoleic acid supplementation in humans – metabolic effects. *Lipids* 2001; 36: 773–781.
- 266 Spriet LL, Howlett RA. Caffeine. In Maughan RJ. *Nutrition in Sports*. Oxford : Blackwell Science, 2000, s. 379–392.
- 267 Stear SJ, Burke LM, Castell LM. A–Z of nutritional supplements: dietary supplements, sports nutrition foods and ergogenic aids for health and performance. Part 3. *Br J Sports Med* 2009; 43: 890–892.
- 268 Steidl L, Zbránková B. Význam karnitinu a jeho použití v medicíně. Praha : Triton, 2000.
- 269 Stejskal P, Hejnová J. Praktické problémy preskripce intenzity v rámci programu tělesné aktivity. *Med Sport Bohemoslov* 1993; 2: 76–81.
- 270 Stejskal P, Hejnová J. Preskripce intenzity zatížení v programu kontinuální tělesné aktivity. *Med Sport Bohemoslov* 1992; 1: 11–16.

- 271 Stejskal P. Preskripce trvání tréninku, jeho energetického výdeje a týdenní frekvence v rámci aerobní části programu tělesné aktivity. *Med Sport Boh Slov* 1993; 2: 93–98.
- 272 Stellingwerff T. β -Alanine and Carnosine: The science and its application. Lausanne : Nestlé Research Center, 2011.
- 273 Stoppani J, Scheett T, Pena J et al. Consuming a supplement containing branched-chain amino acids during a resistance-training program increases lean mass, muscle strength and fat loss. *J Int Soc Sports Nutr* 2009; 6(Suppl. 1): P1.
- 274 Stout JR, Cramer JT, Zoeller RF et al. Effects of beta-alanine supplementation on the onset of neuromuscular fatigue and ventilatory threshold in women. *Amino Acids (Vienna)* 2007; 32: 381–386.
- 275 Suminski R et al. Acute effect of amino acid ingestion and resistance exercise on plasma growth hormone concentration in young men. *Int J Sport Nutr* 1997; 7: 48–60.
- 276 Takanami Y et al. Vitamin E supplementation and endurance exercise: Are there benefits? *Sports Med* 2000; 29: 73–83.
- 277 Telford R et al. The effect of 7 to 8 months of vitamin/mineral supplementation on athletic performance. *Int J Sport Nutr* 1992; 2: 135–153.
- 278 Terblanche S, Noakes TD, Dennis SC et al. Failure of magnesium supplementation to influence marathon running performance or recovery in magnesium-replete subjects. *Int J Sports Nutr* 1992; 2: 154.
- 279 Thom E, Wadstein J, Gudmundsen O. Conjugated linoleic acid reduces body fat in healthy exercising humans. *J Int Med Res* 2001; 29: 392–396.
- 280 Thomas C, Perrey S, Lambert K et al. Monocarboxylate transporters, blood lactate removal after supramaximal exercise, and fatigue indexes in humans. *J Appl Physiol* 2005; 98: 804–809.
- 281 Tipton K, Wolfe R. Exercise, protein metabolism, and muscle growth. *Int J Sport Nutr Exerc Metabol* 2001; 11: 109–132.
- 282 Tipton KD, Ferrando AA, Phillips SM et al. Postexercise net protein synthesis in human muscle from orally administered amino acids. *Am J Physiol Endocrin Metab* 1999; 276: E628–E634.
- 283 Tipton KD, Rasmussen BB, Miller SL et al. Timing of amino acid-carbohydrate ingestion alters anabolic response of muscle to resistance exercise. *Am J Physiol Endocrin Metab* 2001; 281: E197–E206.
- 284 Tomita M, Sendju Y. Über Die Oxyaminverbindungen welche die Biuretreaktion zeigen. *Z Physiol Chem* 1927; 196: 263–267. In Steidl L, Zbránková B. Význam karnitinu a jeho použití v medicíně. Praha : Triton, 2000.
- 285 Tong BC, Barbul A. Cellular and physiological effects of arginine. *Mini Rev Med Chem* 2004; 4: 823–832.
- 286 Trent LK, Thieding-Cancel D. Effects of chromium picolinate on body composition. *J Sports Med Phys Fitness* 1995; 35: 273.
- 287 Uebelhart D, Malaise M, Marcolongo R et al. Intermittent treatment of knee osteoarthritis with oral chondroitin sulfate: A one year, randomized, double-blind, multicenter study versus placebo. *Osteoarthritis Cartilage* 2004; 12: 269–276.

- 288 Uffelmann K, Vogler W, Fruth C. Der Einsatz hydrolytischer Enzyme beim extra-artikulären Rheumatismus. *Allgemeinmedizin* 1990; 19: 151–153.
- 289 Van der Beek E. Vitamin supplementation and physical exercise performance. *J Sport Sci* 1991; 92: 77–79.
- 290 van Someren KA, Edwards AJ, Howatson G. Supplementation with beta-hydroxy-beta-methylbutyrate (HMB) and alphaketoisocaproic acid (KIC) reduces signs and symptoms of exercise-induced muscle damage in man. *Int J Sport Nutr Exerc Metab* 2005; 15: 413–424.
- 291 Vandenberghe K, Goris M, van Hecke P et al. Long-term creatine intake is beneficial to muscle performance during resistance training. *J Appl Physiol* 1997; 83: 2055–2063.
- 292 Vangsness CT Jr, Spiker W, Erickson J. A Review of evidence-based medicine for glucosamine and chondroitin sulfate use in knee osteoarthritis. *Arthroscopy* 2009; 25: 86–94.
- 293 Vávra J, Máček M. *Fyziologie a patofyziologie tělesné zátěže*. Praha : Avicenum, 1982.
- 294 Vecchiet L et al. Influence of L-carnitine administration on maximal physical exercise. *Eur J Appl Physiol* 1990; 61: 486–490.
- 295 Verbitsky O, Mizrahi J, Levin M et al. Effect of ingested sodium bicarbonate on muscle force, fatigue, and recovery. *J Appl Physiol* 1997; 83: 333–337.
- 296 Vilikus Z, Brandejský P, Novotný V. *Tělovýchovné lékařství*. Praha : Karolinum, 2004. 257 s.
- 297 Vincent JB. The biochemistry of chromium. *J Nutr* 2000; 130: 715–721.
- 298 Volek JS, Duncan ND, Mazzetti SA et al. Performance and muscle fiber adaptations to creatine supplementation and heavy resistance training. *Med Sci Sports Exerc* 1999; 31: 1147–1156.
- 299 Volek JS, Kraemer WJ, Bush JA et al. Creatine supplementation enhances muscular performance during high-intensity resistance exercise. *J Am Diet Assoc* 1997; 97: 765–770.
- 300 Volpi E, Kobayashi H, Sheffield-Moore M et al. Essential amino acids are primarily responsible for the amino acid stimulation of muscle protein anabolism in healthy elderly adults. *Am J Clin Nutr* 2003; 78: 250–258.
- 301 Vráblík M. Omega-3 mastné kyseliny a kardiovaskulární onemocnění. *Medicína pro praxi* 2007; 4: 12, www.medicinapropraxi.cz.
- 302 Vukovich MD, Slater G, Macchi MB et al. Beta-hydroxy-beta-methylbutyrate (HMB) kinetics and the influence of glucose ingestion in humans. *J Nutr Biochem* 2001; 12: 631–639.
- 303 Wall BT, Stephens FB, Constantin-Teodosiu D et al. Increasing muscle carnitine content alters muscle fuel metabolism and improves exercise performance in humans. *Japan J Phys Fitness Sports Med* 2011; 60: 85.
- 304 Walzem RM, Dillard CJ, German JB. Whey components: millennia of evolution create functionalities for mammalian nutrition: what we know and what we may be overlooking. *Crit Rev Food Sci Nutr* 2002; 42: 353–375.

- 305 Warren GL, Park ND, Maresca RD et al. Effect of caffeine ingestion on muscular strength and endurance: A meta-analysis. *Med Sci Sports Exerc* 2010; 42: 1375–1387.
- 306 Watras AC, Buchholz AC, Close RN et al. The role of conjugated linoleic acid in reducing body fat and preventing holiday weight gain. *Int J Obes* 2007; 31: 481.
- 307 Watson P et al. The effect of acute branched-chain amino acid supplementation on prolonged exercise capacity in a warm environment. *Eur J Appl Physiol* 2004; 93: 306–314.
- 308 Weight ML, Noakes TD, Labadarios D et al. Vitamin and mineral status of trained athletes including the effects of supplementation. *Am J Clin Nutr* 1988; 47: 186–191.
- 309 West DB, Delany JP, Camet PM et al. Effects of conjugated linoleic acid on body fat and energy metabolism in the mouse. *Am J Physiol* 1998; 275: 667–672.
- 310 Whigham LD, Watras AC, Schoeller DA. Efficacy of conjugated linoleic acid for reducing fat mass: a meta-analysis in humans. *Am J Clin Nutr* 2007; 85: 1203–1211.
- 311 Wildi LM, Raynauld J-P, Martel-Pelletier J et al. Chondroitin sulphate reduces both cartilage volume loss and bone marrow lesions in knee osteoarthritis patients starting as early as 6 months after initiation of therapy: a randomised, double-blind, placebo-controlled pilot study using MRI. *Ann Rheum Dis* 2011; 70: 982–989.
- 312 Wildman R, Miller B. *Sports and Fitness Nutrition*. USA : Thomson Wadsworth, 2004.
- 313 Williams MH, Leutholtz BC. Nutritional ergogenic aids. In Maughan RJ. *Nutrition in Sports*. Oxford : Blackwell Science, 2000, s. 356–366.
- 314 Williams MH. Dietary supplements and sports performance: Introduction and vitamins. *J Int Soc Sports Nutr* 2004; 1: 1–6.
- 315 Williams MH. Facts and fallacies of purported ergogenic amino acid supplements. *Clin Sports Med* 1999; 18: 633–649.
- 316 Williams MH. *Nutrition for Health, Fitness & Sports*. Boston : McGraw-Hill. 2005.
- 317 Williams MH. Nutritional supplements for strength trained athletes. *Sports Sci Exchange* 1993; 6: 1–7.
- 318 Williams MH. The use of nutritional ergogenic aids in sports: is it an ethical issue? *Int J Sport Nutr* 1994; 4: 120–131.
- 319 Williamson JR, Walajtys-Rode E, Coll KE. Effects of branched chain alpha-ketoacids on the metabolism of isolated rat liver cells. Regulation of branched chain alpha-ketoacid metabolism. *J Biol Chem* 1979; 254: 11511–11520.
- 320 Willoughby DS, Rosene J. Effects of oral creatine and resistance training on myosin heavy chain expression. *Med Sci Sports Exerc* 2001; 33: 1674–1681.
- 321 Wittenborg PR, Hanisch J, Sallet R et al. Comparative epidemiological study in patients with rheumatic diseases illustrated in an example of a treatment with non-steroidal anti-inflammatory drugs versus an oral enzyme combination. *Drug Res* 2000; 50: 728–738.
- 322 Wolfe RR. Regulation of muscle protein by amino acids. *J Nutr* 2002; 132: 3219S–3224S.

