

Bibliografie

Benatti FB, Ried-Larsen M. (2015) „The effects of breaking up prolonged sitting time: a review of experimental studies.“ *Med Sci Sports Exerc* 47(10): 2053–2061.

Buckley JP, Hedge A, Yates T, Copeland RJ, Loosemore M, Hamer M, Bradley G, Dunstan DW. (2015) „The sedentary office: an expert statement on the growing case for change toward better health and productivity.“ *Br J Sports Med* 49(21): 1357–62. doi: 10.1136/bjsports-2015-094618.

Casperan CJ, Powell KE, Christenson, GM. (1985) „Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research.“ *Public Health Rep* 100(2): 126–131. www.ncbi.nlm.nih.gov/pmc/articles/PMC1424733/

Chatterjee, R. (12 February 2020) Podcast: „How Exercise Changes Your Brain and Reduces Your Risk of Depression with Brendon Stubbs.“ drchatterjee.com/how-exercise-changes-your-brain-and-reduces-your-risk-of-depression/

Dempsey, PC, Larsen, RN, Winkler, EAH, Owen, N, Kingwell, BA, Dunstan, DW. (2018) „Prolonged uninterrupted sitting elevates postprandial hyperglycaemia proportional to degree of insulin resistance.“ *Diabetes, Obesity and Metabolism* 20(6): 1526–30.

doi.org/10.1111/dom.13254

Dempsey, PC, Sacre, JW, Larsen, RN, Straznicky, NE, Sethi, P, Cohen, ND, Cerin, E, Lambert, GW, Owen, N, Kingwell, BA, Dunstan, DW. (2016) „Interrupting prolonged sitting with brief bouts of light walking or simple resistance activities reduces resting blood pressure and plasma noradrenaline in type 2 diabetes.“ *Journal of Hypertension* 34(12): 2376–82. doi.org/10.1097/HJH.0000000000001101

Fukushima, N, Kitabayashi, M, Kikuchi, H, Sasai, H, Oka, K, Nakata, Y, Tanaka, S, Inoue, S. (2018) „Comparison of accelerometer-measured sedentary behavior, and light- and moderate-to-vigorous-intensity physical activity in white- and blue-collar workers in a Japanese manufacturing plant.“ *Journal of Occupational Health* 60(3): 246–253. doi: 10.1539/joh.2017-0276-OA.

Helgadóttir, B, Forsell, Y, Hallgren, M, Möller, J, Ekblom, Ö. (2017) „Long-term effects of exercise at different intensity levels on depression: A randomized controlled trial.“ *Prev Med* 105: 37–46. doi: 10.1016/j.ypmed.2017.08.008.

Jackson, B, O'Connell, SE, Waheed, G, Munir, F. (2018) „Effectiveness of the stand more at (SMaRT) work intervention: Cluster randomised controlled trial.“ *BMJ* 363: 3870. doi.org/10.1136/bmj.k3870

Kandola, A, Rees, J, Stubbs, B, Dunstan, DW, Genevieve GN, Hayes JF. (2020) „Breaking up excessive sitting with light activity.“ cms.wellcome.org/sites/default/files/2021-05/breaking-up-excessive-sitting-light-activity-wellcome-workplace-mental-health.pdf

Loh, R, Stamatakis, E, Folkerts, D, Allgrove, JE, Moir, HJ. (2020) „Effects of Interrupting Prolonged Sitting with Physical Activity Breaks on Blood Glucose, Insulin and Triacylglycerol Measures: A Systematic Review and Meta-analysis.“ *Sports Medicine* 50(20): 295–330. doi.org/10.1007/s40279-019-01183-w

Mackie, P, Weerasekara, I, Crowfoot, G, Janssen, H, Holliday, E, Dunstan, D, English, C. (2019) „What is the effect of interrupting prolonged sitting with frequent bouts of physical activity or standing on first or recurrent stroke risk factors? A scoping review.“ *PLOS ONE* 14(6): e0217981. doi.org/10.1371/journal.pone.0217981

Munir, F, Biddle, SJH, Davies, MJ, Dunstan, D, Esliger, D, Gray, LJ, Jackson, BR, O'Connell, SE, Yates, T, Edwardson, CL. (2018) „Stand More AT Work (SMARt Work): Using the behaviour change wheel to develop an intervention to reduce sitting time in the workplace.“ *BMC Public Health* 18, 319. doi.org/10.1186/s12889-018-5187-1

Prince, SA, Cardilli, L, Reed, JL, Saunders, TJ, Kite, C, Douillette, K, Fournier, K, Buckley, JPage (2020) „A comparison of self-reported and device measured sedentary behaviour in adults: A systematic review and meta-analysis.“ *International Journal of Behavioral Nutrition and Physical Activity* 17(1): 1–17. BioMed Central Ltd. doi.org/10.1186/s12966-020-00938-3

Straker, L, Abbott, RA, Heiden, M, Mathiassen, SEtoomingas, A. (2013) „Sit-stand desks in call centres: Associations of use and ergonomics awareness with sedentary behavior.“ *Appl Ergon* 44(4): 517–22. doi.org/10.1016/j.apergo.2012.11.001