

20. Literatura

- ABADAL-CENTELLAS, JM., LLOMPART-POU, JA., HOMAR-RAMIREZ, J., et al. *Neurologic outcome of posttraumatic refractory intracranial hypertension treated with external lumbar drainage*. J Trauma, 2007, 62, p. 282–286.
- ABBRUSCATO, TJ., DAVIS, TP. *Combination of hypoxialaglycemia compromises in vitro blood-brain barrier integrity*. JPET, 1999, 289 (2), p. 668–675.
- ABE, KK., YAMAMOTO, LG., ITOMAN, EM., et al. *Lumbar puncture needle length determination*. Am J Emerg Med, 2005, 23, p. 742–746.
- ADAM, P. *Cytologie likvoru*. Pardubice : Stapro, 1995.
- ADAM, P., KRATOCHVÍLA, J., SOBEK, O., et al. *Cytologie mozkomíšního moku*. Praha : SEKK, 2000.
- ADAM, P., ANDRÝS, C., FRÝDECKÝ, B., et al. *Doporučení k vyšetřování mozkomíšního moku*. Klin Biochem Metab, 2006, 14 (35), s. 125–127.
- ALARCON, A., GARCIA ALIX, A., CABAÑAS, F., et al. *Beta₂-microglobulin concentrations in cerebrospinal fluid correlate with neuroimaging findings in newborns with symptomatic congenital cytomegalovirus infection*. Eur J Pediatr, 2006, 165, p. 636–645.
- ALBECK, MJ., BØRGESEN, SE., GJERRIS, F., et al. *Intracranial pressure and cerebrospinal fluid outflow conductance in healthy subjects*. J Neurosurg, 1991, 74, p. 597–600.
- ALBECK, MJ., SKAK, C., NIELSEN, PR., et al. *Age-dependency of resistance to cerebrospinal fluid outflow*. J Neurosurg, 1998, 89, p. 275–278.
- ALLEN, SH. *How to perform a lumbar puncture with the patient in the seated position*. Br J Hosp Med, 2006, 67, p. M46–M47.
- AMBLER, Z., BEDNAŘÍK, J., RŮŽIČKA, E., et al. *Klinická neurologie*. Praha : Triton, 2004.
- ANDERSSON, C., BLENNOW, K., JOHANSSON, SE., et al. *Differential CSF biomarker levels in APOE-epsilon4-positive and -negative patients with memory impairment*. Dement Geriatr Cogn Disord, 2007, 23 (2), p. 87–95.
- ANDERSON, L., MARSHALL, S., BRYDON, C., SERPELL, MG. *Comparison of dural indenting between Whitacre and Quincke tipped spinal needles*. Br J Anaesth, 1997, 78, p. 468–469P.
- ANDERSSON, M., ALVAREZ-CERMEÑO, J., BERNARDI, G., et al. *Cerebrospinal fluid in the diagnosis of multiple sclerosis: a consensus report*. J Neurol Neurosurg Psychiatry, 1994, 57, p. 897–902.
- AOKI, N. *Lumboperitoneal shunt: Clinical applications, complications, and comparison with ventriculoperitoneal shunt*. Neurosurgery, 1990, 26, p. 998–1004.
- AZZONI, R., GEREVINI, S., CABITZA, P. *Spinal cord sonography in newborns: anatomy and diseases*. J Pediatr Orthop B, 2005, 14, p. 185–188.

- BACHMANN, G., PETEREIT, H. *β -trace protein as sensitive marker for liquorrhea.* Acta Neurol Scand, 2004, 110, p. 337–338.
- BALAŠ, V., VLČEK, J. *Akademik Arnold Jirásek – český chirurg.* Praha : Avicenum, 1988.
- BALDWIN, HZ., REKATE, HL. *Preliminary experience with controlled external lumbar drainage in diffuse pediatric head injury.* Pediatr Neurosurg, 1991–1992, 17, p. 115–120.
- BANIK, R., LIN, D., MILLER, NR. *Prevalence of Chiari I malformation and cerebellar ectopia in patients with pseudotumor cerebri.* J Neurol Sci, 2006, 247, p. 71–75.
- BANNISTER, R., et al. *Isotope encephalography in the diagnosis of dementia due to communicating hydrocephalus.* Lancet, 1967, 2, p. 1014–1101.
- BARKOVICH, AJ., WIPPOLD, FJ., SHERMAN, JL. *Significance of cerebellar tonsillar position on MR.* Am J Neuroradiol, 1986, 7, p. 795–799.
- BARRY, A., SINGER, MD. *ECTRIMS 2006. Diagnosis and Monitoring of Multiple Sclerosis: Focus on Cerebrospinal Fluid Analysis and Brain Imaging.* [online] Dostupné z URL: www.medscape.com/viewarticle/548062.
- BARSON, AJ. *The vertebral level of termination of the spinal cord during normal and abnormal development.* J Anat, 1970, 106, p. 489–497.
- BARTOŠ, A., PIŤHA, J. *Axonální patologie v časných stádiích roztroušené sklerózy.* Cesk Slov Neurol N, 2004, 67 (5), s. 381.
- BARTOŠ, A., FIALOVÁ, L., SOUKUPOVÁ, J., et al. *Elevated intrathecal antibodies against the medium neurofilament subunit in multiple sclerosis.* J Neurol, 2007, 254, p. 20–25.
- BAXTER, AL., WELCH, JC., BURKE, BL., ISAACMAN, DJ. *Pain, position, and stylet styles. Infant lumbar puncture practices of pediatric emergency attending physicians.* Pediatr Emerg Care, 2004, 20, p. 816–820.
- BAXTER, AL., FISHER, RG., BURKE, BL., et al. *Local anesthetic and stylet styles: Factors associated with resident lumbar puncture success.* Pediatrics, 2006, 117, p. 876–881.
- BEDNÁŘOVÁ, J., ADAM, P. *Likvorový profil u roztroušené sklerózy mozkomíšni.* Neurologie pro praxi, 2002, 5, s. 249–251.
- BEDNÁŘOVÁ, J., ŠTOURAC, P., KRBKOVÁ, L. *The diagnostic significance of intrathecal occurrence of selected antiviral antibodies in multiple sclerosis.* Scripta Medica, 2005, 78 (4), s. 249–254.
- BEJJANI, GK. *Association of the adult Chiari malformation and idiopathic intracranial hypertension: more than a coincidence.* Med Hypotheses, 2003, 60, p. 859–863.
- BERGAMINI, L., DURELLI, L. *Multiple sclerosis. I. The immune pathogenetic hypothesis.* Riv Neurol, 1989, 59 (5), p. 176–190.
- BERGER, T., RUBNER, P., SCHAUTZER, E., et al. *Antimyelin antibodies as a predictor of clinically definite multiple sclerosis after a first demyelinating event.* N Engl J Med, 2003, 349, p. 139–145.
- BERGSTRAND, G., OXENSTIERNA, G., FLYCKT, L., et al. *Radionuclide cisternography and computed tomography in 30 healthy volunteers.* Neuroradiology, 1986, 28, p. 154–160.
- BILIC, E., BILIC, E., DADIC, M., BOBAN, M. *Calculating lumbar puncture depth in children.* Coll Antropol, 2003, 27, p. 623–626.
- BINDER, DK., DILLON, WP., FISHMAN, RA., SCHMIDT, MH. *Intrathecal saline infusion in the treatment of obtundation associated with spontaneous intracranial hypotension: Technical case report.* Neurosurgery, 2002, 51, p. 830–837.

- BLENNOW, K. *Cerebrospinal fluid protein biomarkers for Alzheimer's disease*. NeuroRx, 2004, 1 (2), p. 213–225.
- BOESIGER, BM., SHIBER, JR. *Subarachnoid hemorrhage diagnosis by computed tomography and lumbar puncture: are fifth generation CT scanners better at identifying subarachnoid hemorrhage?* J Emerg Med, 2005, 29, p. 23–27.
- BOJAR, M. *Neuroborrelióza – současné názory na patogenezi, diagnostiku a léčbu*. Interní medicína pro praxi, 2002, 4, s. 154–159.
- BONADIO, WA. *Interpreting the traumatic lumbar puncture*. Contemp Pediatr Res Q, 1992, 1, p. 23–32.
- BOON, AJW., TANS, JTJ., DELWEL, EJ., et al. *Dutch normal-pressure hydrocephalus study: prediction of outcome after shunting by resistance to outflow of cerebrospinal fluid*. J Neurosurg, 1997, 87, p. 687–693.
- BOON, AJW., TANS, JTJ., DELWEL, EJ., et al. *The Dutch normal – pressure hydrocephalus study. How to select patients for shunting? An analysis of four diagnostic criteria*. Surg Neurol, 2000, 53, p. 201–207.
- BOON, JM., ABRAHAMS, PH., MEIRING, JH., WELCH, T. *Lumbar puncture: Anatomical review of a clinical skill*. Clin Anat, 2004, 17, p. 544–553.
- BØRGESEN, SE., GJERRIS, F., SORENSEN, SC. *The resistance to cerebrospinal fluid absorption in humans. A method of evaluation by lumbo-ventricular perfusion, with particular reference to normal pressure hydrocephalus*. Acta Neurol Scand, 1978, 57, p. 88–96.
- BØRGESEN, SE., GJERRIS, F. *The predictive value of conductance to outflow of CSF in normal pressure hydrocephalus*. Brain, 1982, 105, p. 65–86.
- BREUER, AC., TYLER, HR., MARZEWSKI, DJ., ROSENTHAL, DS. *Radicular vessels are the most probable source of needle-induced blood in lumbar puncture: significance for the thrombocytopenic cancer patient*. Cancer, 1982, 49, p. 2168–2172.
- BROADBENT, CR., MAXWELL, WB., FERRIE, R., et al. *Ability of anaesthetists to identify a marked lumbar interspace*. Anaesthesia, 2000, 55, p. 1122–1126.
- BRUN-BUISSON, CJ., DE GIALULY, E. *Fatal nonmeningitic listeria rhombencephalitis. Report of two cases*. Arch Intern Med, 1985, 145, p. 1982–1985.
- CABANA, MD. *Pseudotumor cerebri u 16letého mladíka*. Medicina, 2002, 9, s. 11.
- CARSON, D., SERPELL, M. *Choosing the best needle for diagnostic lumbar puncture*. Neurology, 1996, 47, p. 33–37.
- CAUDIE, C., ALLAUZEN, O., BANCEL, J., LATER, R. *Diagnostic usefulness of oligoclonal immunoglobulin G bands in cerebro-spinal fluid using isoelectric focusing in early diagnosis of multiple sclerosis*. Ann Biol Clin, 2000, 2 (58), p. 187–193.
- COGNARD, C., CASASCO, A., TOEVI, M., et al. *Dural arteriovenous fistulas as a cause of intracranial hypertension due to impairment of cranial venous outflow*. J Neurol Neurosurg Psychiatry, 1998, 65, p. 308–316.
- COLD, GE., MØLLER, K. *Treatment of intracranial hypertension and aspects on lumbar dural puncture in severe bacterial meningitis. Letter*. Acta Anaesthesiol Scand, 2002, 46, p. 1283–1284.
- CONE, LA., LEUNG, MM., BYRD, RG., et al. *Multiple cerebral abscesses because of Listeria monocytogenes: Three case reports and a literature review of supratentorial listerial brain abscess(es)*. Surg Neurol, 2003, 59, p. 320–328.

- CORBETT, JJ., MEHTA, MP. *Cerebrospinal fluid pressure in normal obese subjects and patients with pseudotumour cerebri*. Neurology, 1983, 33, p. 1386–1388.
- CORREALE, J., DE LOS MILAGROS BASSANI MOLINAS, M. *Oligoclonal bands and antibody responses in multiple sclerosis*. J Neurol, 2002, 249, p. 375–389.
- CROLS, R., SAERENS, J., NOPPE, M., LOWENTHAL, A. *Increased GFAP levels in CSF as a marker of organicity in patients with Alzheimer's disease and other types of irreversible chronic organic brain syndrome*. J Neurol, 1986, 233, p. 1432–1459.
- CRUICKSHANK, RH., HOPKINSON, JM. *Fluid flow through dural puncture sites. An in vitro comparison of needle point types*. Anaesthesia, 1989, 44, p. 415–418.
- CZOSNYKA, ZH., CZOSNYKA, M., WHITFIELD, PC., et al. *Cerebral autoregulation among patients with symptoms of hydrocephalus*. Neurosurgery, 2002, 50, p. 526–533.
- DAGNEW, E., VAN LOVEREN, HR., TEW jr., JM. *Acute foramen magnum syndrome caused by an acquired Chiari malformation after lumbar drainage of cerebrospinal fluid: Report of three cases*. Neurosurgery, 2002, 51, p. 823–829.
- DALE, RC., DE SOUSA, C., CHONG, WK., et al. *Acute disseminated encephalomyelitis, multiphasic disseminated encephalomyelitis and multiple sclerosis in children*. Brain, 2000, 123, p. 2407–2422.
- DAVSON, H., HOLLINGSWORTH, M., SEGAL, MB. *The mechanism and drainage of the cerebrospinal fluid*. Brain, 1970, 93, p. 665–678.
- DERFUSS, T., HOHLFELD, R., MEINL, E. *Intrathecal antibody (IgG) production against human herpesvirus type 6 occurs in about 20% of multiple sclerosis patients and might be linked to a polyspecific B-cell response*. J Neurol, 2005, 252, p. 968–971.
- DI CHIRO, G. *New radiographic and isotopic procedures in neurosurgical diagnosis*. JAMA, 1964, 188, p. 524–529.
- DI CHIRO, G. *Observations on the circulation of the cerebrospinal fluid*. Acta Radiol Scand, 1966, 5, p. 988–1002.
- DIGIOVANNI, AJ., GALBERT, MW., WAHLE, WM. *Epidural injection of autologous blood for postlumbar puncture headache. II. Additional clinical experiences and laboratory investigation*. Anesth Analg, 1972, 51, p. 226–232.
- DOMENICUCCI, M., RAMIERI, A., PAOLINI, S., et al. *Spinal subarachnoid hematomas: our experience and literature review*. Acta Neurochir, 2005, 147, p. 741–750.
- DRAYER, BP., ROSENBAUM, AE. *Studies of the third circulation. Amipaque CT cisternography and ventriculography*. J Neurosurg, 1978, 48, p. 946–956.
- DYER, EJ., FARAGHER, IG. *Computed tomography finding of portal venous gas in a patient who survived mesenteric infarction*. ANZ J Surg, 2005, 75, p. 363–364.
- EBINGER, F., KOSEL, C., PIETZ, J., et al. *Strict bed rest following lumbar puncture in children and adolescents is of no benefit*. Neurology, 2004, 62, p. 1003–1005.
- ECK, H. *Encephalomyelitis listeriaca apostematosa*. Schweiz Med Wochenschr, 1957, 87, S. 210–214.
- EHLING, R., LUTTEROTTI, A., WANSCHITZ, J., et al. *Increased frequencies of serum antibodies to neurofilament light in patients with primary chronic progressive multiple sclerosis*. Mult Scler, 2004, 10, p. 601–606.

- EIDE, PK., BREAN, A. *Intracranial pulse pressure amplitude levels determined during preoperative assessment of subjects with possible idiopathic normal pressure hydrocephalus*. Acta Neurochir, 2006, 148, p. 1151–1156.
- EIKELENBOOM, MJ., PETZOLD, A., LAZERON, RH., et al. *Multiple sclerosis: Neurofilament light chain antibodies are correlated to cerebral atrophy*. Neurology, 2003, 60, p. 219–223.
- EKSTEDT, J. *CSF hydrodynamic studies in man. Normal hydrodynamic variables related to CSF pressure and flow*. J Neurol Neurosurg Psychiatry, 1978, 41, p. 345–353.
- ELLIOTT, KAC., JASPER, HH. *Physiological salt solutions for brain surgery*. J Neurosurg, 1949, 6, p. 140–152.
- ERBAY, SH., O'CALLAGHAN, MG., BHADDELIA, R. *Is lumbar puncture contraindicated in patients with Chiari I malformation? Letter*. AJNR Am J Neuroradiol, 2005, 26, p. 985.
- ERDOGAN, C., HAKYEMEZ, B., YILDIRIM, N., PARLAK, M. *Brain abscess and cystic brain tumor. Discrimination with dynamic susceptibility contrast perfusion-weighted MRI*. J Comput Assist Tomogr, 2005, 29, p. 663–667.
- ETCHEPARE, F., ROCHE, B., ROZENBERG, S., et al. *Post-lumbar puncture arachnoiditis. The need for directed questioning*. Joint Bone Spine, 2005, 72, p. 180–182.
- FELGENHAUER, K., REIBER, H. *The diagnostic significance of antibody specificity indices in multiple sclerosis and herpes virus induced diseases of the nervous system*. Clin Investig, 1992, 70, p. 28–37.
- FELGENHAUER, K. *Laboratory Diagnosis of Neurological Diseases*. In THOMAS, L. (Ed.), *Clinical Laboratory Diagnostics. Use and Assessment of Clinical Laboratory Results*. Frankfurt am Main : TH-Books, 1998, p. 1308–1326.
- FERGUSON, AH. *Intraperitoneal diversion of the cerebrospinal fluid in cases of hydrocephalus*. NY Med J, 1898, 67, p. 902.
- FERGUSON, J., BRIN, M., GOLDMAN, R., et al. *Review of the literature on spinal ultrasound for the evaluation of back pain and radicular disorders: Report of the Therapeutics and Technology Assessment subcommittee of the American Academy of Neurology*. Neurology, 1998, 51, p. 343–344.
- FISHER, A., LUPU, L., GUREVITZ, B., et al. *Hip flexion and lumbar puncture: A radiological study*. Anaesthesia, 2001, 56, p. 262–266.
- FISHMAN, RA., DILLON, WP. *Dural enhancement and cerebral displacement secondary to intracranial hypotension*. Neurology, 1993, 43, p. 609–611.
- FLÍDROVÁ, E. *Patogeneze roztroušené sklerózy*. Bakalářská práce. Praha : Univerzita Karlova, Přírodovědecká fakulta, 2007, 48 s. Vedoucí bakalářské práce: doc. MUDr. Eva Havrdová, CSc.
- FLOETH, FW., PAULEIT, D., SABEL, M., et al. *¹⁸F-FET PET differentiation of ring-enhancing brain lesions*. J Nucl Med, 2006, 47, p. 776–782.
- FOLDES, EF., ARROWOOD, JG. *Changes in cerebrospinal fluid pressure under the influence of continuous subarachnoid infusion of normal saline*. J Clin Invest, 1948, 27, p. 346–351.
- FRANK, LR., PAXSON, A., BRAKE, J. *Spontaneous intracranial hypotension – a case report*. J Emerg Med, 2005, 28, p. 427–430.
- FREDERIKS, JAM., KOEHLER, PJ. *The first lumbar puncture*. J Hist Neurosci, 1997, 6, p. 147–153.
- GALIMBERTI, D., SCHOONENBOOM, N., SCHELTENS, P., et al. *Intrathecal chemokine synthesis in mild cognitive impairment and Alzheimer disease*. Arch Neurol, 2006, 63 (4), p. 538–543.

- GAUCHER jr., DJ., PEREZ jr., JA. *Subdural hematoma following lumbar puncture*. Arch Intern Med, 2002, 162, p. 1904–1905.
- GELWAN, MJ., CHOI, IS., BERENSTEIN, A., et al. *Dural arteriovenous malformations and papilledema*. Neurosurgery, 1988, 22, p. 1079–1084.
- GIBSON, T., NORRIS, W. *Skin fragments removed by injection needles*. Lancet, 1958, 2, p. 983–985.
- GJERRIS, F., SNORRASON, E. *The history of hydrocephalus*. J Hist Neurosci, 1992, 1, p. 285–312.
- GJERRIS, F., BECH-AZEDDINE, R. *Comment*. Neurosurgery, 2001, 49, p. 1185–1186.
- GLEASON, CA., MARTIN, RJ., ANDERSON, JV., et al. *Optimal position for a spinal tap in preterm infants*. Paediatrics, 1983, 71, p. 31–35.
- GOODMAN, JM. Spontaneous intracranial hypotension: A potential complication of meningeal diverticula or cysts. In KAUFMAN, HH. (Ed.), *Cerebrospinal fluid collections*. New York : Thieme, 1998, p. 183–186.
- GORELICK, PB., BILLER, J. *Lumbar puncture. Technique, indications, and complications*. Postgrad Med, 1986, 79, p. 257–268.
- GOSCH, UW., HUEPPE, M., HALLSCHMID, M., et al. *Post-dural puncture headache in young adults: comparison of two small-gauge spinal catheters with different needle design*. Br J Anaesth, 2005, 94, p. 657–661.
- GRAEF, IT., HENZE, I., REIBER, H. *Polyspezifische Immunreaktion im ZNS bei Autoimmunerkrankungen mit ZNS-Beteiligung*. Z Arztl Fortbild, 1994, 88, S. 587–591.
- GRAFF-RADFORD, NR., GODERSKY, JC., JONES, MP. *Variables predicting surgical outcome in symptomatic hydrocephalus in the elderly*. Neurology, 1989, 39, p. 1601–1604.
- GRÄNDE, P.-O., MYHRE, EB., NORDSTRÖM, CH., SCHLIAMSER, S. *Treatment of intracranial hypertension and aspects on lumbar dural puncture in severe bacterial meningitis*. Acta Anaesthesiol Scand, 2002, 46, p. 264–270.
- GRÄNDE, P.-O. *Mechanisms behind postspinal headache and brain stem compression following lumbar dural puncture – a physiological approach*. Acta Anaesthesiol Scand, 2005, 49, p. 619–626.
- GREENE, HM. *Lumbar puncture and the prevention of postpuncture headache*. JAMA, 1926, 86, p. 391–392.
- GRIEBEL, RW., BLACK, PM., PILE-PELLMAN, J., et al. *The importance of »accessory« outflow pathways in hydrocephalus after experimental subarachnoid hemorrhage*. Neurosurgery, 1989, 24, p. 187–192.
- HAKIM, R., BLACK, P. *Correlation between lumboventricular perfusion and MRI-CSF flow studies in idiopathic normal pressure hydrocephalus*. Surg Neurol, 1998, 49, p. 14–20.
- HANSEN, PB., JENSEN, TH., LYKKEGAARD, S., KRISTENSEN, HS. *Listeria monocytogenes meningitis in adults. Sixteen consecutive cases 1973–1982*. Scand J Infect Dis, 1987, 19, p. 55–60.
- HANSSON, O., ZETTERBERG, H., BUCHHAVE, P., et al. *Association between CSF biomarkers and incipient Alzheimer's disease in patients with mild cognitive impairment: a follow-up study*. Lancet Neurol, 2006, 5 (3), p. 228–234.
- HANSSON, O., ZETTERBERG, H., BUCHHAVE, P., et al. *Prediction of Alzheimer's disease using the CSF Abeta42/Abeta40 ratio in patients with mild cognitive impairment*. Dement Geriatr Cogn Disord, 2007, 23 (5), p. 316–320.

- HASEGAWA, K., YAMAMOTO, N. *Nerve root herniation secondary to lumbar puncture in the patient with lumbar canal stenosis. A case report.* Spine, 1999, 24, p. 915–917.
- HAVRDOVÁ, E., et al. *Neuroimunologie.* Praha : Maxdorf, 2001.
- HE, M., OHRUI, T., MARUYAMA, M., et al. *ACE activity in CSF of patients with mild cognitive impairment and Alzheimer disease.* Neurology, 2006, 67 (7), p. 1309–1310.
- HEARMON, C.J., GHOSH, S.K. *Listeria monocytogenes meningitis in previously healthy adults.* Postgrad Med J, 1989, 65, p. 74–78.
- HEBB, A.O., CUSIMANO, M.D. *Idiopathic normal pressure hydrocephalus: a systematic review of diagnosis and outcome.* Neurosurgery, 2001, 49, p. 1166–1186.
- HERUKKA, S.K., HELISALMI, S., HALLIKAINEN, M., et al. *CSF Abeta42, Tau and phosphorylated Tau, APOE epsilon4 allele and MCI type in progressive MCI.* Neurobiol Aging, 2007, 28 (4), p. 507–514.
- HERZIG, R., SCHNEIDERKA, P., HORAK, D., et al. *S-100β protein as a predictor of the size and clinical outcome in ischemic stroke. Review [abstract].* Cerebrovasc Dis, 2005, 19, Suppl. 2, p. 128.
- HOLST, D., MOLLMANN, M., EBEL, C., et al. *In vitro investigation of cerebrospinal fluid leakage after dural puncture with various spinal needles.* Anesth Analg, 1998, 87, p. 1331–1335.
- HOUDEK, M., KALA, M., PAVLÍČEK, V. *Mišní spasticita – nové léčebné možnosti.* Prakt Lék, 1996, 76, s. 110–112.
- HUGHES, L.E., BONELL, S., NATT, R.S., et al. *Antibody Responses to Acinetobacter spp. and Pseudomonas aeruginosa in Multiple Sclerosis: Prospects for Diagnosis Using the Myelin-Acinetobacter-Neurofilament Antibody Index.* Clin and Diagn Lab Immunol, 2001, 8 (6), p. 1181–1188.
- HUMPHREYS, R.P. *Computed tomography and the early diagnostic lumbar puncture.* Can Med Assoc J, 1979, 121, p. 150–151.
- HUMPHREYS, R.P. *Is lumbar puncture alive and well?* Pediatr Neurosurg, 1991–1992, 17, p. 114.
- CHAN, Y.C., HO, K.H., TAMBYAH, P.A., et al. *Listeria meningoenzephalitis: two cases and review of the literature.* Ann Acad Med Singapore, 2001, 30, p. 659–663.
- CHUDÁČKOVÁ, J., BOHUNĚK, V. *Vyšetření mozkomíšního moku.* In HARTL, J., et al., *Základní vyšetřovací metody v neurologii.* 2., přeprac. vyd. Olomouc : Lékařská fakulta Univerzity Palackého, 1991.
- CHUMAS, P.D., ARMSTRONG, D.C., DRAKE, J.M., et al. *Tonsillar herniation: the rule rather than the exception after lumboperitoneal shunting in the pediatric population.* J Neurosurg, 1993, 78, p. 568–573.
- CHUMAS, P.D., KULKARNI, A.V., DRAKE, J.M., et al. *Lumboperitoneal shunting: a retrospective study in the pediatric population.* Neurosurgery, 1993, 32, p. 376–383.
- ILLANES, S.D., ARAYA, P.C., FERRER, S.D., et al. *Rhomboencephalitis caused by listeria monocytogenes infection. Report of three cases.* Rev Med Chil, 2003, 131, p. 921–928.
- IQBAL, J., DAVIS, L.E., ORRISON jr., W.W. *An MRI study of lumbar puncture headaches.* Headache, 1995, 35, p. 420–422.
- JACKSON, I.J., SNODGRASS, S.R. *Peritoneal shunts in the treatment of hydrocephalus and increased intracranial pressure: a 4-year survey of 62 patients.* J Neurosurg, 1955, 12, p. 216–222.

- JANSSEN, J.C. *The prevalence of oligoclonal bands in the CSF of patients with primary neurodegenerative dementia.* J Neurol, 2004, 251 (2), p. 184–188.
- JIRÁSEK, A. *Česká a slovenská chirurgie v letech 1898–1945.* Praha : Státní zdravotnické nakladatelství, 1956.
- JIRÁSKOVÁ, N., ROZSÍVAL, P. *Výsledky 62 dekompresí obalů zrakového nervu.* Čs Oftal, 1999, 55, s. 136–144.
- JOHNSON, J.C., DEEB, ZL. *Coaxial needle technique for lumbar puncture in the morbidly obese patient.* Radiology, 1991, 179, p. 874.
- KALA, M., VAVERKA, M., BUŘVAL, S. *Hnisavé procesy v páteřním kanále – klinické zkušenosti.* Čes a Slov Neurol Neurochir, 1997, 60/93, s. 288–292.
- KALA, M. *Měření nitrolebečního tlaku a Pascalův zákon.* Ces Slov Neurol Neurochir, 2007, 70, 103 (2), s. 340–341.
- KALA, M. *Hydrocefalus.* Praha : Galén, 2005.
- KANG, S. *Efficacy of lumbo-peritoneal versus ventriculo-peritoneal shunting for management of chronic hydrocephalus following aneurysmal subarachnoid haemorrhage.* Acta Neurochir, 2000, 142, p. 45–49.
- KARAOŠMANOĞLU, D., OKTAR, SÖ., ARAÇ, M., ERBAŞ, G. *Portal and systemic venous gas in a patient after lumbar puncture.* Br J Radiol, 2005, 78, p. 767–769.
- KATZMAN, R., HUSSEY, F. *A simple constant-infusion manometric test for measurement of CSF absorption. I. Rationale and method.* Neurology, 1970, 20, p. 534–544.
- KELBICH, P., SLAVÍK, S., JASANSKÁ, J., et al. *Hodnocení energetických poměrů v likvorovém kompartmentu pomocí vyšetřování vybraných parametrů metabolismu glukosy v CSF.* Klin Biochem Metab, 1998, 6 (27), s. 213–225.
- KELBICH, P., JÍLEK, D., KRÁL, V., et al. *Likvorové nálezy u pacientů s neurosyfilis.* Klin Biochem Metab, 2000, 8 (29), s. 229–239.
- KELBICH, P., KOUDELKOVÁ, M., MACHOVÁ, H., et al. *Význam urgentního vyšetření mozkomíšního moku pro včasnou diagnostiku neuroinfekcí.* Klin mikrobiol inf lék, 2007, 13 (1), s. 9–20.
- KESSLER, LA., NOVELLI, PM., REIGEL, DH. *Surgical treatment of benign intracranial hypertension – subtemporal decompression revisited.* Surg Neurol, 1998, 50, p. 73–76.
- KINNUNEN, E., LIMONEN, T., PIRTILÄ, T., et al. *Effects of recombinant alpha-2b-interferon therapy in patients with progressive MS.* Acta Neurol Scand, 1993, 87 (6), p. 457–460.
- KNEEN, R., SOLOMON, T., APPLETON, R. *The role of lumbar puncture in children with suspected central nervous system infection.* BMC Pediatr, 2002, 2, p. 8.
- KNOBLOCH, J. *Obecná chirurgie. 6., část. přeprac. vyd.* Praha : Avicenum, 1973.
- KOLÁŘ, M., LÁTAL, T., ČERMÁK, P. *Klinicko-mikrobiologické podklady racionální antibiotické léčby.* Praha : Trios, 2002.
- KOLÁŘ, O. *Přehled likvorologie.* In KOLEKTIV AUTORŮ, *Přehled vyšetřovacích metod v neurologii.* Praha : Státní pedagogické nakladatelství Praha, 1966, s. 69–70.
- KOSTELJANETZ, M., NEHEN, AM., KAALUND, J. *Cerebrospinal fluid outflow resistance measurements in the selection of patients for shunt surgery in the normal pressure hydrocephalus syndrome. A controlled trial.* Acta Neurochir, 1990, 104, p. 48–53.

- KOUL, R., CHACKO, A., JAVED, H., et al. *Syndrome of cerebrospinal fluid hypovolemia following lumbar puncture cerebrospinal fluid leak in a patient with idiopathic intracranial hypertension.* J Child Neurol, 2002, 17, p. 77–79.
- KREJSEK, J., KOPECKÝ, O. *Klinická imunologie.* Hradec Králové : Nucleus, 2004.
- KRUGER, H., ENGLERT, D., PFLUGHAUPT, KW. *Demonstration of oligoclonal immunoglobulin G in Guillain-Barre syndrome and lymphocytic meningoradiculitis by isoelectric focusing.* J Neurol, 1981, 226 (1), p. 15–24.
- KUHLE, J., POHL, C., MEHLING, M., et al. *Lack of association between antimyelin antibodies and progression to multiple sclerosis.* N Engl J Med, 2007, 356 (4), p. 371–378.
- LAI, PH., HO, JT., CHEN, WL., et al. *Brain abscess and necrotic brain tumor: Discrimination with proton MR spectroscopy and diffusion-weighted imaging.* AJNR Am J Neuroradiol, 2002, 23, p. 1369–1377.
- LEE, AC., LAU, Y., LI, CH., et al. *Intraspinal and intracranial hemorrhage after lumbar puncture.* Pediatr Blood Cancer, 2005, 48, p. 233–237.
- LENNON, VA., WINGERCHUK, DM., KRYMR, TJ., et al. *A serum autoantibody marker of neuromyelitis optica: distinction from multiple sclerosis.* Lancet, 2004, 364, p. 2106–2112.
- LEVINE, JF., HIESIGER, EM., WHELAN, MA., et al. *Pneumococcal meningitis associated with retroperitoneal abscess.* JAMA, 1982, 248, p. 2308–2309.
- LEVY, DI., REKATE, HL., CHERNY, WB., et al. *Controlled lumbar drainage in pediatric head injury.* J Neurosurg, 1995, 83, p. 453–460.
- LEWCZUK, P., KORNHUBER, J., VANDERSTICHELE, H., et al. *Multiplexed quantification of dementia biomarkers in the CSF of patients with early dementias and MCI: a multicenter study.* Neurobiol Aging, 2008, 29 (6), p. 812–818.
- LIN, LC., LIAO, YM., YANG, RC. *Pneumorrhachis following lumbar puncture: a case report.* Kaohsiung J Med Sci, 2006, 22, p. 39–43.
- LINK, H., TIBBLING, G. *Principles of albumin and IgG disorders. Evaluation of IgG synthesis within the central nervous system in multiple sclerosis.* Scand J Clin Lab Invest, 1977, 37, p. 397–401.
- LINKER, G., MIRZA, N., MANETTI, BA., et al. *Fine-needle, negative-pressure lumbar puncture: A safe technique for collecting CSF.* Neurology, 2002, 59, p. 2008–2009.
- LUNDAR, T., NONES, H. *Determination of ventricular fluid outflow resistance in patients with ventriculomegaly.* J Neurol Neurosurg Psychiatry, 1990, 53, p. 896–898.
- LYNCH, J., KRINGS-ERNST, I., STRICK, K., et al. *Use of a 25 gauge Whitacre needle to reduce the incidence of post-dural puncture headache.* Br J Anaesth, 1991, 67, p. 692–693.
- MACDONALD, A., CHATRATH, P., SPECTOR, T., et al. *Level of termination of the spinal cord and the dural sac: a magnetic resonance study.* Clin Anat, 1999, 12, p. 149–152.
- MAREŠ, J., MUCHOVÁ, B., HERZIG, R., et al. *Změny vzorce oligoklonálních proužků v likvoru nemocných s roztroušenou sklerózou během terapie interferony β , resp. glatiramer acetátem. Pilotní studie.* Čes a slov Neurol Neurochir, 2003, 66/99 (2), s. 118–120.
- MAREŠ, J., HERZIG, R., URBÁNEK, K., et al. *Využití stanovení cystatinu C u pacientů s neurodegenerativními chorobami.* Čes a slov Neurol Neurochir, 2006, 69/102 (3), s. 195–199.
- MARTÍN-MILLÁN, M., HERNÁNDEZ, JL., MATORRAS, P., et al. *Multiple subdural haematomas following lumbar puncture.* Eur Neurol, 2005, 53, p. 159–160.

- MASSERMAN, JH. *Cerebrospinal hydrodynamics. IV. Clinical experimental studies.* Arch Neurol Psychiat, 1934, 32, p. 523.
- MAYER, M. Principy hodnocení likvorologického nálezu. In URBÁNEK, K., et al., *Přehled vyšetřovacích metod v neurologii.* Olomouc : Univerzita Palackého, 1996, s. 112–122.
- MAZOR, SS., MCNULTY, JE., ROOSEVELT, GE. *Interpretation of traumatic lumbar punctures: Who can go home?* Pediatrics, 2003, 111, p. 525–528.
- MCDONALD, JV., KLUMP, TE. *Epidermoid spinal cord tumors caused by lumbar puncture.* Arch Neurol, 1986, 43, p. 936–939.
- MCDONALD, WI., COMPSTON, A., EDAN, G., et al. *Recommended diagnostic criteria for multiple sclerosis: guidelines from the International Panel on the diagnosis of multiple sclerosis.* Ann Neurol, 2001, 50, p. 121–127.
- MCLEAN, BN., LUXTON, RW., THOMPSON, EJ. *A study of immunoglobulin G in the cerebrospinal fluid of 1007 patients with suspected neurological disease using isoelectric focusing and the Log IgG-Index. A comparison and diagnostic applications.* Brain, 1990, 113, p. 1269–1289.
- MCLONE, DG. *Comment.* Neurosurgery, 1993, 32, p. 383.
- MEIER, U., ZEILINGER, FS., SCHÖNHERR, B. *Endoscopic ventriculostomy versus shunt operation in normal pressure hydrocephalus: Diagnostics and indications.* Minim Invas Neurosurg, 2000, 43, p. 87–90.
- MENGE, T., LALIVE, PH., VON BUDINGEN, HC., et al. *Antibody responses against galactocerebroside are potential stage-specific biomarkers in multiple sclerosis.* J Allergy Clin Immunol, 2005, 116, p. 453–459.
- Mezinárodní soustava jednotek SI ve zdravotnictví.* Praha : Avicenum, 1979.
- MILHORAT, TH. *Failure of choroid plexectomy as a treatment for hydrocephalus.* Surg Gynecol Obstet, 1974, 193, p. 505.
- MILLER, JR., BURKE, AM., BEVER, CT. *Occurrence of oligoclonal bands in multiple sclerosis and other CNS diseases.* Ann Neurol, 1983, 13 (1), p. 53–58.
- MISHRA, AM., GUPTA, RK., JAGGI, RS., et al. *Role of diffusion-weighted imaging and in vivo proton magnetic resonance spectroscopy in the differential diagnosis of ring-enhancing intracranial cystic mass lesions.* J Comput Assist Tomogr, 2004, 28, p. 540–547.
- MIYAKE, S., KOBAYASHI, N., MURAI, N., et al. *Acquired lumbar epidermoid cyst in an adult. Case report.* Neurol Med Chir (Tokyo), 2005, 45, p. 277–279.
- MORI, T., MORIMOTO, K., HAYAKAWA, T., et al. *Radioimmunoassay of Astroprotein (An Astrocyte-Specific Cerebroprotein) in Cerebrospinal Fluid and Its Clinical Significance.* Neurol Med Chir (Tokyo), 1978, 18, p. 25–31.
- MULDOON, T. *Obtaining fluid samples and measuring intrathecal pressure may require different approaches.* BMJ, 1998, 316, p. 1018.
- MUMENTHALER, M., MATTLE, H. *Neurologie.* 1. čes. vyd. Praha : Grada Publishing, 2001.
- MURSHID, WR. *Ventricular failure as a cause of unsuccessful endoscopic third ventriculostomy.* Minim Invas Neurosurg, 2002, 45, p. 65–71.
- NAKAISO, M., UNO, M., HARADA, M., et al. *Brain abscess and glioblastoma identified by combined proton magnetic resonance spectroscopy and diffusion-weighted magnetic resonance imaging.* Neurol Med Chir (Tokyo), 2002, 42, p. 346–348.

- NEL, MR., ROBINSON, N. *Epidural blood patching can be used to treat headache*. BMJ, 1998, 316, p. 1019.
- NELSON, JR., GOODMAN, SJ. *An evaluation of the cerebrospinal fluid infusion test for hydrocephalus*. Neurology, 1971, 21, p. 1037–1053.
- O'FARELL, R., THORNTON, J., BRENNAN, P., et al. *Spinal cord infarction and tetraplegia - rare complications of meningococcal meningitis*. Br J Anaesth, 2000, 84, p. 514–517.
- O'CONNOR, KC., CHITNIS, T., GRIFFIN, DE., et al. *Myelin basic protein-reactive autoantibodies in the serum and cerebrospinal fluid of multiple sclerosis patients are characterized by low-affinity interactions*. J Neuroimmunol, 2003, 136, p. 140–148.
- OERTEL, MF., KORINTH, MC., REINGES, MH., et al. *Pathogenesis, diagnosis and management of pneumorrhachis*. Eur Spine J, 2006, 15, Suppl. 17, p. 636–643.
- OLIVER, WJ., SHOPE, TC., KUHNS, LR. *Fatal lumbar puncture: Fact versus fiction-an approach to a clinical dilemma*. Pediatrics, 2003, 112, p. 174–176.
- OWLER, BK., PARKER, G., HALMAGYI, GM., et al. *Pseudotumor cerebri syndrome: venous sinus obstruction and its treatment with stent placement*. J Neurosurg, 2003, 98, p. 1045–1055.
- OYELESE, AA., STEINBERG, GK., HUHNS, SL., WIJMAN, CAC. *Paradoxical cerebral herniation secondary to lumbar puncture after decompressive craniectomy for large space-occupying hemispheric stroke: Case report*. Neurosurgery, 2005, 57, p. 594.
- PADMANABHAN, R., CROMPTON, D., BURN, D., BIRCHALL, D. *Acquired Chiari 1 malformation and syringomyelia following lumboperitoneal shunting for pseudotumor cerebri*. J Neurol Neurosurg Psychiatry, 2005, 76, p. 298.
- PATIN, DJ., ECKSTEIN, EC., HARUM, K., PALLARES, VS. *Anatomic and biomechanical properties of human lumbar dura mater*. Anesth Analg, 1993, 76, p. 535–540.
- PEARCE, JMS. *Queckenstedt's manoeuvre*. J Neurol Neurosurg Psychiatry, 2006, 77, p. 728.
- PERICOT, I., RÍO, J., ROVIRA, A., et al. *Rombencefalitis por listeria. Hallazgos neurorradiológicos*. Neurología, 2003, 18, p. 34–37.
- PETERSON, MA., ABELE, J. *Bedside ultrasound for difficult lumbar puncture*. J Emerg Med, 2005, 28, p. 197–200.
- PORTER, RW., HIBBERT, C., WELLMAN, P. *Backache and the lumbar spinal canal*. Spine, 1980, 5, p. 99–105.
- PROCHÁZKA, J. *Současné možnosti látek používaných k intratékální analgezi*. Bolest, 2001, 4, s. 220–231.
- QUINCKE, HI. *Die lumbalpunktion des Hydrocephalus*. Berl Klin Wochenschr, 1891, 28, S. 929–933, S. 965–968.
- RABIN, BM., ROYCHOWDHURY, S., MEYER, JR., et al. *Spontaneous intracranial hypotension: spinal MRI findings*. AJNR, 1998, 19, p. 1034–1039.
- RAMENGGHI, LA., WEBB, AV., SHEVLIN, PM., et al. *Intra-oral administration of sweet-tasting substances and infants' crying response to immunization: a randomized, placebo-controlled trial*. Biol Neonate, 81, 2002, p. 163–169.
- RASTOGI, S., LIEBESKIND, DS., ZAGER, EL., et al. *Rapid cognitive decline following lumbar puncture in a patient with a dural arteriovenous fistula*. Surg Neurol, 2004, 62, p. 341–345.

- REIBER, H., UNGEFEHR, S., JACOBI, CH. *The intrathecal, polyspecific and oligoclonal immune response in multiple sclerosis*. Mult Scler, 1998, 4, p. 111–117.
- REIBER, H., OTTO, M., TRENDELENBURG, CH., et al. *Reporting Cerebrospinal Fluid Data: Knowledge Base and Interpretation Software*. Clin Chem Lab Med, 2001, 39 (4), p. 324–332.
- REIBER, H., PETER, JB. *Cerebrospinal fluid analysis: disease-related data patterns and evaluation programs*. J Neurol Sci, 2001, 1, p. 101–122.
- REINA, MA., LOPEZ, A., VILLANUEVA, MC., et al. *Possibility of cauda equina nerve root damage from lumbar punctures performed with 25-gauge Quincke and Whitacre needles*. Rev Esp Anesthesiol Reanim, 2005, 52, p. 267–275.
- REINDL, M., LININGTON, C., BREHM, U., et al. *Antibodies against the myelin oligodendrocyte glycoprotein and the myelin basic protein in multiple sclerosis and other neurological diseases: a comparative study*. Brain, 1999, 122, p. 2047–2056.
- REKATE, HL. *Classification of slit – ventricle syndrome using intracranial pressure monitoring*. Pediatr Neurosurg, 1993, 19, p. 15–20.
- REKATE, HL. *Comment*. Neurosurgery, 1993, 32, p. 383.
- RENDER, CA. *The reproducibility of the iliac crest as a marker of lumbar spine level*. Anaesthesia, 1996, 51, p. 1070–1071.
- REYNOLDS, F., O'SULLIVAN, G. *»Atraumatic needle« is a better term than »blunt needle«*. BMJ, 1998, 316, p. 1018.
- RICKMANN, H., OBHOF, W., RAUH, J., DRUSCHKY, K.-F. *Rhombenzephalitis durch Listeria monocytogenes*. Nervenarzt, 1994, 65, S. 557–559.
- RIEDL, O. *Základy punkční techniky*. In FUČÍK, M., et al., *Základní vyšetření ve vnitřním lékařství*. 2., oprav. a dopln. vyd. Praha : Avicenum, 1978, s. 377–385.
- RIFAAT, M., EL-SHAFAI, I., SAMRA, K., SOROUR, O. *Intramedullary spinal abscess following spinal puncture. Case report*. J Neurosurg, 1973, 38, p. 366–367.
- ROSENBLUM, ML., HOFF, JT., NORMAN, D. *Nonoperative treatment of brain abscesses in selected high-risk patients*. J Neurosurg, 1980, 52, p. 217–225.
- RUBIN, RC., HENDERSON, ES., OMMAYA, AK., et al. *The production of cerebrospinal fluid in man and its modification by acetazolamide*. J Neurosurg, 1966, 25, p. 430–436.
- RUDICK, RA., COOKFAIR, DL., SIMONIAN, NA., et al. *Cerebrospinal fluid abnormalities in a phase III trial of Avonex (IFN β -1a) for relapsing multiple sclerosis. The Multiple Sclerosis Collaborative Research Group*. J Neuroimmunol, 1999, 93 (1–2), p. 8–14.
- RUPP, SM., WILSON, CB. *Treatment of spontaneous cerebrospinal fluid leak with epidural blood patch. Case report*. J Neurosurg, 1989, 70, p. 808–810.
- RUTKA, JT. *Cerebrospinal fluid production in patients with hydrocephalus*. J Neurosurg, 2002, 97, p. 1269–1270.
- SAHEBKAR-MOGHADDAM, F., ADORNATO, BT. *The failed lumbar puncture*. Neurology, 2005, 64, p. E24.
- SAIFUDDIN, A., BURNETT, SJ., WHITE, J. *The variation of the position of the conus medullaris in an adult population. A magnetic resonance imaging study*. Spine, 1998, 23, p. 1452–1456.
- SAJJADI, SA., HARIRCHIAN, MH., SHEIKHBAHAIEI, N., et al. *The relation between intracranial and intraocular pressures: Study of 50 patients*. Ann Neurol, 2006, 59, p. 867–870.

- SAMADANI, U., HUANG, JH., BARANOV, D., et al. *Intracranial hypotension after intraoperative lumbar cerebrospinal fluid drainage*. Neurosurgery, 2003, 52, p. 148–152.
- SANDOVAL, M., SHESTAK, W., STÜRMAN, K., HSU, C. *Optimal patient position for lumbar puncture, measured by ultrasonography*. Emerg Radiol, 2004, 10, p. 179–181.
- SATHI, S., STIEG, PE. *«Acquired» Chiari I malformation after multiple lumbar punctures: Case report*. Neurosurgery, 1993, 32, p. 306–309.
- SERPELL, MG., RAWAL, N. *Headaches after diagnostic dural punctures*. BMJ, 2000, 321, p. 973–974.
- SHIRAISHI, N., MATSUMURA, G. *What is the true location of Jacoby's line?* Okajimas Folia Anat Jpn, 2006, 82, p. 111–115.
- SHIRAISHI, N., MATSUMURA, G. *Establishing intercrestal line by posture: a radiographic evaluation*. Okajimas Folia Anat Jpn, 2006, 82, p. 139–146.
- SCHALTENBRAND, G. *Neuere Anschauungen zur Pathophysiologie der Liquorzirkulation*. Zentralbl Neurochir, 1938, 3, S. 290–299.
- SCHIFFER, P., PODVINEC, M., HASLER, P. *Hearing loss after lumbar myelography*. Schweiz Rundsch Med Prax, 2006, 95, S. 855–857.
- SCHMIDT, B., CZOSNYKA, M., SCHWARZE, JJ., et al. *Evaluation of a method for noninvasive intracranial pressure assessment during infusion studies in patients with hydrocephalus*. J Neurosurg, 2000, 92, p. 793–800.
- SCHWARZ, M., SPECTOR, L., GORTLER, M., et al. *Serum anti-Glc(alpha1,4)Glc(alpha) antibodies as a biomarker for relapsing-remitting multiple sclerosis*. J Neurol Sci, 2006, 244, p. 59–68.
- SILBER, E., SEMRA, YK., GREGSON, NA., SHARIEF, MK. *Patients with progressive multiple sclerosis have elevated antibodies to neurofilament subunit*. Neurology, 2002, 58 (9), p. 1372–1381.
- SILVERBERG, GD., HUHN, S., JAFFE, RA., et al. *Downregulation of cerebrospinal fluid production in patients with chronic hydrocephalus*. J Neurosurg, 2002, 97, p. 1271–1275.
- SINDIC, CHJM., VAN ANTWERPEN, MP., GOFFETTE, S. *The Intrathecal Humoral Immune Response: Laboratory Analysis and Clinical Relevance*. Clin Chem Lab, 2001, 39 (4), p. 333–340.
- SINĚLNIKOV, RD. *Atlas anatomie člověka. Svazek třetí*. Praha : Státní zdravotnické nakladatelství, 1965.
- SOTELO, J., RUBALCAVA, MA., GOMEZ-LLATA, S. *A new shunt for hydrocephalus that relies on CSF production rather than on ventricular pressure: initial clinical experiences*. Surg Neurol, 1995, 43, p. 324–332.
- SPETZLER, RE., WILSON, CB., GROLLMUS, JM. *Percutaneous lumboperitoneal shunt. Technical note*. J Neurosurg, 1975, 43, p. 770–773.
- SRIRAM, S., STRATTON, CW., YAO, S., et al. *Chlamydia pneumoniae infection of the central nervous system in multiple sclerosis*. Ann Neurol, 1999, 46 (1), p. 6–14.
- STAEBLER, M., AZZI, N., SEKHARA, T., et al. *Complications of lumbar puncture in a child treated for leukaemia*. Pediatr Radiol, 2005, 35, p. 1121–1124.
- STINGL, J., LEBEDA, P., MARKOVÁ, H., ŠTĚŇHOVÁ, H. *Poznámky ke klinické anatomii páteřního kanálu*. Bolest, 2000, 3, s. 4–9.
- STRACHAN, A., TRAIN, J. *Aspirating cerebrospinal fluid speeds up procedure*. BMJ, 1998, 316, p. 1018–1019.

- SUGERMAN, HJ. *Lumboperitoneal shunting for pseudotumor cerebri. To the Editor.* Neurology, 1998, 50, p. 835–836.
- SUCHOMEL, P., MAJDAN, M., NESTÁVAL, A. *Význam izotopové cisternografie pro indikaci zkratové operace u komunikujícího hydrocefalu.* Čs Neurol Neurochir, 1993, 56/89, s. 150–154.
- SULLIVAN, HC. *Fatal tonsillar herniation in pseudotumor cerebri.* Neurology, 1991, 41, p. 1142–1144.
- ŠTOURÁČ, P. *Diferenciálně-diagnostický význam funkce hematolikvorové bariéry a oligoklonální syntézy IgG u sclerosis multiplex.* Klin Biochem Metab, 1998, 4 (6/27), s. 210–212.
- ŠTOURÁČ, P. *Oligoklonální IgG syntéza u autoimunitních neurologických paraneoplastických syndromů.* Klin Biochem Metab, 2000, 4 (8/29), s. 224–228.
- TANAKA, K., WAGA, S., SHIMOSAKA, S. *Papilledema and spinal cord tumors.* Surg Neurol, 1988, 29, p. 462–466.
- THOENNISSSEN, J., HERKNER, H., LANG, W., et al. *Does bed rest after cervical or lumbar puncture prevent headache? A systematic review and meta-analysis.* CMAJ, 2001, 165, p. 1311–1316.
- THOMAS, SR., JAMIESON, DRS., MUIR, KW. *Randomised controlled trial of atraumatic versus standard needles for diagnostic lumbar puncture.* BMJ, 2000, 321, p. 986–990.
- TICHÝ, J. *Klinická biochemie a likvorologie.* In NEVŠÍMALOVÁ, S., RŮŽIČKA, E., TICHÝ, J., et al., *Neurologie.* Praha : Galén, 2002, s. 145–150.
- TICHÝ, J. *Záněty CNS.* In NEVŠÍMALOVÁ, S., RŮŽIČKA, E., TICHÝ, J. et al., *Neurologie.* Praha : Galén, 2002, s. 235–246.
- TOMOSVARI, A., MENCSEK, Z., FUTO, J., et al. *Preliminary experience with controlled lumbar drainage in medically refractory intracranial hypertension.* Orv Hetil, 2005, 146, p. 159–164.
- TOURTELLOTTE, WW. *On cerebrospinal fluid immunoglobulin-G (IgG) quotients in multiple sclerosis and other diseases. A review and a new formula to estimate the amount of IgG synthesized per day by the central nervous system.* J Neurol Sci, 1970, 10, p. 279–304.
- TRUPP, M. *Stylet injury syndrome.* JAMA, 1977, 237, p. 2524.
- TUBBS, RS., SMYTH, MD., WELLONS 3rd, JC., OAKES, WJ. *Intramedullary hemorrhage in a neonate after lumbar puncture resulting in paraplegia: A case report.* Pediatrics, 2004, 113, p. 1403–1405.
- TURNBULL, D., MCKEVITT, F. *Letter.* BMJ, 2001, 322, p. 994.
- ULDRY, PA., KUNTZER, T., BOGOUSSLAWSKY, J., et al. *Early symptoms and outcome of Listeria monocytogenes rhomboencephalitis: 14 adult cases.* J Neurol, 1993, 240, p. 235–242.
- VAN DELLEN, JR., BILL, PLA. *Lumbar puncture- an innocuous diagnostic procedure?* S Afr Med J, 1978, 53, p. 666–668.
- VANĚK, P., SAMEŠ, M. *Onemocnění syndromem normotenzního hydrocefalu – význam lumbálního infúzního testu pro diagnostiku a efekt zkratové operace (Analýza souboru pacientů z let 1992–1999).* Čes a Slov Neurol Neurochir, 2002, 65/98, s. 256–263.
- VANĚK, P., SAMEŠ, M. *Lumbální infúzní test a transkraniální dopplerometrie v diagnostice a léčbě normotenzního hydrocefalu. Prospektivně hodnocená sestava pacientů 2000–2002.* Čas Lék čes, 2003, 142, s. 545–550.
- VAPHIADES, MS., EGGENBERGER, ER., MILLER, NR., et al. *Resolution of papilledema after neurosurgical decompression for primary Chiari I malformation.* Am J Ophthalmol, 2002, 133, p. 673–678.

- VAVERKA, M., KALA, M., BUŘVAL, S. *Význam CT cisternografie při léčbě mokových píštělí*. Čes Radiol, 1995, 49, s. 112–116.
- VERTOSICK jr., F. *When the air hits your brain*. New York, London : W.W. Norton & Company, 1996.
- WANG, JC., SPENCINER, D., ROBINSON, JC. *SPIRE spinous process stabilisation plate: biomechanical evaluation of a novel technology*. J Neurosurg Spine, 2006, 4, p. 160–164.
- WANIFUCHI, H., SHIMIZU, T., MARUYAMA, T. *Age-related changes in the proportion of intracranial cerebrospinal fluid space measured using volumetric computerized tomography scanning*. J Neurosurg, 2002, 97, p. 607–610.
- WARD, E., GUSHURST, CA. *Uses and technique of pediatric lumbar puncture*. Am J Dis Child, 1992, 146, p. 1160–1165.
- WEKERLE, H., HOHLFELD, R. *Molecular mimicry in multiple sclerosis*. N Engl J Med, 2003, 349, p. 185–186.
- WELCH, K. *The intracranial pressure in infants*. J Neurosurg, 1980, 52, p. 693–699.
- WIRTZ, PW., BLOEM, BR., VAN DER MEER, FJM., BROUWER, OF. *Paraparesis after lumbar puncture in a male with leukemia*. Pediatr Neurol, 2000, 23, p. 67–68.
- YENTIS, S. *Replacing the stylet before withdrawing the needle may help to prevent headache*. BMJ, 1998, 316, p. 1018.
- ZARZUR, E. *Genesis of the »true« negative pressure in the lumbar epidural space. A new hypothesis*. Anaesthesia, 1984, 39, p. 1101.
- ZUNIGA, FA., SHI, G., HALLER, JF., et al. *A three-dimensional model of the human facilitative glucose transporter glut1*. J Biol Chem, 2001, 276 (48), p. 44970–44975.