

References

Annersten, H. 1974: Mössbauer studies of natural biotites. Amer. Min. 59, 143-151.

Annersten, H. 1975: Mössbauer study of iron in natural and synthetic biotites. Fortschr. Miner. 52, 583-590.

Ericsson, T. & Wäppling, R. 1976: Texture effects in $3/2-1/2$ Mössbauer spectra. J. Phys. 37:C6, 719-723.

- Eugster, H. 1957: Heterogeneous reactions involving oxidation and reduction at high pressures and temperatures. J. Chem. Phys. 26, 1760.
- Eugster, H. & Wones, D. 1962: Stability relation of the ferrigenous biotite. annite. J. Petrol. 3, 82-125.
- Hazen, R. & Wones, D. 1972: The effect of cation substitutions on the physical parameters of trioctahedral micas. Amer. Min. 57, 103-129.
- Heller-Kallai, L. & Rozenson, I. 1981: The use of Mössbauer spectroscopy of iron in clay mineralogy. Phys. Chem. Min. 7, 223-238.
- Hogg, C. S. & Meads, R. E. 1975: A Mössbauer study of thermal decomposition of biotites. Min. Mag. 40, 79-88.
- Häggström, L., Wäppling, R. & Annersten, H. 1969: Mössbauer study of iron rich biotites. Chem. Phys. Lett. 4, 107-108.
- Radoslovich, E. W. & Norrish, K. 1962: The cell dimensions and symmetry of layer lattice silicates. I. Some structural considerations. Amer. Min. 47, 599-616.
- Rutherford, M. J. 1973: The phase relations of aluminous iron biotites in the system KAlSi_3O_8 - KAlSiO_4 - Al_2O_3 -Fe-O-H. J. Petrol. 14, 159-180.
- Wones, D. 1963: Phase equilibria of ferriannite, $\text{KFe}_3\text{FeSi}_3\text{O}_{10}(\text{OH})_2$. Amer. J. Sci. 261, 581-596.

