

9 Referenční seznam

- AACE/ACE Obesity Task Force. (2012). AACE/ACE position statement on the prevention, diagnosis, and treatment of obesity. *Endocrine Practice*, 18(5), 642-648. doi:10.4158/EP12160.PS
- Adams, M., Frank, L., Schipperijn, J., Smith, G., Chapman, J., Christiansen, L., . . . Sallis, J. (2014). International variation in neighborhood walkability, transit, and recreation environments using geographic information systems: the IPEN adult study. *International Journal of Health Geographics*, 13(1), 43. doi:10.1186/1476-072X-13-43
- Afzal, S., Tybjaerg-Hansen, A., Jensen, G. B., & Nordestgaard, B. G. (2016). Change in body mass index associated with lowest mortality in Denmark, 1976-2013. *JAMA*, 315(18), 1989-1996. doi:10.1001/jama.2016.4666
- Ainsworth, B. E., Haskell, W. L., Herrmann, S. D., Meckes, N., Bassett, D. R., Jr., Tudor-Locke, C., . . . Leon, A. S. (2011). 2011 Compendium of physical activities: a second update of codes and MET values. *Medicine and Science in Sports and Exercise*, 43(8), 1575-1581. doi:10.1249/MSS.0b013e31821ece12
- Albanese, C. V., Diessel, E., & Genant, H. K. (2003). Clinical applications of body composition measurements using DXA. *Journal of Clinical Densitometry*, 6(2), 75-85. doi:10.1385/JCD:6:2:75
- Anderson, L. J., Erceg, D. N., & Schroeder, E. T. (2012). Utility of multifrequency bioelectrical impedance compared with dual-energy x-ray absorptiometry for assessment of total and regional body composition varies between men and women. *Nutrition Research*, 32(7), 479-485. doi:10.1016/j.nutres.2012.05.009
- Arden, N. K., & Spector, T. D. (1997). Genetic influences on muscle strength, lean body mass, and bone mineral density: a Twin Study. *Journal of Bone and Mineral Research*, 12(12), 2076-2081. doi:10.1359/jbmr.1997.12.12.2076
- Atkin, L.-M., & Davies, P. S. (2000). Diet composition and body composition in preschool children. *American Journal of Clinical Nutrition*, 72(1), 15-21.
- Babb, T. G., Wyrick, B. L., Chase, P. J., DeLorey, D. S., Rodder, S. G., Feng, M. Y., & Ranasinghe, K. G. (2011). Weight loss via diet and exercise improves exercise breathing mechanics in obese men. *Chest*, 140(2), 454-460. doi:10.1378/chest.10-1088
- Baker, G., Gray, S., Wright, A., Fitzsimons, C., Nimmo, M., Lowry, R., & Mutrie, N. (2008). The effect of a pedometer-based community walking intervention "Walking for Wellbeing in the West" on physical activity levels and health outcomes: a 12-week randomized controlled trial. *International Journal of Behavioral Nutrition and Physical Activity*, 5(1), 44. doi:10.1186/1479-5868-5-44
- Barbosa-Silva, M. C. G., & Barros, A. J. (2005). Bioelectrical impedance analysis in clinical practice: a new perspective on its use beyond body composition equations. *Current Opinion in Clinical Nutrition and Metabolic Care*, 8(3), 311-317.
- Barbosa-Silva, M. C. G., Barros, A. J., Wang, J., Heymsfield, S. B., & Pierson, R. N. (2005). Bioelectrical impedance analysis: population reference values for phase angle by age and sex. *American Journal of Clinical Nutrition*, 82(1), 49-52.

- Barnett, T. A., O'Loughlin, J., Sabiston, C. M., Karp, I., Bélanger, M., Van Hulst, A., & Lambert, M. (2010). Teens and screens: the influence of screen time on adiposity in adolescents. *American Journal of Epidemiology*, *172*(3), 255-262. doi:10.1093/aje/kwq125
- Barreira, T. V., Harrington, D. M., Staiano, A. E., Heymsfield, S. B., & Katzmarzyk, P. T. (2011). Body adiposity index, body mass index, and body fat in white and black adults. *JAMA*, *306*(8), 828-830. doi:10.1001/jama.2011.1189
- Barros, A. J., Santos, L. P., Wehrmeister, F., Motta, J. V., Matijasevich, A., Santos, I. S., . . . Barros, F. C. (2017). Caesarean section and adiposity at 6, 18 and 30 years of age: results from three Pelotas (Brazil) birth cohorts. *BMC Public Health*, *17*(1), 256. doi:10.1186/s12889-017-4165-3
- Bassett, D. R., Troiano, R. P., McClain, J. J., & Wolff, D. L. (2015). Accelerometer-based physical activity: a total volume per day and standardized measures. *Medicine and Science in Sports and Exercise*, *47*(4), 833-838. doi:10.1249/mss.0000000000000468
- Baumgartner, R. N., Heymsfield, S. B., Lichtman, S., Wang, J., & Pierson, R. N. (1991). Body composition in elderly people: effect of criterion estimates on predictive equations. *American Journal of Clinical Nutrition*, *53*(6), 1345-1353.
- Beets, M. W., Bornstein, D., Dowda, M., & Pate, R. R. (2011). Compliance with national guidelines for physical activity in U.S. preschoolers: measurement and interpretation. *Pediatrics*, *127*(4), 658-664. doi:10.1542/peds.2010-2021
- Behnke, A. R. (1942). Physiologic studies pertaining to deep sea diving and aviation, especially in relation to the fat content and composition of the body: the Harvey lecture, March 19, 1942. *Bulletin of the New York Academy of Medicine*, *18*(9), 561-585.
- Behnke, A. R., Feen, B. G., & Welham, W. C. (1942). The specific gravity of healthy men: Body weight ÷ volume as an index of obesity. *JAMA*, *118*(7), 495-498. doi:10.1001/jama.1942.02830070001001
- Behnke, A. R., Osserman, E. F., & Welham, W. C. (1953). Lean body mass: Its clinical significance and estimation from excess fat and total body water determinations. *Archives of Internal Medicine*, *91*(5), 585-601. doi:10.1001/archinte.1953.00240170011002
- Belarmino, G., Gonzalez, M. C., Torrinhas, R. S., Sala, P., Andraus, W., D'Albuquerque, L. A., . . . Waitzberg, D. L. (2017). Phase angle obtained by bioelectrical impedance analysis independently predicts mortality in patients with cirrhosis. *World Journal of Hepatology*, *9*(7), 401-408. doi:10.4254/wjh.v9.i7.401
- Bera, T. K. (2014). Bioelectrical impedance methods for noninvasive health monitoring: a review. *Journal of Medical Engineering*, *2014*, 381251. doi:10.1155/2014/381251
- Berghöfer, A., Pischon, T., Reinhold, T., Apovian, C., Sharma, A., & Willich, S. (2008). Obesity prevalence from a European perspective: a systematic review. *BMC Public Health*, *8*(1), 200-210. doi:10.1186/1471-2458-8-200
- Bergman, R. N., Stefanovski, D., Buchanan, T. A., Sumner, A. E., Reynolds, J. C., Sebring, N. G., . . . Watanabe, R. M. (2011). A better index of body adiposity. *Obesity*, *19*(5), 1083-1089. doi:10.1038/oby.2011.38
- Bernhard, A. B., Scabim, V. M., Serafim, M. P., Gadducci, A. V., Santo, M. A., & Cleva, R. d. (2017). Modified body adiposity index for body fat estimation in severe obesity. *Journal of Human Nutrition and Dietetics*, *30*(2), 177-184. doi:10.1111/jhn.12404
- Berrington de Gonzalez, A., Hartge, P., Cerhan, J. R., Flint, A. J., Hannan, L., MacInnis, R. J., . . . Thun, M. J. (2010). Body-mass index and mortality among 1.46 million white adults. *New England Journal of Medicine*, *363*(23), 2211-2219. doi:10.1056/NEJMoa1000367
- Bhaskaran, K., Douglas, I., Forbes, H., dos-Santos-Silva, I., Leon, D. A., & Smeeth, L. (2014). Body-mass index and risk of 22 specific cancers: a population-based cohort study of 5.24 million UK adults. *Lancet*, *384*(9945), 755-765. doi:10.1016/S0140-6736(14)60892-8

- Bigaard, J., Frederiksen, K., Tjønneland, A., Thomsen, B. L., Overvad, K., Heitmann, B. L., & Sørensen, T. I. A. (2004). Body fat and fat-free mass and all-cause mortality. *Obesity Research, 12*(7), 1042-1049. doi:10.1038/oby.2004.131
- Bin, Y. S., Marshall, N. S., & Glozier, N. (2012). Secular trends in adult sleep duration: a systematic review. *Sleep Medicine Reviews, 16*(3), 223-230. doi:10.1016/j.smr.2011.07.003
- Biospace. (2008). InBody 720 – The precision body composition analyzer (User's Manual). Retrieved from <http://www.inbody.com/>
- Bland, J. M., & Altman, D. G. (2010). Statistical methods for assessing agreement between two methods of clinical measurement. *International Journal of Nursing Studies, 47*(8), 931-936. doi:10.1016/j.ijnurstu.2009.10.001
- Boneva-Asiova, Z., & Boyanov, M. (2011). Age-related changes of body composition and abdominal adipose tissue assessed by bioelectrical impedance analysis and computed tomography. *Endocrinología y Nutrición, 58*(9), 472-477. doi:10.1016/j.endonu.2011.07.004
- Bornstein, D. B., Beets, M. W., Byun, W., Welk, G., Bottai, M., Dowda, M., & Pate, R. (2011). Equating accelerometer estimates of moderate-to-vigorous physical activity: in search of the Rosetta Stone. *Journal of Science and Medicine in Sport, 14*(5), 404-410. doi:10.1016/j.jsams.2011.03.013
- Borrud, L. G., Flegal, K. M., Looker, A. C., Everhart, J. E., Harris, T. B., & Shepherd, J. A. (2010). Body composition data for individuals 8 years of age and older: U.S. population, 1999-2004. *Vital and Health Statistics. Series 11: Data from the National Health Survey, 250*, 1-87.
- Bosaeus, M., Karlsson, T., Holmäng, A., & Ellegård, L. (2014). Accuracy of quantitative magnetic resonance and eight-electrode bioelectrical impedance analysis in normal weight and obese women. *Clinical Nutrition, 33*(3), 471-477. doi:10.1016/j.clnu.2013.06.017
- Bossingham, M. J., Carnell, N. S., & Campbell, W. W. (2005). Water balance, hydration status, and fat-free mass hydration in younger and older adults. *American Journal of Clinical Nutrition, 81*(6), 1342-1350.
- Bosy-Westphal, A., Danielzik, S., Dörhöfer, R.-P., Later, W., Wiese, S., & Müller, M. J. (2006). Phase angle from bioelectrical impedance analysis: population reference values by age, sex, and body mass index. *Journal of Parenteral and Enteral Nutrition, 30*(4), 309-316. doi:10.1177/0148607106030004309
- Bouchard, C., & Perusse, L. (1988). Heredity and body fat. *Annual Review of Nutrition, 8*(1), 259-277. doi:10.1146/annurev.nu.08.070188.001355
- Bouchard, C., Simoneau, J. A., Lortie, G., Boulay, M. R., Marcotte, M., & Thibault, M. C. (1986). Genetic effects in human skeletal muscle fiber type distribution and enzyme activities. *Canadian Journal of Physiology and Pharmacology, 64*(9), 1245-1251. doi:10.1139/y86-210
- Bouchi, R., Asakawa, M., Ohara, N., Nakano, Y., Takeuchi, T., Murakami, M., . . . Ogawa, Y. (2016). Indirect measure of visceral adiposity 'A Body Shape Index' (ABSI) is associated with arterial stiffness in patients with type 2 diabetes. *BMJ Open Diabetes Research and Care, 4*(1). doi:10.1136/bmjdr-2015-000188
- Bradbury, K. E., Guo, W., Cairns, B. J., Armstrong, M. E. G., & Key, T. J. (2017). Association between physical activity and body fat percentage, with adjustment for BMI: a large cross-sectional analysis of UK Biobank. *BMJ Open, 7*(3), e011843. doi:10.1136/bmjopen-2016-011843
- Brage, S., Wedderkopp, N., Andersen, L. B., & Froberg, K. (2003). Influence of step frequency on movement intensity predictions with the CSA accelerometer: A field validation study in children. *Pediatric Exercise Science, 15*(3), 277-287.

- Brambilla, P., Bedogni, G., Heo, M., & Pietrobelli, A. (2013). Waist circumference-to-height ratio predicts adiposity better than body mass index in children and adolescents. *International Journal of Obesity*, 37(7), 943-946. doi:10.1038/ijo.2013.32
- Brantlov, S., Ward, L. C., Jødal, L., Rittig, S., & Lange, A. (2017). Critical factors and their impact on bioelectrical impedance analysis in children: a review. *Journal of Medical Engineering and Technology*, 41(1), 22-35. doi:10.1080/03091902.2016.1209590
- Bravata, D. M., Smith-Spangler, C., Sundaram, V., & et al. (2007). Using pedometers to increase physical activity and improve health: A systematic review. *JAMA*, 298(19), 2296-2304. doi:10.1001/jama.298.19.2296
- Brazendale, K., Beets, M. W., Bornstein, D. B., Moore, J. B., Pate, R. R., Weaver, R. G., . . . van Sluijs, E. M. F. (2016). Equating accelerometer estimates among youth: The Rosetta Stone 2. *Journal of Science and Medicine in Sport*, 19(3), 242-249. doi:10.1016/j.jsams.2015.02.006
- Breij, L. M., Steegers-Theunissen, R. P., Briceno, D., & Hokken-Koelega, A. C. (2015). Maternal and fetal determinants of neonatal body composition. *Hormone Research in Paediatrics*, 84(6), 388-395. doi:10.1159/000441298
- Brožek, J. (1965). *Human body composition: approaches and applications* (1st ed.). Oxford, New York,: Symposium Publications Division.
- Brožek, J., Grande, F., Anderson, J. T., & Keys, A. (1963). Densitometric analysis of body composition: revision of some quantitative assumptions. *Annals of the New York Academy of Sciences*, 110(1), 113-140. doi:10.1111/j.1749-6632.1963.tb17079.x
- Brožek, J., & Prokopec, M. (2001). Historical note: Early history of the anthropometry of body composition. *American Journal of Human Biology*, 13(2), 157-158. doi:10.1002/1520-6300(200102/03)13:2<157::aid-ajhb1023>3.0.co;2-l
- Buffa, R., Saragat, B., Cabras, S., Rinaldi, A. C., & Marini, E. (2013). Accuracy of specific BIVA for the assessment of body composition in the United States population. *PLoS One*, 8(3), e58533. doi:10.1371/journal.pone.0058533
- Buono, M. J., Burke, S., Endemann, S., Graham, H., Gressard, C., Griswold, L., & Michalewicz, B. (2004). The effect of ambient air temperature on whole-body bioelectrical impedance. *Physiological Measurement*, 25(1), 119-123.
- Butte, N. F., Hopkinson, J. M., Wong, W. W., Smith, E. O. B., & Ellis, K. J. (2000). Body composition during the first 2 years of life: an updated reference. *Pediatric Research*, 47(5), 578-585.
- Cain, K. L., Sallis, J. F., Conway, T. L., Van Dyck, D., & Calhoun, L. (2013). Using accelerometers in youth physical activity studies: a review of methods. *Journal of Physical Activity and Health*, 10(3), 437-450.
- Cífková, R., Skodová, Z., Bruthans, J., Adámková, V., Jozífová, M., Galovcová, M., . . . Lánská, V. (2010). Longitudinal trends in major cardiovascular risk factors in the Czech population between 1985 and 2007/8. Czech MONICA and Czech post-MONICA. *Atherosclerosis*, 211(2), 676-681. doi:10.1016/j.atherosclerosis.2010.04.007
- Clarys, J. P., Martin, A. D., Marfell-Jones, M. J., Janssens, V., Caboor, D., & Drinkwater, D. T. (1999). Human body composition: a review of adult dissection data. *American Journal of Human Biology*, 11(2), 167-174. doi:10.1002/(SICI)1520-6300(1999)11:2<167::AID-AJHB4>3.0.CO;2-G
- Cleland, V. J., Schmidt, M. D., Dwyer, T., & Venn, A. J. (2008). Television viewing and abdominal obesity in young adults: is the association mediated by food and beverage consumption during viewing time or reduced leisure-time physical activity? *American Journal of Clinical Nutrition*, 87(5), 1148-1155.
- Cliff, D. P., Hesketh, K. D., Vella, S. A., Hinkley, T., Tsiros, M. D., Ridgers, N. D., . . . Lubans, D. R. (2016). Objectively measured sedentary behaviour and health and development in children and adolescents: systematic review and meta-analysis. *Obesity Reviews*, 17(4), 330-344. doi:10.1111/obr.12371

- Codognotto, M., Piazza, M., Frigatti, P., & Piccoli, A. (2008). Influence of localized edema on whole-body and segmental bioelectrical impedance. *Nutrition*, *24*(6), 569-574. doi:10.1016/j.nut.2008.02.009
- Cohen, D. A., Ghosh-Dastidar, B., Conway, T. L., Evenson, K. R., Rodriguez, D. A., Beckman, R., . . . Lytle, L. (2014). Energy balance in adolescent girls: the trial of activity for adolescent girls cohort. *Obesity*, *22*(3), 772-780. doi:10.1002/oby.20536
- Cohn, S. H., & Dombrowski, C. S. (1971). Measurement of total-body calcium, sodium chlorine, nitrogen, and phosphorus in man by *in vivo* neutron activation analysis. *Journal of Nuclear Medicine*, *12*(7), 499-505.
- Coin, A., Sergi, G., Minicuci, N., Giannini, S., Barbiero, E., Manzato, E., . . . Enzi, G. (2008). Fat-free mass and fat mass reference values by dual-energy X-ray absorptiometry (DEXA) in a 20-80 year-old Italian population. *Clinical Nutrition*, *27*(1), 87-94. doi:10.1016/j.clnu.2007.10.008
- Cole, K. S. (1932). Electric phase angle of cell membranes. *Journal of General Physiology*, *15*(6), 641-649.
- Cole, T. J. (1986). Weight/height^p compared to weight/height² for assessing adiposity in childhood: influence of age and bone age on p during puberty. *Annals of Human Biology*, *13*(5), 433-451.
- Cole, T. J., Bellizzi, M. C., Flegal, K. M., & Dietz, W. H. (2000). Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*, *320*(7244), 1240-1243. doi:10.1136/bmj.320.7244.1240
- Cole, T. J., Flegal, K. M., Nicholls, D., & Jackson, A. A. (2007). Body mass index cut offs to define thinness in children and adolescents: international survey. *BMJ*, *335*(7612), 194. doi:10.1136/bmj.39238.399444.55
- Cole, T. J., & Lobstein, T. (2012). Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatric Obesity*, *7*(4), 284-294. doi:10.1111/j.2047-6310.2012.00064.x
- Collings, P. J., Brage, S., Ridgway, C. L., Harvey, N. C., Godfrey, K. M., Inskip, H. M., . . . Ekelund, U. (2013). Physical activity intensity, sedentary time, and body composition in preschoolers. *American Journal of Clinical Nutrition*, *97*(5), 1020-1028. doi:10.3945/ajcn.112.045088
- Connor Gorber, S., Tremblay, M., Moher, D., & Gorber, B. (2007). A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obesity Reviews*, *8*(4), 307-326. doi:10.1111/j.1467-789X.2007.00347.x
- Craig, C. L., Cameron, C., & Tudor-Locke, C. (2012). CANPLAY pedometer normative reference data for 21,271 children and 12,956 adolescents. *Medicine and Science in Sports and Exercise*, *45*(1), 123-129. doi:10.1249/MSS.0b013e31826a0f3a
- Cuberek, R., Gába, A., Svoboda, Z., Pelclová, J., Chmelík, F., Lehnert, M., . . . Frömel, K. (2014). *Walking in the lifestyle of elderly women with a sedentary occupation*. Olomouc: Vydavatelství Univerzity Palackého v Olomouci.
- Damilakis, J., Adams, J. E., Guglielmi, G., & Link, T. M. (2010). Radiation exposure in X-ray-based imaging techniques used in osteoporosis. *European Radiology*, *20*(11), 2707-2714. doi:10.1007/s00330-010-1845-0
- Daniels, P., Burns, R. D., Brusseau, T. A., Hall, M. S., Davidson, L., Adams, T. D., & Eisenman, P. (2017). Effect of a randomised 12-week resistance training programme on muscular strength, cross-sectional area and muscle quality in women having undergone Roux-en-Y gastric bypass. *Journal of Sports Sciences*, Ahead of Print. doi:10.1080/02640414.2017.1322217
- De Lorenzo, A., Bianchi, A., Maroni, P., Iannarelli, A., Di Daniele, N., Iacopino, L., & Di Renzo, L. (2013). Adiposity rather than BMI determines metabolic risk. *International Journal of Cardiology*, *166*(1), 111-117. doi:10.1016/j.ijcard.2011.10.006
- De Lorenzo, A., Deurenberg, P., Pietrantuono, M., Di Daniele, N., Cervelli, V., & Andreoli, A. (2003). How fat is obese? *Acta Diabetologica*, *40*(1), 254-257. doi:10.1007/s00592-003-0079-x
- de Onis, M., & Lobstein, T. (2010). Defining obesity risk status in the general childhood population: Which cut-offs should we use? *International Journal of Pediatric Obesity*, *5*(6), 458-460. doi:10.3109/17477161003615583

- de Onis, M., Onyango, A. W., Borghi, E., Siyam, A., Nishida, C., & Siekmann, J. (2007). Development of a WHO growth reference for school-aged children and adolescents. *Bulletin of the World Health Organization*, 85(9), 660-667.
- De Palo, T., Messina, G., Edefonti, A., Perfumo, F., Pisanello, L., Peruzzi, L., . . . Piccoli, A. (2000). Normal values of the bioelectrical impedance vector in childhood and puberty. *Nutrition*, 16(6), 417-424. doi:10.1016/S0899-9007(00)00269-0
- Dehghan, M., & Merchant, A. T. (2008). Is bioelectrical impedance accurate for use in large epidemiological studies? *Nutrition Journal*, 7(1), 26. doi:10.1186/1475-2891-7-26
- Dempster, P., & Aitkens, S. (1995). A new air displacement method for the determination of human body composition. *Medicine and Science in Sports and Exercise*, 27(12), 1692-1697.
- Demura, S., Sato, S., & Kitabayashi, T. (2004). Percentage of total body fat as estimated by three automatic bioelectrical impedance analyzers. *Journal of Physiological Anthropology and Applied Human Science*, 23(3), 93-99. doi:10.2114/jpa.23.93
- Demura, S., Yamaji, S., Goshi, F., & Nagasawa, Y. (2002). The influence of transient change of total body water on relative body fats based on three bioelectrical impedance analyses methods. Comparison between before and after exercise with sweat loss, and after drinking. *Journal of Sports Medicine and Physical Fitness*, 42(1), 38-44.
- den Hoed, M., & Westerterp, K. R. (2008). Body composition is associated with physical activity in daily life as measured using a triaxial accelerometer in both men and women. *International Journal of Obesity*, 32(8), 1264-1270. doi:10.1038/ijo.2008.72
- Dencker, M., Thorsson, O., Karlsson, M. K., Lindén, C., Eiberg, S., Wollmer, P., & Andersen, L. B. (2006). Daily physical activity related to body fat in children aged 8-11 years. *The Journal of Pediatrics*, 149(1), 38-42. doi:10.1016/j.jpeds.2006.02.002
- Department of Health and Ageing. (2014a). Make your Move – Sit less – Be active for life! (Adults) [brochure]. Retrieved from [http://www.health.gov.au/internet/main/publishing.nsf/content/F01F92328EDADA5BCA257BF0001E720D/\\$File/brochure%20PA%20Guidelines_A5_18-64yrs.PDF](http://www.health.gov.au/internet/main/publishing.nsf/content/F01F92328EDADA5BCA257BF0001E720D/$File/brochure%20PA%20Guidelines_A5_18-64yrs.PDF)
- Department of Health and Ageing. (2014b). Make your move – Sit less – Be active for life! (Children) [brochure]. Retrieved from [http://www.health.gov.au/internet/main/publishing.nsf/content/F01F92328EDADA5BCA257BF0001E720D/\\$File/brochure%20PA%20Guidelines_A5_5-12yrs.PDF](http://www.health.gov.au/internet/main/publishing.nsf/content/F01F92328EDADA5BCA257BF0001E720D/$File/brochure%20PA%20Guidelines_A5_5-12yrs.PDF)
- Department of Health and Ageing. (2014c). Make your move – Sit less – Be active for life! (Young People) [brochure]. Retrieved from [http://www.health.gov.au/internet/main/publishing.nsf/content/F01F92328EDADA5BCA257BF0001E720D/\\$File/brochure%20PA%20Guidelines_A5_13-17yrs.PDF](http://www.health.gov.au/internet/main/publishing.nsf/content/F01F92328EDADA5BCA257BF0001E720D/$File/brochure%20PA%20Guidelines_A5_13-17yrs.PDF)
- Deurenberg, P., & Schutz, Y. (1995). Body composition: overview of methods and future directions of research. *Annals of Nutrition and Metabolism*, 39(6), 325-333. doi:10.1159/000177881
- Deurenberg, P., Weststrate, J. A., Paymans, I., & van der Kooy, K. (1988). Factors affecting bioelectrical impedance measurements in humans. *European Journal of Clinical Nutrition*, 42(12), 1017-1022.
- Deurenberg, P., Weststrate, J. A., & Seidell, J. C. (1991). Body mass index as a measure of body fatness: age-and sex-specific prediction formulas. *British Journal of Nutrition*, 65(2), 105-114.
- Dey, D. K., Bosaeus, I., Lissner, L., & Steen, B. (2009). Changes in body composition and its relation to muscle strength in 75-year-old men and women: A 5-year prospective follow-up study of the NORA cohort in Göteborg, Sweden. *Nutrition*, 25(6), 613-619. doi:10.1016/j.nut.2008.11.023
- Dixon, C. B., Ramos, L., Fitzgerald, E., Reppert, D., & Andreacci, J. L. (2009). The effect of acute fluid consumption on measures of impedance and percent body fat estimated using segmental bioelectrical impedance analysis. *European Journal of Clinical Nutrition*, 63(9), 1115-1122. doi:10.1038/ejcn.2009.42