

**Literatura:**

1. Marzullo P, Minocci A, Tagliaferri MA, et al. Investigations of thyroid hormones and antibodies in obesity: leptin levels are associated with thyroid autoimmunity independent of bioanthropometric, hormonal, and weight-related determinants. *J Clin Endocrinol Metab.* 2010 Aug; 95(8): 3965-3972.
2. Rosenbaum M, Hirsch J, Murphy E, Leibel RL. Effects of changes in body weight on carbohydrate metabolism, catecholamine excretion, and thyroid function. *Am J Clin Nutr.* 2000 Jun; 71(6): 1421-1432.
3. Tiller D, Ittermann T, Greiser KH, et al.. Association of Serum Thyrotropin with Anthropometric Markers of Obesity in the General Population. *Thyroid.* 2016 Sep; 26(9): 1205-1214.
4. Fontenelle LC, Feitosa MM, Severo JS, et al. Thyroid Function in Human Obesity: Underlying Mechanisms. *Horm Metab Res.* 2016 Dec; 48(12): 787-794.
5. Knudsen N Laurberg P, Rasmussen LB, et al. Small differences in thyroid function may be important for body mass index and the occurrence of obesity in the population. *J Clin Endocrinol Metab.* 2005 Jul; 90(7): 4019-4024.
6. Abdi H, Kazemian E, Gharibzadeh S, et al. Association between Thyroid Function and Body Mass Index: A 10-Year Follow-Up. *Ann Nutr Metab.* 2017; 70(4): 338-345.
7. Sjouke B, Langslet G, Ceska R, et al. Eprotirome in patients with familial hypercholesterolaemia (the AKKA trial): a randomised, double-blind, placebo-controlled phase 3 study. *Lancet Diabetes Endocrinol.* 2014 Jun; 2(6): 455-463.
8. Cao Y, Matsubara T, Zhao C, et al. Antisense oligonucleotide and thyroid hormone conjugates for obesity treatment. *Sci Rep.* 2017 Aug 24; 7(1): 9307.
9. American Thyroid Association (ATA) Guidelines Taskforce on Thyroid Nodules and Differentiated Thyroid Cancer: Cooper DS, Doherty GM, Haugen BR, et al. Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid.* 2009 Nov; 19(11): 1167-1214.
10. Miyauchi A, Ito Y. Conservative Surveillance Management of Low-Risk Papillary Thyroid Microcarcinoma. *Endocrinol Metab Clin North Am.* 2019 Mar; 48(1): 215-226.
11. Smallridge RC, Meek SE, Morgan MA, et al. Monitoring thyroglobulin in a sensitive immunoassay has comparable sensitivity to recombinant human tsh-stimulated thyroglobulin in follow-up of thyroid cancer patients. *J Clin Endocrinol Metab.* 2007 Jan; 92(1): 82-87.
12. Lezaic L, Rep S, Sever MJ, et al. <sup>18</sup>F-Fluorocholine PET/CT for localization of hyperfunctioning parathyroid tissue in primary hyperparathyroidism: a pilot study. *Eur J Nucl Med Mol Imaging.* 2014 Nov; 41(11): 2083-2089.
13. Kluijfhout WP, Vorselaars WM, van den Berk SA, et al. Fluorine-18 fluorocholine PET-CT localizes hyperparathyroidism in patients with inconclusive conventional imaging: a multicenter study from the Netherlands. *Nucl Med Commun.* 2016 Dec; 37(12): 1246-1252.