

## 10 REFERENČNÍ SEZNAM

Anker, L. C., Weerdesteyn, V., Nes, I. J. W, Nienhuis, B., Straatman, H. & Guerts, A. C. (2008). The relation between postural stability and weight distribution in healthy subjects. *Gait & Posture*, 27, 471-477.

Anonymous (2003). *The Rehabilitation of People with Amputations*. USA: World Health Organization.

Anonymous (2008). Činnost oboru diabetologie, péče o diabetiky v roce 2008. *UZIS*. Retrieved 10. 7. 2008 from World Wide Web: Retrieved 17. 7. 2008 from The Cochrane Database on the World Wide Web: [http://www.uzis.cz/news.php?mnu\\_id=1100](http://www.uzis.cz/news.php?mnu_id=1100).

Aruin, A. S., Nicholas, J. J., & Latash, M. L. (1997). Anticipatory postural adjustments during standing in below the knee amputees. *Clinical Biomechanics*, 12(1), 52-59.

Arya, A. P., Lees, A., Nirula, H. C., & Klenerman L. (1995). A biomechanical comparison of SACH, Seattle and Jaipur feet using ground reaction forces. *Prosthetic and Orthotic International*, 19(1), 37-45.

Baum, B. S., Schnall, B. L., Tis, J. E., & Lipton, J. S. (2008). Correlation of residual limb length and gait parameters in amputees. *Injury*, 39(7), 728-733.

Beekman, C. E. & Axtell, L. A. (1987). Prosthetic use in elderly patients with dysvascular above-knee and through-knee amputation. *Physical Therapy*, 67, 510-516.

Berencsi, A., Ishihara, M. & Imanaka, K. (2005). The functional role of central and peripheral vision in the control of posture. *Human Movement Science*, 24 (5-6), 689-709.

Berke, G. M. (2007). Transtibial Protheses. In M. M. Lusardi & C. C. Nielsen (Eds). *Orthotics and Prosthetics in Rehabilitation (2nd ed.)* (pp 593-641). St. Louis: Saunders.

Birgusová, G. (2006). Standard fyzioterapie doporučený UNIFY ČR – *Amputace dolní končetiny*. Praha: UNIFY ČR.

Birgusová, G. (2007). Včasná pooperační péče a příprava pacientů k aplikaci protézy. *Sborník semináře Komplexní péče o amputované na dolní končetině*, Ostrava: Ostravská univerzita v Ostravě.

Blaszczyk, J., K. Hansen, P. D., & Lowe, D. L. (1993). Postural sway and perception of the upright stance stability borders. *Perception*, 22 (11), 1333-1341.

Blaszczyk, J. W., Lowe, D. L., & Hansen, P. D. (1994). Ranges of postural stability and their changes in the elderly. *Gait & Posture*, 2, 11-17.

Blaszczyk, J. W., Prince, F., Raiche, M., & Hebert, R. (2000). Effect of ageing and vision on limb load asymmetry during quiet stance. *Journal of Biomechanics*, 33, 1243-1248.

Blaszczyk, J. W., & Klonovski, W. (2001). Postural Stability and Fractal Dynamics. *Acta Neurobiologiae Experimentalis*. 61(6), 105-112.

Bowker, J. H. (2004). Transfemoral amputation surgical management. In D. G. Smith, J. W. Michael & J. H. Bowker (Eds.), *Atlas of Amputations and Limb Deficiencies. Surgical, Prosthetic, and Rehabilitation Principles (3rd ed.)* (pp. 589-619). Rosemont, IL.

Branemark, R., Rydevik, B., & Myers, R. R. (2001). Osseointegration in skeletal reconstruction and rehabilitation: a review. *Journal of Research and Development*, 38(2), 175-181.

Broomhead, P., Dales, D., Hale C., Lambert, A., & Shepherd, R. (2003). *Evidence base clinical guidelines for the physiotherapy management of adults with lower limb prostheses*. London: British Association of Chartered Physiotherapists in Amputation Rehabilitation.

Brozmanová, B. (1990). *Ortopedická protetika*. Martin: Osveta.

Buckley, J. G, O'Driscoll, D., & Bennett, S. J. (2002). Postural sway and active balance performance in highly active lower-limb amputees. *American Journal of Physical Medicine & Rehabilitation*, 81(1), 13-20.

Burke, M. J., Roman, V., & Wright, V. (1978). Bone and joint changes in lower limb amputees. *Annals of the Rheumatic Diseases*, 37(3), 252-254.

Carroll, K., & Binder, K. (2006). Transtibial Prosthetic Designs. In K. Carroll, & J. E. Edelstein, (Eds.). *Prosthetic and Patient Management: a Comprehensive Clinical Approach*. Thorofare: SLACK.

Carroll, K., & Edelstein, J. E. (2006). Clinic team approach to rehabilitation. In K. Carroll, & J. E. Edelstein, (Eds.). *Prosthetic and Patient Management: a Comprehensive Clinical Approach*. Thorofare (USA): SLACK.

Casillas, J., Dulieu, V., Cohen, M., & Didier, J. P. (1995). Bioenergetic comparison of a new energy-restoring foot and SACH-foot in traumatic below-

knee vascular amputations. *Archives of Physical Medicine and Rehabilitation*, 76, 39-44.

Cornilleau-Pérès, V., Shabana, N., Droulez, J., Goh, J. C., Lee, G. S., & Cheb, P. T. (2005). Measurement of the visual contribution to postural steadiness from the COP movement: methodology and reliability. *Gait & Posture*, 22 (2), 96-106.

Day, B. L., Steiger, M. J., Thompson, P. D., & Marsden, C.D. (1993). Effects of vision and stance width on human body motion when standing: implications for afferent control of lateral sway. *Journal of Physiology*, 469, 479-499.

Dietz, V. (1992). Human neuronal control of automatic functional movements: interaction between central programs and afferent input. *Physiological Reviews*, 72, 33-69.

Dolan, N. C., Liu, K., Criqui, M. H., Greebland, P., Guralnik, J. M., & Chan, Ch. et al. (2002). Peripheral artery disease, diabetes and reduced lower extremity functioning. *Diabetes Care*, 25, 113-120.

Dornan, J., Fernie, G. R., & Holliday, R. J. (1978). Visual input: its importance in the control of postural sway. *Archives of Physical Medicine and Rehabilitation*, 59, 586-591.

Duclos, C., Roll, R., Kavounoudias, A., Mongeau, J. P., Roll, J. P., & Forget, R. (2009). Postural changes after sustained neck muscle contraction in persons with a lower leg amputation. *Journal of Electromyography and Kinesiology*, 19, e214-e222.

Dungl, P et al. (2005). *Ortopedie*. Praha: Grada Publishing.

Ehde, D. M., & Smith, D. G. (2004). Chronic pain Management. In D. G. Smith, J. W. Michael & J. H. Bowker (Eds.), *Atlas of Amputations and Limb Deficiencies. Surgical, Prosthetic, and Rehabilitation Principles (3rd ed.)* (pp. 711-726). Rosemont, IL.

Elbert, T., Sterr, A., Flor, H., Rockstroh, B., Knecht, S., & Pantev, C. et al. (1997). Input-increase and input-decrease types of cortical reorganization after upper extremity amputation in humans. *Experimental Brain Research*, 117, 161-164.

Engsberg, J. R., Lee, A. G., Tedford, K. G., & Harder, J. A. (1993). Normative ground reaction force data for able-bodied and below-knee amputee children during walking. *Journal of pediatric orthopedics*, 13(2), 169-173.

Engstrom, B. & Van de Ven, C. (Eds.). (1999). *Therapy for amputees (3th ed.)*. London: Churchill Livingstone.

Falstie-Jensen, N., Christensen, K. S., & Brochner-Mortensen, J. (1989). Long Posterior Flap Versus Equal sagittal Flaps in Below-Knee amputation for Ischaemia. *Journal of Bone and Joint Surgery*, 71(1), 102-104.

Fergason, J. (2007). Prosthetic feet. In M. M. Lusardi & C. C. Nielsen (Eds). *Orthotics and prosthetics in Rehabilitation (2nd ed.)* (pp. 593-641). St. Louis: Saunders.

Fernie, G. R., & Holliday, P. J. (1978). Postural sway in amputees and normal subjects. *Journal of Bone and Joint Surgery*. 60, 895-898.

Fitzpatrick, R., Rogers, D. K., & McCloskey, D. I. (1994). Stable human standing with lower limb muscle afferents providing the only sensory input. *Journal of Physiology*, 480, 395-403.

Flor, H., Elbert, T., Knechti, S., Wienbruch, C., Pantevi, C., & Birbaumer, N. et al. (1995). Phantom limb pain as a perceptual correlate of cortical reorganization following arm amputation. *Nature*, 375, 482-484.

Gagnon-Gauthier, Ch., Grise, M. C., & Potvin, D. (1998). Predisposing Factors Related to Prosthetic Use by People with a Transtibial and Transfemoral Amputation. *Journal of Prosthetic and Orthotics*, 10, 99-109.

Gailey, R. S. (1994). *One Step Ahead – An Integrated Approach to Lower Extremity Prosthetics and Amputee Rehabilitation*, Miami: Advanced rehabilitation Therapy, Inc.

Gailey, R. S., & Clark, R. S. (2004). Physical Therapy. In D. G. Smith, J. W. Michael & J. H. Bowker (Eds.). *Atlas of Amputations and Limb Deficiencies Surgical, Prosthetic, and Rehabilitation Principles, (3rd ed.)* (pp. 589-619). Rosemont: American Academy of Orthopaedic Surgeons.

Gailey, R. S. (2007). „How do you do“ – *Amputee Mobility and Prosthetic Training Workshop 2007*, Miami: Advanced Rehabilitation Therapy, Inc.

Gauthier-Gagnon, C., St-Pierre, D., & Drouin G. et al. (1986). Augmental sensory feedback in the early training of standing balance of below knee amputees. *Physiotherapy Can*, 38, 137-142.

Genthon, N. & Rougier, P. (2005). Influence of an asymmetrical body weight distribution on the control of undisturbed upright stance. *Journal of Biomechanics*, 38, 2037-2049.

Giacomini, P. G., Bruno, E., Monticone, G., Girolamo, S. Di., Magrini, A., & Parisi, L. et al. (1996). Postural rearrangement in IDDM patients with peripheral neuropathy. *Diabetes Care*, 19, 372-374.

Gillespie, J. (2007). Konstrukce lůžka TF protézy. *Sborník semináře Komfort pahýlového lůžka.*, Frýdek Místek: Ossur Academy East Europe.

Gottschalk, F. (2004). Transfemoral Amputation: Surgical Management. In D. G. Smith, J. W. Michael & J. H. Bowker (Eds.). *Atlas of Amputations and Limb Deficiencies Surgical, Prosthetic, and Rehabilitation Principles, (3rd ed.) (pp. 533-540)*. Rosemont: AAOS.

Group, T. G. (2000). Epidemiology of lower extremity amputation in centres in Europe, North America and East Asia. The global lower extremity amputation group. *British Journal of Surgery*, 87(3), 328-337.

Guerts, A. C., Mulder, T. W., & Nienhuis, B. et al. (1991). Dual-task assessment of reorganization of postural control in persons with lower limb amputation. *Archives of Physical Medicine and Rehabilitation*, 72 (13), 1059-64.

Guerts, A. C., Mulder, T. W., & Nienhuis, B. et al. (1992). Postural reorganization following lower limb amputation. Possible motor and sensory determinants of recovery. *Scandinavian Journal of Rehabilitation Medicine*, 24, 83-90.

Hagberg, K., & Brenemark, R. (2001). Consequences of non-vascular transfemoral amputation: a survey of quality of life, prosthetic use and problems. *Prosthetics and Orthotics International*, 25, 186-194.

Hagberg, K., Haggstrom, E., Uden, M., & Branemark R. (2005). Socket versus bone-anchored transfemoral prostheses: Hip range of motion and sitting comfort. *Prosthetics and Orthotics International*, 29(2), 153-163.

Harker, J. (2006). *Wound healing complications associated with lower limb amputation*. World wide wounds journal. Retrieved 25.5. 2009 from World Wide Web:

<http://www.worldwidewounds.com/2006/september/Harker/Wound-Healing-Complications-Limb-Amputation.html>.

Hermodsson, Y., Ekdahl, C., & Persson, B. M. et al. (1994). Standing balance in trans-tibial amputees following vascular disease or trauma: a comparative study with healthy subjects. *Prosthetic and Orthotic International*, 18, 150-158.

Holden, J. M., & Fernie, G. R. (1987). Extent of artificial limb use following rehabilitation. *Journal of Orthopaedic Research*, 5, 562-568.

Horak, F. B., Shupert, C. L. & Mirka, A. (1989). Components of postural sway dyscontrol in the elderly: a review. *Neurobiology of Aging*, 10, 727-738.

Horak, F. B. (1997). Clinical assessment of balance disorders. *Gait & Posture*, 6, 76-84.

Horak, F. B. (2006). Postural orientation and equilibrium: what do we need to know about neural control of balance to prevent falls? *Age and Ageing*, 35(2), ii7 - ii11.

Huang, G. F., Chou, Y. L., & Su, F. C. (2000). Gait analysis and energy consumption of below-knee amputees wearing free different prosthetic feet. *Gait & Posture*, 12(2), 162-168.

Hurwitz, D. E., Sumner, D. R., & Block, J. A. (2001). Bone density, dynamic joint loading and joint degeneration. *Cells Tissues Organs*, 169(3), 201-209.

Chen, R., Cohen, L. G., & Hallett, M. (2002). Nervous system reorganization following injury. *Neuroscience*, 111, 761-773.

Isakov E., Mizrahi, J., & Ring, H. et al. (1992). Standing sway and weight-bearing distribution in people with below knee amputations. *Archives of Physical and Medicine and Rehabilitation*, 73(2), 174-178.

Isakov, E., Burdoragin, N., & Shenhav, S. et al. (1995). Anatomic sites of foot lesions resulting in amputation among diabetics and non-diabetics. *American Journal of Physical Medicine and Rehabilitation*, 74(2), 130-133.

Isakov, E., Keren, O., & Benjuya, N. (2000). Trans-tibial amputee gait: time - distance parameters and EMG activity. *Prosthetic and Orthotic International*, 24(3), 216-220.

Jaegers, S., Vos, L., & Rispens, P. (1993). The relationship between comfortable and most metabolically efficient walking speed in persons with unilateral above-knee amputation. *Archives of Physical Medicine and Rehabilitation*, 75, 521-525.