

## REFERENČNÍ SEZNAM

- Ambler, Z. (2002). *Neurologie pro studenty lékařské fakulty*. Praha: Universita Karlova v Praze, Karolinum.
- Ambler, Z. (2006). *Základy neurologie*. Praha: Galén.
- Ambler, Z., Bednařík, J., Růžička, E. et al. (2008). *Klinická neurologie, část obecná*. 2. vydání, Praha: Triton.
- Blennerhassett, J. M., Carey, L. M., & Matyas, T. A. (2006). Grip force regulation during pinch grip lifts under somatosensory guidance: comparison between people with stroke and healthy controls. *Archives of Physical Medicine and Rehabilitation*, 87, 418-429.
- Blennerhassett, J. M., Matyas, T. A., & Carey, L. M. (2007). Impaired discrimination of surface friction contributes to pinch grip deficit after stroke. *Neurorehabilitation and Neural Repair*, 21, 263-272.
- Blennerhassett, J. M., Carey, L. M., & Matyas, T. A. (2008). Clinical measures of handgrip limitation relate to impaired pinch grip force control after stroke. *Journal of Hand Therapy*, 21, 245-253.
- Bobath, B. (1997). *Hemiplégia dospělých*. Bratislava: Liečreh Gúth.
- Busse, M., & Tyson, S. F. (2009). How many body locations need to be tested when assessing sensation after stroke? An investigation of redundancy in the Rivermead Assessment of Somatosensory Performance. *Clinical Rehabilitation*, 23, 91-95.
- Byl, N., Roderick, J., Mohamed, O., Hanny, M., Kotler, J., Smith, A., Tang, M., & Abrams, G. (2003). Effectiveness of Sensory and Motor Rehabilitation of the Upper Limb Following the Principles of Neuroplasticity: Patients Stable Poststroke. *Neurorehabilitation and Neural Repair*, 17 (3), 176-191.

- Carr, J. H., & Shepherd, R. B. (1998). *Neurological Rehabilitation: Optimizing Motor Performance*. Oxford: Butterworth – Heinemann.
- Carey, L. M., Matyas, T. A., & Oke, L. E. (1993). Sensory loss in stroke patients: effective training of tactile and proprioceptive discrimination. *Archives of Physical Medicine and Rehabilitation*, 74, 602-611.
- Carey, L. M. (1995). Somatosensory loss after stroke. *Critical Reviews in Physical and Rehabilitation Medicine*, 7, 51-91.
- Carey, L. M., Oke, L. E., & Matyas, T. A. (1996). Impaired limb position sense after stroke: a quantitative test for clinical use. *Archives of Physical Medicine and Rehabilitation*, 77, 1271-1278.
- Carey, L. M., Oke, L. E., & Matyas, T. A. (1997). Impaired touch discrimination after stroke: a quantitative test. *Journal of Neurologic Rehabilitation*, 11, 219-232.
- Carey, L. M., Matyas, T. A., & Oke, L. E. (2002). Evaluation of impaired fingertip texture discrimination and wrist position sense in patients affected by stroke: comparison of clinical and new quantitative measures. *Journal of Hand Therapy*, 15, 71-82.
- Carey, L. M., & Matyas, T. A. (2005). Training of somatosensory discrimination after stroke: Facilitation of stimulus generalization. *Archives of Physical Medicine and Rehabilitation*, 88, 428-442.
- Carey, L. M., Abbott, D. F., Egan, G. F., O'Keefe, G. J., Bernhardt, J., & Donnan, G. A. (2005). Motor Impairment and Recovery in the Upper Limb after Stroke. *Stroke*, 36, 625-629.
- Carey, L. M., Abbott, D. F., Egan, G. F., O'Keefe, G. J., Jackson, G. D., Bernhardt, J., & Donnan, G. A. (2006). Evolution of Brain Activation with Good and Poor Motor Recovery after Stroke. *Neurorehabilitation and Neural Repair*, 20, 24-41.
- Carey, L. M., & Matyas, T. A. (2011). Frequency of discriminative sensory loss in the hand after stroke in a rehabilitation setting. *Journal of Rehabilitation Medicine*, 43, 82-88.

- Cambell, J. A., Lahuerta, J., & Bowsher, D. (1996). Quantitative assessment of somatosensory function. *British Journal of Therapy and Rehabilitation*, 3, 135-141.
- Cordo, P., & Harnad, S. R. (1994). *Movement control*. United States: Cambridge University Press.
- Čihák, R. (1999). *Anatomie 3*. Praha: Grada Publishing.
- Dannenbaum, R. M., Michaelsen, S. M., Desrosiers, J., & Levin, M. F. (2002). Development and validation of two new sensory tests of the hand for patients with stroke. *Clinical Rehabilitation*, 16 (6), 630-639.
- Dannenbaum, R. M., & Dykes, R. W. (1990). Evaluating sustained touch-pressure in severe sensory deficits: Meeting an unanswered need. *Archives of Physical Medicine and Rehabilitation*, 71, 455-459.
- Davies, P. M. (2000). *Steps to Follow*. 2nd. ed. Berlin, Heidelberg: Springer – Verlag.
- Duncan, P. W., Werner, D. K., Chander, J., & Studenski, S. (1990). Functional Reach: A New Clinical Measure of Balance. *Journal of Gerontology*, 45(6), 192-197.
- Dylevský, I., Druga, R., & Mrázková, O. (2000). *Funkční anatomie člověka*. Praha: Grada Publishing.
- Feydy, A., Carlier, R., Roby-Brami, A., Bussel, B., Cazalis, F., Pierot, L., Burnod, Y., & Maier, M. A. (2002). Longitudinal Study of Motor Recovery after Stroke: Recruitment and Focusing of Brain Activation. *Stroke*, 33, 1610-1617.
- Ganong, W. F. (2005). *Přehled lékařské fyziologie*. Praha: Galén.
- Goldstone, R. L. (1998). Perceptual Learning. *Annual Review of Psychology*, 49, 585-612.
- Howle, J. M. (2002). *Neuro-Developmental Treatment Approach. Theoretical Foundations and Principles of Clinical Practice*. Canada, Laguna Beach: Neuro-Developmental Treatment Association.