
10 REFERENČNÍ SEZNAM

- Alton, F., Baldey, L., Caplan, S., & Morrissey, M. C. (1998). A kinematic comparison of overground and treadmill walking. *Clinical Biomechanics*, *13*(6), 434-440.
- Anderson, F. C., & Pandy, M. G. (2003). Individual muscle contributions to support in normal walking. *Gait & Posture*, *17*(2), 159-169. doi: 10.1016/S0966-6362(02)00073-5
- Annaswamy, T. M., Giddings, C. J., Della Croce, U., & Kerrigan, C. (1999). Rectus femoris: Its role in normal gait. *Archives of Physical Medicine and Rehabilitation*, *80*(8), 930-934. doi: 10.1016/S0003-9993(99)90085-0
- Arendt-nielsen, L., Sinkjaer, T., Nielsen, J., & Kallesoe, K. (1991). Electromyographic patterns and knee joint kinematics during walking at various speeds. *Journal of Electromyography and Kinesiology*, *1*(2), 89-95. doi: 10.1016/1050-6411(91)90002-m
- Basmajian, J. V., & De Luca, C. J. (1985). *Muscles alive: Their functions revealed by electromyography*. Baltimore: Williams & Wilkins.
- Batlkhani, B., Oyunaa, C., & Odongua, N. (2014). A kinematic comparison of overground and treadmill walking. *Value Health*, *17*(7), A774. doi: 10.1016/j.jval.2014.08.337
- BioSpace. (2008). *InBody 720. Precision in body composition analysis. User's Manual*. Retrieved from http://inbody.com/_admin/uploads/GLOBAL/InBody720Nawi_CDmanual_Eng_A.pdf.
- Bobbert, A. C. (1960). Energy expenditure in level and grade walking. *Journal of Applied Physiology*, *15*(6), 1015-1021. doi: 10.1152/jappl.1960.15.6.1015
- Bompa, T. O., & Buzzichelli, C. (2015). *Periodization training for sports* (3rd ed.). Champaign, IL: Human Kinetics.
- Booth, F. W., Roberts, C. K., & Laye, M. J. (2012). Lack of exercise is a major cause of chronic diseases. *Comprehensive Physiology*, *2*(2), 1143-1211. doi: 10.1002/cphy.c110025

- Borg, G. (1990). Psychophysical scaling with applications in physical work and the perception of exertion. *Scandinavian Journal of Work, Environment & Health*, 16, 55-58. doi: 10.5271/sjweh.1815
- Bortolan, L., Pellegrini, B., & Schena, F. (2009). *Development and validation of a system for poling force measurement in cross-country skiing and nordic walking*. Paper presented at the 27th International Conference on Biomechanics in Sports, Rovereto, IT.
- Bronstein, A. M., Brandt, T., & Woollacott, M. H. (1996). *Clinical disorders of balance, posture and gait*. London: Arnold.
- Bruijn, S. M., Meijer, O. G., Van Dieen, J. H., Kingma, I., & Larnoth, C. J. C. (2008). Coordination of leg swing, thorax rotations, and pelvis rotations during gait: The organisation of total body angular momentum. *Gait & Posture*, 27(3), 455-462. doi: 10.1016/j.gaitpost.2007.05.017
- Burke, E. R. (1998). *Precision heart rate training*. Champaign, IL: Human Kinetics.
- Busse, R., Blümel, M., Scheller-Kreinsen, D., & Zentner, A. (2010). *Tackling chronic disease in Europe: Strategies, interventions and challenges*. Copenhagen: WHO, Regional Office for Europe.
- Byrne, C. A., O'Keeffe, D. T., Donnelly, A. E., & Lyons, G. M. (2007). Effect of walking speed changes on tibialis anterior EMG during healthy gait for FES envelope design in drop foot correction. *Journal of Electromyography and Kinesiology*, 17(5), 605-616. doi: 10.1016/j.jelekin.2006.07.008
- Caballero, B., Allen, L., & Prentice, A. (2005). *Encyclopedia of human nutrition* (2nd ed.). Oxford: Elsevier Academic Press.
- CDC. (2015). Target heart rate and estimated maximum heart rate. Retrieved from <https://www.cdc.gov/physicalactivity/basics/measuring/heartrate.htm>.
- Cibulka, M. T., Winters, K., Kampwerth, T., McAfee, B., Payne, L., Roeckenhaus, T., & Ross, S. A. (2016). Predicting foot progression angle during gait using two clinical measures in healthy adults, a preliminary study. *International Journal of Sports Physical Therapy*, 11(3), 400-408.
- Craik, R. L., & Oatis, C. A. (1995). *Gait analysis: Theory and application*. St. Louis, MO: Mosby.

- Criswell, E. (2011). *Cram's introduction to surface electromyography* (2nd ed.). Sudbury, MA: Jones and Bartlett Publishers.
- Čihák, R. (2011). *Anatomie 1. Třetí, upravené a doplněné vydání*. Praha: Grada.
- De Luca, C. J., & Merletti, R. (1988). Surface myoelectric signal cross-talk among muscles of the leg. *Electroencephalography and Clinical Neurophysiology*, 69(6), 568-575. doi: 10.1016/0013-4694(88)90169-1
- Dedieu, P., & Zanone, P. G. (2012). Effects of gait pattern and arm swing on intergirdle coordination. *Human Movement Science*, 31(3), 660-671. doi: 10.1016/j.humov.2011.07.009
- Dechman, G., Appleby, J., Carr, M., & Haire, M. (2012). Comparison of treadmill and over-ground Nordic walking. *European Journal of Sport Science*, 12(1), 36-42. doi: 10.1080/17461391.2010.551411
- Di Blasio, A., Morano, T., Bucci, I., Di Santo, S., D'Arielli, A., Castro, C. G., . . . Napolitano, G. (2016). Physical exercises for breast cancer survivors: Effects of 10 weeks of training on upper limb circumferences. *Journal of Physical Therapy Science*, 28(10), 2778-2784. doi: 10.1589/jpts.28.2778
- Donker, S. F., Beek, P. J., Wagenaar, R. C., & Mulder, T. (2001). Coordination between arm and leg movements during locomotion. *Journal of Motor Behavior*, 33(1), 86-102. doi: 10.1080/00222890109601905
- Downer, D. (2006). *Nordic walking step by step*. [s.l.]: Nordic Walking Publications.
- Dunġl, P. (2014). *Ortopedie (2. přepracované a doplněné vydání)*. Praha: Grada Publishing, a.s.
- Dvořák, R. (2007). *Základy kinezioterapie*. Olomouc: Univerzita Palackého v Olomouci.
- EC. (2008). EU physical activity guidelines; Recommended policy actions in support of health-enhancing physical activity. Retrieved from http://ec.europa.eu/sport/library/policy_documents/eu-physical-activity-guidelines-2008_en.pdf.
- Egerton, T., Danoudis, M., Huxham, F., & Ianssek, R. (2011). Central gait control mechanisms and the stride length - cadence relationship. *Gait & Posture*, 34(2), 178-182. doi: 10.1016/j.gaitpost.2011.04.006
- Encarnación-Martínez, A., Pérez-Soriano, P., & Llana-Belloch, S. (2015). Differences in Ground Reaction Forces and Shock Impacts Between

- Nordic Walking and Walking. *Research Quarterly for Exercise and Sport*, 86(1), 94-99. doi: 10.1080/02701367.2014.975178
- Etnyre, B. R., & Abraham, L. D. (1988). Antagonist muscle-activity during stretching: A paradox re-assessed. *Medicine and Science in Sports and Exercise*, 20(3), 285-289. doi: 10.1249/00005768-198806000-00013
- Figard-Fabre, H., Fabre, N., Leonardi, A., & Schena, F. (2011). Efficacy of Nordic walking in obesity management. *International Journal of Sports Medicine*, 32(6), 407-414. doi: 10.1055/s-0030-1268461
- Finley, F. R., Cody, K. A., & Finizie, R. V. (1969). Locomotion patterns in elderly women. *Archives of Physical Medicine and Rehabilitation*, 50(3), 140-146.
- FiTRONiC. (n.d.). *FiTRO Light Gates. Bezdrôtová fotobunková časomiera. Návod na použitie*. Retrieved from http://fitronic.sk/images/upload/fitro/NENG/gates/download/manual_Gates_SK.pdf.
- Fleck, S. J., & Kraemer, W. J. (2014). *Designing resistance training programs* (4th ed.). Champaign, IL: Human Kinetics.
- Franz, J. R., & Kram, R. (2012). The effects of grade and speed on leg muscle activations during walking. *Gait & Posture*, 35(1), 143-147. doi: 10.1016/j.gaitpost.2011.08.025
- Gamble, P. (2010). *Strength and conditioning for team sports : Sport-specific physical preparation for high performance*. New York, NY: Routledge.
- Gao, Y., Arfat, Y., Wang, H., & Goswami, N. (2018). Muscle Atrophy Induced by Mechanical Unloading: Mechanisms and Potential Countermeasures. *Frontiers in Physiology*, 9, 235. doi: 10.3389/fphys.2018.00235
- Gerilovsky, L., Tsvetinov, P., & Trenkova, G. (1986). H-reflex potentials shape and amplitude changes at different length of relaxed soleus muscle. *Electromyography and Clinical Neurophysiology*, 26(8), 641-653.
- Gimmon, Y., Riemer, R., Rashed, H., Shapiro, A., Debi, R., Kurz, I., & Melzer, I. (2015). Age-related differences in pelvic and trunk motion and gait adaptability at different walking speeds. *Journal of Electromyography and Kinesiology*, 25(5), 791-799. doi: 10.1016/j.jelekin.2015.05.003
- Gram, B., Christensen, R., Christiansen, C., & Gram, J. (2010). Effects of nordic walking and exercise in type 2 diabetes mellitus: A randomized controlled