

## 10 Referenční seznam

- Anonymous (2019a). Retrieved 13. 1. 2019 from World Wide Web  
<https://docs.vicon.com/display/Nexus25/Calibrate+a+Vicon+system>
- Anonymous (2018b). Retrieved 21. 1. 2019 from World Wide Web  
[blog.strazcenter.org/tag/pierre-beauchamp/](http://blog.strazcenter.org/tag/pierre-beauchamp/)
- Anonymous (2019c). Retrieved 21. 1. 2019 from World Wide Web  
<https://bestpointe.com/category/shoes/pointe-shoes/smart-pointe-by-grishko/>
- Anonymous (2019d). Retrieved 21. 1. 2019 from World Wide Web  
[https://www.urbantrend.cz/produkt/baletni-spicky/?attribute\\_pa\\_color=s-vycpavkou&attribute\\_pa\\_6f6cb72d544962fa333e2e=32&gclid=CjwKCAjwqZPrBRBnEiwAmNJsNt2wbi2KbpmlIWKJcG5aVvZoOAK34CLCilaZJ9LtuQWOJ6v\\_mHJEuRoCDNMQAvD\\_BwE](https://www.urbantrend.cz/produkt/baletni-spicky/?attribute_pa_color=s-vycpavkou&attribute_pa_6f6cb72d544962fa333e2e=32&gclid=CjwKCAjwqZPrBRBnEiwAmNJsNt2wbi2KbpmlIWKJcG5aVvZoOAK34CLCilaZJ9LtuQWOJ6v_mHJEuRoCDNMQAvD_BwE)
- Anonymous (2019e). Retrieved 21. 1. 2019 from World Wide Web  
[https://www.decathlon.cz/platene-baletni-piskoty?id\\_8527698.html?gclid=CjwKCAjwqZPrBRBnEiwAmNJsNnb1dbrA5nEeDh31Y1COwP00o9LcZOitkvPTfZNTv1bx8m5VLPcgzhoC5jsQAvD\\_BwE](https://www.decathlon.cz/platene-baletni-piskoty?id_8527698.html?gclid=CjwKCAjwqZPrBRBnEiwAmNJsNnb1dbrA5nEeDh31Y1COwP00o9LcZOitkvPTfZNTv1bx8m5VLPcgzhoC5jsQAvD_BwE)
- Aalten, A. (2005). In the Presence of the Body: Theorizing Training, Injuries and Pain in Ballet. *Dance Research Journal*, 37(2), 55-72. doi: 10.1017/S0149767700008561
- Adams, J. M., & Perry, J. (2006). Gait analysis: Clinical Decision Making. In Rose, J. & Gamble J. G. (Eds.), *Human Walking* (pp. 165-183). Baltimore: Williams & Wilkins.
- Ahonen, J. (2008). Biomechanics of the foot in dance. A literature review. *Journal of Dance Medicine & Science*, 12(3), 99-108.
- Albert, S. F., & Curran, S. A. (2018). *Lower extremity biomechanics: Theory and practice* (1st ed.). Colorado, USA: Bipedmed.
- Alderson, J., Hopper, L., Elliott, B. & Ackland, T. (2009). Risk factors for lower back injury in male dancers performing ballet lifts. *Journal of Dance Medicine & Science*, 13(3), 83-89.
- Anderson, F. C., Goldberg, S. R., Pandy, M. G., & Scott, L. D. (2004). Contributions of muscle forces and toe-off kinematics to peak flexion during the swing phase of normal gait: an induced position analysis. *Journal of Biomechanics*, 37(5), 731-737.
- Ardestani, M. M., Ferrigno, C., Moazen, M., & Wimmer, M. A. From normal to fast walking: Impact of cadence and stride length on lower extremity joint moments. *Gait & Posture*, 46(May), 118-125. doi: 10.1016/j.gaitpost.2016.02.005
- Ayyappa, E. (1997). Normal human locomotion, part 1: Basic concepts and terminology. *Journal of Prosthetic and Orthotics*, 9(1), 10-17.

- Barwick, A., Smith, J., & Chuter, V. (2012). The relationship between foot motion and lumbopelvic-hip function. A review of the literature. *The Foot*, 22(3), 224-231.
- Batson, G. (2009). Update on proprioception considerations for dance education. *Journal of Dance Medicine & Science*, 13(2), 35-41.
- Bejjani, F. J. (1990). Occupational biomechanics. In Jahss, M. (Eds.), *Disorders of the foot and ankle*. Philadelphia: WB Saunders.
- Bennel, K., Khan, K. M., Matthews, B., de Gruyter, M., Cook, E., Holzer, K., & Wark, J. D. (1999). Hip and ankle range of motion and hip muscle strength in young female ballet dancers and controls. *British Journal of Sports Medicine*, 33(5), 340-346.
- Bennel, K., Khan, K. M., Matthews, B., & Singleton, C. (2001). Changes in hip and ankle range of motion and hip muscle strength in 8-11 year old novice female ballet dancers and controls: a 12 month follow up study. *British Journal of Sports Medicine*, 35(1), 54-59. doi: 10.1136/bjism.35.1.54
- Bickle, C., Deighan, M., & Theis, N. (2018). The effect of pointe shoe deterioration on foot and ankle kinematics and kinetics in professional ballet dancers. *Human Movement Science*, 60(Aug.), 72-77. doi: 10.1016/j.humov.2018.05.011
- Bisi, M. C., & Stagni, R. (2016). Complexity of human gait pattern at different ages assessed using multiscale entropy: From development to decline. *Gait & Posture*, 47(Jun.), 37-42. doi: 10.1016/j.gaitpost.2016.04.001
- Biz, C., Favero, L., Stecco, C., & Aldegheri, R. (2012). Hypermobility of the first ray in ballet dancer. *Muscles, Ligaments and Tendons Journal*, 2(4), 282-288.
- Boling, M. C., Padua, D. A., & Marshall, S. W. (2009). A prospective investigation of biomechanical risk factors for patellofemoral pain syndrome: The joint undertaking to monitor and prevent acl injury. *The American Journal of Sports Medicine*, 24(Sep), 1-7. doi: 10.1177/0363546509337934
- Bowerman, E. A., Whatman, C., Harris, N., & Bradshaw, E. (2015). A Review of the Risk Factors for Lower Extremity Overuse Injuries in Young Elite Female Ballet Dancers. *Journal of Dance Medicine & Science*, 19(2), 51-56. doi: 10.12678/1089-313X.19.2.51
- Bracilović, A. (2009). *Essential dance medicine*. New York: Humana Press.
- Brodská, B. (2007). *Romantický balet*. Praha, Česká republika: Akademie múzických umění.
- Bronner, S. (2012). Differences in segmental coordination and postural control in a multijoint dance movement: développé arabesque. *Journal of Dance Medicine & Science*, 16(1), 26-35.
- Bronner, S., Ojofeitimi, S., Lora, J. B., Southwick, H., Kulak, M. C., Gamboa, J., ... Gibbs, R. (2014). A preseason cardiorespiratory profile of dancers in nine professional ballet and modern companies. *Journal of Dance Medicine & Science*, 18(2), 74-85.

- Bronner, S., Ojofeitimi, S., & Rose, D. (2003). Injuries in a Modern Dance Company. Effect of Comprehensive Management on Injury Incidence and Time Loss. *The American Journal of Sports Medicine*, 31(3), 365-373.
- Buldt, A. K., Murley, G. S., Butterworth, P., Levinger, P., Menz, H. B., & Landorf, K. B. (2013). The relationship between foot posture and lower limb kinematics during walking: A systematic review. *Gait & Posture*, 38(3), 363-372. doi: 10.1016/j.gaitpost.2013.01.010
- Bursová, M. (2005). *Kompenzační cvičení: uvolňovací, protahovací, posilovací*. Praha, Česká republika: Grada.
- Busselová, D. (1995). *Abeceda baletu*. Praha: Ikar.
- Bryan, N., & Smith, B. M. (1992). The ballet dancer. *Occupational Medicine: State of the Art Reviews*, 7(1), 67-75.
- Cachupe, W. C. J., Shifflett, B., Kahanov, L. & Wughalter, E. H. (2001). Reliability of biodex balance system measures. *Physical Education and Exercise Science*, 5(2), 97-108. doi: 10.1207/S15327841MPEE0502\_3
- Carter, S. L., Bryant, A. R., & Hopper, L. S. (2017). Lower leg and foot contributions to turnout in university level female ballet dancers: a preliminary investigation. *Journal of the American Podiatric Medical*, 12(10).doi: 10.7547/15-142
- Celletti, C., Galli, M., Cimolin, V., Castori, M., Tenore, N., Albertini, G., & Camerota, F. (2013). Use of the Gait Profile Score for the evaluation of patients with joint hypermobility syndrome/Ehlers-Danlos syndrome hypermobility type. *Research in Developmental Disabilities*, 34(11), 4280-4285. doi: 10.1016/j.ridd.2013.09.019
- Cimelli, S. N. & Curran, S. A. (2012). Influence of turnout on foot posture and its relationship to overuse musculoskeletal injury in professional contemporary dancers. *Journal of the American Podiatric Medical Association*, 102 (1), 25-33.
- Clippinger, K. (2007). *Dance anatomy and kinesiology*. Champaign, USA: Human Kinetics.
- de Cock, A-M., Fransen, E., Perkisas, Verhoeven, V., Beauchet, O., Remmen, R., & Vandewoude, M. (2017). Gait characteristic under different walking conditions: Association with the Presence of Cognitive impairment in community-dwelling older people. *PLoS ONE*, 12(6), 1-19. doi 10.1371/journal.pone.0178566
- Coplan, J. A. (2002). Ballet dancer's turnout and its relationship to self-reported injury. *Journal of Orthopaedic and Sports Physical Therapy*, 32(11), 579-584. doi: 10.2519/jospt.2002.32.11.579
- Coughlan, G., & Caulfield, B. (2007). A 4-week neuromuscular training program and gait patterns at the ankle joint. *Journal of Athletic Training*, 42(1), 51-59.