

9 REFERENČNÍ SEZNAM

- Abramowitz, M. K., Hall, C. B., Amodu, A., Sharma, D., Androga, L., & Hawkins, M. (2018) Correction: Muscle mass, BMI, and mortality among adults in the United States: A population-based cohort study. *Plos One*, 13(5): e0198318.
- Abu-Omar, K., Rütten, A., & Lehtinen, V. (2004). Mental health and physical activity in the European Union. *Social and Preventive Medicine*, 49(5), 301–309.
- Afari, N., Cuneo, J., G., Herbert, M., Miller, I., Webb-Murphy, J., Delaney, E., ... Wisbach, G. (2019). Design for a cohort-randomized trial of an acceptance and commitment therapy-enhanced weight management and fitness program for Navy personnel. *Contemporary Clinical Trials Communications*, 15.
- Alzeidan, R., Fayed, A., Rabiee, F., Hersi, A., & Elmorshedy, H. (2020). Diagnostic performance of waist-to-height ratio in identifying cardiovascular risk factors and metabolic syndrome among adult Saudis. A cross-sectional study. *Saudi medical journal*, 41(3), 253–260.
- Anděl, J. (1985). *Matematická statistika*. Praha: SNTL/Alfa.
- Anděl, J. (2007). *Základy matematické statistiky*. Druhé opravené vydání. Praha: Matfyzpress.
- Aoyagi, Y., & Shephard, R., J. (2009). Steps per day: The road to senior health? *Sports Medicine*, 39, 423–438.
- Arimbawa, M., Putra, S., Suryawan, B., Windiani, T., & Widyadharma, E. (2019). Association of Body Fat Thickness with Vitamin D Levels in Obese Adolescent. *International Journal of Medical Reviews and Case Reports*, 3(3), 94–98.
- Ashwell, M., & Gibson, S. (2016). Waist-to-height ratio as an indicator of 'early health risk': simpler and more predictive than using a 'matrix' based on BMI and waist circumference. *British Medical Journal Open*, 6(3), e010159.
- Ashwell, M., Gunn, P., & Gibson, S. (2012). Waist-to-height ratio is a better screening tool than waist circumference and BMI for adult cardiometabolic risk factors: systematic review and meta-analysis. *Obesity Reviews*, 13, 275–286.
- Atkinson, G., & Nevill, A. M. (1998). Statistical Methods For Assessing Measurement Error (Reliability) in Variables Relevant to Sports Medicine. *Sports Medicine*, 26(4), 217–238.
- Avila, J. A., Lima, F., Paulo, D., Páscoa, M. A., & Tessutti, L. S. (2013). Efeito de 13 semanas de treinamento físico militar sobre a composição corporal e o desempenho

- físico dos alunos da escola preparatória de cadetes do exército. *Revista Brasileira de Medicina do Esporte*, 19(5), 363–366.
- Ayhan, Savaş & Ayhan, S. (2016). The Effect of Different Sport Branches on Body Fat Percentage. *The Journal of international medical research*, 8, 41–47.
- Bartůňková, S. (2007). *Fyziologie člověka a tělesných cvičení*. UK Praha.
- Bauman, A., Bull, F., Chey, T., Craig, C. L., Ainsworth, B., E., & The IPS GROUP. (2009). *The International Prevalence Study on Physical Activity*. Retrieved 21. 1. 2021 from the World Wide Web: <http://www.ijbnpa.org/content/6/1/21>.
- Benešová, D., Hranáč, P., & Švátora, K. (2018). Vliv ideomotorické intervence v průběhu senzomotorického učení. *Studia Kinanthropologica*, 19(3), 159–163.
- Bentley, D. J., Roels, B., Hellard, P., Faucuet, S., & Millet, G. (2005). Physiological responses during submaximal interval swimming training: Effects of interval duration. *Journal of Science and Medicine in Sport*, 8(4), 392–402.
- Berghofer, A., Pischon, T., Reinhold, T., Apovian, C. M., Sharma, A. M., & Willich, S. N. (2008). Obesity prevalence from a European perspective: a systematic review. *BMC Public Health*, 8(200).
- Bernstein, S. A., Lo, M., & Davis, W. S. (2017). Proposing Using Waist-to-Height Ratio as the Initial Metric for Body Fat Assessment Standards in the U.S. Army. *Military Medicine*, 182(1), 304–309.
- Bertalanffy, L. (1970). General System Theory. In Stefanov, N. et al. *Riadenie, modelovanie, prognózovanie*, 12–13. Bratislava: Pravda.
- Blahuš, P. (1976). *K teorii testování pohybových schopností*. Praha: Karlova universita.
- Blahuš, P. (1996). *K systémovému pojetí statistických metod v metodologii empirického výzkumu chování*. Praha: Karolinum.
- Blasio, A. D., Donato, F. D., & Mazzocco, Ch. (2005). *IPAQ - AUTOMATIC REPORT - English version - self-admin short - Di Blasio et al.* From World Wide Web: <https://docs.google.com/viewer?a=v&pid=sites&srcid=ZGVmYXVsdGRvbWFpbmN0aGVpcGFxfGd4OjUxNjA1MzU5ZDI3NjViNmQ>.
- Blanchonette, P. (2017). *Australian Defence Force Anthropometry: A Summary of Historic Surveys*. Commonwealth of Australia.
- Bompa, T. (2000). *Total training for young champions*. Champaign, IL: Human Kinetics.
- Bouchard, C. (2000). *Physical activity and Obesity*. Champaign: Human Kinetics.
- Bouchard, C., Blair, S. N., & Haskell, W. L. (2007). *Physical activity and health*. (2nd ed.). Champaign, IL: Human Kinetics.

- Bouchard, C., Stepward, D. J., & Stephens, T. (1994). *Physical activity fitness and health: international proceedings and consensus statement*. Champaign IL: Human Kinetics.
- Bouchard, C., & Katzmarzyk, P. T. (2010). *Physical activity and obesity*. Champaign: Human Kinetics.
- Bös, K., & Mechlin, H. (1983). *Dimension sportmotorischer Leistungen*. Schorndorf: Karl Hofmann.
- Brandon, L. J., Milliken, L. A., Blew, R. M., Lohman, T. G. (2020). Body composition field methods. In: Lohman TG, Milliken LA, editors. ACSM's body composition assessment. *Human Kinetics*, 2020, 59–89.
- Brožová, H. (2007). *Modely pro řízení znalostí a podporu rozhodování*. Praha: Česká zemědělská univerzita.
- Bunc, V. (1990). *Biokybernetický přístup k hodnocení reakce organismu na tělesné zatížení*. Praha: VÚT UK.
- Bunc, V., Hráský, P., & Skalská, M. (2012). Pohybové aktivity seniorů – benefity a problémy. In Štěpánková, H. (Ed.), *Stárnutí 2012 (23-29)*. Praha: Psychiatrické centrum.
- Bunc, V., Hráský, P., & Skalská, M. (2014). Body Composition and Aerobic Fitness like a Result of 6Month Walking program in Senior Women. *Journal of Medical Research*, 2014, 10.
- Bunc, V., & Štilec, M. (2003). Possibilities of body composition and aerobic fitness influence by walking in senior women. In K. Einfeld, U. Wiesmann, H. J. Hannich & P. Hirtz (Eds.), *Gesund und bewegt ins Alter*, 193–200.
- Bunc, V., & Štilec, M. (2007). Tělesné složení jako indikátor aktivního životního stylu seniorek. *Česká kinantropologie*, 11(3), 17–23.
- Bunc, V., Štilec, M., Moravcová, J., & Matouš, M. (2000). Body composition determinativ by whole body bioimpedance measurement in women seniors. *Acta Universitatis Carolinae Kinanthropology*, 36(1), 23–38.
- Bunc, V., Štilec, M., Moravcová, J., & Matouš, M. (2010). Body composition determination by whole body bioimpedance measurement in women seniors. *Acta University Karolinae Kinanthropologica*, 110, 113–140.