

Literatura.

1. Petráš P, Ježek P, Machová I, Bílina P, Fiala J. Další smrtelný případ pneumonie vyvolané kmenem *Staphylococcus aureus* s produkcí Pantonova – Valentinova leukocidinu. Zprávy EM 2009; 18: 15-17.
2. Vandenesch F, Naimi T, Enright MC *et al.* Community-acquired methicillin-resistant *Staphylococcus aureus* carrying Panton-Valentine leukocidin genes: worldwide emergence. *Emerg Inf Dis* 2003; 9: 978-984.
3. Okuma K, Iwakawa K, Runidge JD *et al.* Dissemination of new methicillin-resistant *Staphylococcus aureus* clones in the community. *J Clin Microbiol* 2002; 40: 4289-4294.
4. Fusco V, Quero GM, Morea M, Blaiotta G, Visconti A. Rapid and reliable identification of *Staphylococcus aureus* harbouring the enterotoxin gene cluster (*egc*) and quantitative detection in raw milk by real time PCR. *Int J Food Microbiol* 2011; 144: 528-537.
5. Vasconcelos NG, Cunha MLRS. Staphylococcal enterotoxins: Molecular aspects and detection methods. *Journal of Public Health and Epidemiology* 2010; 2: 29-42.
6. Archer GL, Thanassi JA, Niemeyer DM, Pucci MJ. Characterization of IS1272, an insertion sequence-like element from *Staphylococcus haemolyticus*. *Antimicrob Agents Chemother* 1996; 40: 924-929.
7. Wu S, Piscitelli C, de Lencastre H, Tomasz A. Tracking the evolutionary origin of the methicillin resistance gene: cloning and sequencing of a homologue of *mecA* from a methicillin susceptible strains of *Staphylococcus sciuri*. *Microb Drug Resist* 1996; 2: 435-441.
8. Hiramatsu K, Cui L, Kuroda M, Ito T. The emergence and evolution of methicillin-resistant *Staphylococcus aureus*. *Trends Microbiol* 2001; 9: 486-493.
9. Robinson AD, Kearns AM, Holmes A *et al.* Re-emergence of early pandemic *Staphylococcus aureus* as a community-acquired methicillin-resistant clone. *Lancet* 2005; 365:1256-1258.
10. Robinson DA, Enright MC. Multilocus sequence typing and the evolution of methicillin-resistant *Staphylococcus aureus*. *Clin Microbiol Infect* 2004; 10: 92-97.
11. Oliviera DC, Tomasz A, de Lencastre H. Secrets of success of a human pathogen: molecular evolution of pandemic clones of methicillin - resistant *Staphylococcus aureus*. *The Lancet Infect Dis* 2002; 2: 180 - 189.
12. Robinson DA, Enright MC. Evolutionary models of the emergence of methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2003; 47: 3926-3934.
13. Enright MC, Robinson DA, Randle G, Feil EJ, Grundmann H, Spratt BG. The evolutionary history of methicillin-resistant *Staphylococcus aureus* (MRSA). *Proc Natl Acad Sci USA* 2002; 99: 7687-7692.
14. Melter O, Aires de Sousa M, Urbaskova P, Jakubu V, Zemlickova H, de Lencastre H. Update on the major

global types of methicillin-resistant *Staphylococcus aureus* in the Czech Republic. J Clin Microbiol 2003; 41: 4998-5005.

15. Feil EJ, Li BC, Aanensen DM *et al.* eBURST: inferring patterns of evolutionary descent among clusters of related bacterial genotypes from multilocus sequence typing data. J Bacteriol 2004; 186: 1518-1530.

16. Enright MC, Day NP, Davies CE, Peacock SJ, Spratt BG. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. J Clin Microbiol 2000; 38: 1008-1015.

17. Harmsen D, Claus H, Witte W *et al.* Typing of methicillin-resistant *Staphylococcus aureus* in a university hospital setting by using novel software for spa repeat determination and database management. J Clin Microbiol 2003; 41: 5442-5448.

18. Milheirico C, Oliveira DC, de Lancastre H. Update to the multiplex PCR strategy for assignment of *mec* element types in *Staphylococcus aureus*. Antimicrob Agents Chemother 2007; 51: 3374-3377.

19. Chung M, de Lancastre H, Matthews P *et al.* Molecular typing of methicillin-resistant *Staphylococcus aureus* by pulsed-field gel electrophoresis: comparison of results obtained in a multilaboratory effort using identical protocols and MRSA strains. Microb Drug Resist 2000; 6: 189-198.

20. Fisher A, Francois P, Holftreter S, Broeker B, Schrenzi J. Development and evaluation of a rapid strategy to determine enterotoxin gene content in *Staphylococcus aureus*. J Microbiol Methods 2004; 77: 184-190.

21. Holftreter S, Bauer K, Thomas D, Feig C. *Egc*-Encoded Superantigens from *Staphylococcus aureus* are neutralized by human sera much less efficiently than are classical staphylococcal enterotoxins or toxic shock syndrome toxin. J Clin Microbiol 2004; 72: 4061-4071.

22. Růžičková V, Voller R, Pantůček R, Petráš P, Doškař J. Multiplex PCR for detection of three exfoliative toxin serotype genes in *Staphylococcus aureus*. Folia Microbiol 2005; 50: 499-502.

23. Lina G, Piémont Y, Godail-Gamot F, Bes M. Involvement of Pantone-Valentine leukocidin-producing *Staphylococcus aureus* in primary skin infections and pneumonia. Clin Infect Dis 1999; 29: 1128-32.

24. Shopsin B, Mathema B, Alcabes P *et al.* Prevalence of *agr* Specificity Groups among *Staphylococcus aureus* Strains Colonizing Children and Their Guardians. J Clin Microbiol 2003; 41: 456-459.

25. Bergerová T, Hedlová D, Jindrák V, Urbášková P, Chmelík V. Doporučený postup pro kontrolu výskytu kmenů *Staphylococcus aureus* rezistentních k oxacilinu (MRSA) a s jinou nebezpečnou antibiotickou rezistencí ve zdravotnických zařízeních. Doporučený postup ČLS JEP. <http://www.cls.cz/dalsi-odborne-projekty>.