Reference

[1] M. Ristow a S. Schmeisser, „Extending life span by increasing oxidative stress“, *Free Radic Biol Med*, vol. 51, č. 2, s. 327–336, červenec 2011, doi: 10.1016/j.freeradbiomed.2011.05.010.

[2] Y. Wei, S. Jia, Y. Ding, S. Xia a S. Giunta, „Balanced basal-levels of ROS (redox-biology), and very-low-levels of pro-inflammatory cytokines (cold-inflammaging), as signaling molecules can prevent or slow-down overt-inflammaging, and the aging-associated decline of adaptive-homeostasis“, *Exp Gerontol*, vol. 172, s. 112067, únor 2023, doi: 10.1016/j.exger.2022.112067.

[3] D. R. Rogers, D. J. Lawlor a J. L. Moeller, „Vitamin C Supplementation and Athletic Performance: A Review“, *Curr Sports Med Rep*, vol. 22, č. 7, s. 255–259, červenec 2023, doi: 10.1249/JSR.0000000000001083.

[4] A. Thirupathi et al., „Effect of Different Exercise Modalities on Oxidative Stress: A Systematic Review“, *Biomed Res Int*, vol. 2021, č. 1, leden 2021, doi: 10.1155/2021/1947928.

[5] W. Willett, J. Manson a S. Liu, „Glycemic index, glycemic load, and risk of type 2 diabetes“, *Am J Clin Nutr*, vol. 76, č. 1, s. 274S–280S, červenec 2002, doi: 10.1093/ajcn/76.1.274S.

[6] D. Romeres, M. Schiavon, A. Basu, C. Cobelli, R. Basu a C. Dalla Man, „Exercise effect on insulin-dependent and insulin-independent glucose utilization in healthy individuals and individuals with type 1 diabetes: a modeling study“, *American Journal of Physiology-Endocrinology and Metabolism*, vol. 321, č. 1, s. E122–E129, červenec 2021, doi: 10.1152/ajpendo.00084.2021.

[7] A. J. Rose a E. A. Richter, „Skeletal Muscle Glucose Uptake During Exercise: How is it Regulated?“, *Physiology*, vol. 20, č. 4, s. 260–270, srpen 2005, doi: 10.1152/physiol.00012.2005.

[8] D. Pesta et al., „Determinants of increased muscle insulin sensitivity of exercise-trained versus sedentary normal weight and overweight individuals“, *Sci Adv*, vol. 11, č. 1, leden 2025, doi: 10.1126/sciadv.adr8849.

[9] H.-C. Lee a Y.-H. Wei, „Mitochondrial biogenesis and mitochondrial DNA maintenance of mammalian cells under oxidative stress“, *Int J Biochem Cell Biol*, vol. 37, č. 4, s. 822–834, duben 2005, doi: 10.1016/j.biocel.2004.09.010.

[10] I. Y. Kuo a B. E. Ehrlich, „Signaling in Muscle Contraction“, *Cold Spring Harb Perspect Biol*, vol. 7, č. 2, s. a006023, únor 2015, doi: 10.1101/cshperspect.a006023.

[11] F. S. Sjöstrand, „The Ultrastructure of Cells as Revealed by the Electron Microscope“, 1956, s. 455–533. doi: 10.1016/S0074-7696(08)62578-4.

[12] C. Lhuissier et al., „Mitochondrial F0F1-ATP synthase governs the induction of mitochondrial fission“, *iScience*, vol. 27, č. 5, s. 109808, květen 2024, doi: 10.1016/j.isci.2024.109808.

[13] D. Sorriento, E. Di Vaia a G. Iaccarino, „Physical Exercise: A Novel Tool to Protect Mitochondrial Health“, *Front Physiol*, vol. 12, duben 2021, doi: 10.3389/fphys.2021.660068.

[14] M. W. Gray, „Mitochondrial Evolution“, *Cold Spring Harb Perspect Biol*, vol. 4, č. 9, s. a011403–a011403, září 2012, doi: 10.1101/cshperspect.a011403.

[15] A. N. Oliveira a D. A. Hood, „Exercise is mitochondrial medicine for muscle“, *Sports Medicine and Health Science*, vol. 1, č. 1, s. 11–18, prosinec 2019, doi: 10.1016/j.smhs.2019.08.008.

[16] J. O. Holloszy, „Biochemical Adaptations in Muscle“, *Journal of Biological Chemistry*, vol. 242, č. 9, s. 2278–2282, květen 1967, doi: 10.1016/S0021-9258(18)96046-1.

[17] A.-L. Tardy, E. Pouteau, D. Marquez, C. Yilmaz a A. Scholey, „Vitamins and Minerals for Energy, Fatigue and Cognition: A Narrative Review of the Biochemical and Clinical Evidence“, *Nutrients*, vol. 12, č. 1, s. 228, leden 2020, doi: 10.3390/nu12010228.

[18] Z. El-Gammal, M. A. Nasr, A. O. Elmehrath, R. A. Salah, S. M. Saad a N. El-Badri, „Regulation of mitochondrial temperature in health and disease“, *Pflugers Arch*, vol. 474, č. 10, s. 1043–1051, říjen 2022, doi: 10.1007/s00424-022-02719-2.

[19] J. D. Browning, J. Baxter, S. Satapati a S. C. Burgess, „The effect of short-term fasting on liver and skeletal muscle lipid, glucose, and energy metabolism in healthy women and men“, *J Lipid Res*, vol. 53, č. 3, s. 577–586, březen 2012, doi: 10.1194/jlr.P020867.

[20] S. Z. Sun a M. W. Empie, „Fructose metabolism in humans – what isotopic tracer studies tell us“, *Nutr Metab (Lond)*, vol. 9, č. 1, s. 89, 2012, doi: 10.1186/1743-7075-9-89.

[21] S. Jung, H. Bae, W.-S. Song a C. Jang, „Dietary Fructose and Fructose-Induced Pathologies“, *Annu Rev Nutr*, vol. 42, č. 1, s. 45–66, srpen 2022, doi: 10.1146/annurev-nutr-062220-025831.

[22] P. Polčic, „Naming the Cycle: On the Etymology of the Citric Acid Cycle Intermediates“, *J Chem Educ*, vol. 95, č. 10, s. 1894–1896, Oct. 2018, doi: 10.1021/acs.jchemed.8b00059.

[23] N. C. Williams a L. A. J. O'Neill, „A Role for the Krebs Cycle Intermediate Citrate in Metabolic Reprogramming in Innate Immunity and

Inflammation", *Front Immunol*, vol. 9, únor 2018, doi: 10.3389/fimmu.2018.00141.

[24] A. Schurr, „How the 'Aerobic/Anaerobic Glycolysis' Meme Formed a 'Habit of Mind' Which Impedes Progress in the Field of Brain Energy Metabolism", *Int J Mol Sci*, vol. 25, č. 3, s. 1433, leden 2024, doi: 10.3390/ijms25031433.

[25] I. G. Darvey, „What factors are responsible for the greater yield of ATP per carbon atom when fatty acids are completely oxidised to CO2 and water compared with glucose?", *Biochem Educ*, vol. 27, č. 4, s. 209–210, říjen 1999, doi: 10.1016/S0307-4412(98)00269-6.

[26] M. Hargreaves a L. L. Spriet, „Skeletal muscle energy metabolism during exercise", *Nat Metab*, vol. 2, č. 9, s. 817–828, srpen 2020, doi: 10.1038/s42255-020-0251-4.

[27] M. J. Gibala, „Protein Metabolism and Endurance Exercise", *Sports Medicine*, vol. 37, č. 4, s. 337–340, 2007, doi: 10.2165/00007256-200737-040-00016.

[28] M. Clauss a J. Jensen, „Effect of Exercise Intensity, Duration, and Volume on Protein Oxidation During Endurance Exercise in Humans: A Systematic Review With Meta-Analysis", *Scand J Med Sci Sports*, vol. 35, č. 4, duben 2025, doi: 10.1111/sms.70038.

[29] P. Knuiman, M. T. E. Hopman, C. Verbruggen a M. Mensink, „Protein and the Adaptive Response With Endurance Training: Wishful Thinking or a Competitive Edge?", *Front Physiol*, vol. 9, květen 2018, doi: 10.3389/fphys.2018.00598.

[30] J. Achten a A. E. Jeukendrup, „Optimizing fat oxidation through exercise and diet", *Nutrition*, vol. 20, č. 7–8, s. 716–727, červenec 2004, doi: 10.1016/j.nut.2004.04.005.

[31] A. Coletta, D. L. Thompson a H. A. Raynor, „The influence of commercially-available carbohydrate and carbohydrate-protein supplements on endurance running performance in recreational athletes during a field trial", *J Int Soc Sports Nutr*, vol. 10, č. 1, leden 2013, doi: 10.1186/1550-2783-10-17.

[32] C. Kerksick et al., „International Society of Sports Nutrition position stand: Nutrient timing", *J Int Soc Sports Nutr*, vol. 5, č. 1, leden 2008, doi: 10.1186/1550-2783-5-17.

[33] B. Murray a C. Rosenbloom, „Fundamentals of glycogen metabolism for coaches and athletes", *Nutr Rev*, vol. 76, č. 4, s. 243–259, duben 2018, doi: 10.1093/nutrit/nuy001.

[34] M. Kanter, „High-Quality Carbohydrates and Physical Performance", *Nutr Today*, vol. 53, č. 1, s. 35–39, leden 2018, doi: 10.1097/NT.0000000000000238.

[35] D. J. Dearlove, O. K. Faull a K. Clarke, „Context is key: exogenous ketosis and athletic performance“, *Curr Opin Physiol*, vol. 10, s. 81–89, srpen 2019, doi: 10.1016/j.cophys.2019.04.010.

[36] P. J. Cox et al., „Nutritional Ketosis Alters Fuel Preference and Thereby Endurance Performance in Athletes“, *Cell Metab*, vol. 24, č. 2, s. 256–268, srpen 2016, doi: 10.1016/j.cmet.2016.07.010.

[37] D. J. Dearlove, O. K. Faull, E. Rolls, K. Clarke a P. J. Cox, „Nutritional Ketoacidosis During Incremental Exercise in Healthy Athletes“, *Front Physiol*, vol. 10, březen 2019, doi: 10.3389/fphys.2019.00290.

[38] C. P. Bailey a E. Hennessy, „A review of the ketogenic diet for endurance athletes: performance enhancer or placebo effect?“, *J Int Soc Sports Nutr*, vol. 17, č. 1, leden 2020, doi: 10.1186/s12970-020-00362-9.

[39] C. P. Bailey a E. Hennessy, „A review of the ketogenic diet for endurance athletes: performance enhancer or placebo effect?“, *J Int Soc Sports Nutr*, vol. 17, č. 1, leden 2020, doi: 10.1186/s12970-020-00362-9.

[40] S. N. BLAIR et al., „Physical activity, nutrition, and chronic disease“, *Med Sci Sports Exerc*, vol. 28, č. 3, s. 335–349, březen 1996, doi: 10.1097/00005768-199603000-00009.

[41] J. L. Veerman et al., „Physical activity and life expectancy: a life-table analysis“, *Br J Sports Med*, p. bjsports-2024-108125, listopad 2024, doi: 10.1136/bjsports-2024-108125.

[42] J. L. Veerman et al., „Television viewing time and reduced life expectancy: a life table analysis“, *Br J Sports Med*, vol. 46, č. 13, s. 927–930, říjen 2012, doi: 10.1136/bjsports-2011-085662.

[43] C. E. Morris, J. C. Garner, S. G. Owens, M. W. Valliant, H. Debusk a M. Loftin, „A Prospective Study Comparing Distance-based vs. Time-based Exercise Prescriptions of Walking and Running in Previously Sedentary Overweight Adults“, *Int J Exerc Sci*, vol. 10, č. 5, s. 782–797, 2017.

[44] P. T. Williams, „Greater weight loss from running than walking during a 6.2-yr prospective follow-up“, *Med Sci Sports Exerc*, vol. 45, č. 4, s. 706–13, duben 2013, doi: 10.1249/MSS.0b013e31827b0d0a.

[45] P. T. Williams a P. D. Thompson, „Walking versus running for hypertension, cholesterol, and diabetes mellitus risk reduction“, *Arterioscler Thromb Vasc Biol*, vol. 33, č. 5, s. 1085–1091, květen 2013, doi: 10.1161/ATVBAHA.112.300878.

[46] J. Jeppesen et al., „Enhanced Fatty Acid Oxidation and FATP4 Protein Expression after Endurance Exercise Training in Human Skeletal Muscle“, *PLOS One*, vol. 7, č. 1, s. e29391, leden 2012, doi: 10.1371/journal.pone.0029391.

[47] L. S. Sidossis, R. R. Wolfe a A. R. Coggan, „Regulation of fatty acid oxidation in untrained vs. trained men during exercise“, *American Journal*

of Physiology-Endocrinology and Metabolism, vol. 274, č. 3, s. E510–E515, březen 1998, doi: 10.1152/ajpendo.1998.274.3.E510.

[48] E. A. Richter, L. Sylow a M. Hargreaves, „Interactions between insulin and exercise", *Biochemical Journal*, vol. 478, č. 21, s. 3827–3846, listopad 2021, doi: 10.1042/BCJ20210185.

[49] A. M. Fritzen, A.-M. Lundsgaard a B. Kiens, „Tuning fatty acid oxidation in skeletal muscle with dietary fat and exercise", *Nat Rev Endocrinol*, vol. 16, č. 12, s. 683–696, prosinec 2020, doi: 10.1038/s41574-020-0405-1.

[50] C. A. Slentz, J. A. Houmard a W. E. Kraus, „Exercise, Abdominal Obesity, Skeletal Muscle, and Metabolic Risk: Evidence for a Dose Response", *Obesity*, vol. 17, č. S3, prosinec 2009, doi: 10.1038/oby.2009.385.

[51] F. Zurlo, K. Larson, C. Bogardus a E. Ravussin, „Skeletal muscle metabolism is a major determinant of resting energy expenditure", *Journal of Clinical Investigation*, vol. 86, č. 5, s. 1423–1427, listopad 1990, doi: 10.1172/JCI114857.

[52] R. Bruellman, S. Pahlen, J. M. Ellingson, R. P. Corley, S. J. Wadsworth a C. A. Reynolds, „A twin-driven analysis on early aging biomarkers and associations with sitting-time and physical activity", *PLOS One*, vol. 19, č. 9, s. e0308660, září 2024, doi: 10.1371/journal.pone.0308660.

[53] G. Perseghin et al., „Habitual Physical Activity Is Associated With Intrahepatic Fat Content in Humans", *Diabetes Care*, vol. 30, č. 3, s. 683–688, březen 2007, doi: 10.2337/dc06-2032.

[54] S. Zelber-Sagi et al., „Role of leisure-time physical activity in nonalcoholic fatty liver disease: A population-based study", *Hepatology*, vol. 48, č. 6, s. 1791–1798, prosinec 2008, doi: 10.1002/hep.22525.

[55] G. N. Healy et al., „Breaks in Sedentary Time", *Diabetes Care*, vol. 31, č. 4, s. 661–666, duben 2008, doi: 10.2337/dc07-2046.

[56] G. N. Healy et al., „Breaks in Sedentary Time", *Diabetes Care*, vol. 31, č. 4, s. 661–666, duben 2008, doi: 10.2337/dc07-2046.

[57] A. Grøntved, „Television Viewing and Risk of Type 2 Diabetes, Cardiovascular Disease, and All-Cause Mortality", *JAMA*, vol. 305, č. 23, s. 2448, červen 2011, doi: 10.1001/jama.2011.812.

[58] P. T. Williams a P. D. Thompson, „Walking Versus Running for Hypertension, Cholesterol, and Diabetes Mellitus Risk Reduction", *Arterioscler Thromb Vasc Biol*, vol. 33, č. 5, s. 1085–1091, květen 2013, doi: 10.1161/ATVBAHA.112.300878.

[59] B. Strasser, „Survival of the fittest VO sub 2 sub max a key predictor of longevity", *Frontiers in Bioscience*, vol. 23, č. 8, s. 4657, 2018, doi: 10.2741/4657.

[60] C. J. Williams et al., „Genes to predict VO2max trainability: a systematic review“, *BMC Genomics*, vol. 18, č. S8, s. 831, listopad 2017, doi: 10.1186/s12864-017-4192-6.

[61] B. Strasser, „Survival of the fittest VO sub 2 sub max a key predictor of longevity“, *Frontiers in Bioscience*, vol. 23, č. 8, s. 4657, 2018, doi: 10.2741/4657.

[62] T. Scribbans, S. Vecsey, H. Paul, W. Foster a B. Gurd, „The Effect of Training Intensity on VO2max in Young Healthy Adults: A Meta-Regression and Meta-Analysis“, *Int J Exerc Sci*, vol. 9, č. 2, s. 230–247, 2016, doi: 10.70252/HHBR9374.

[63] K. Mandsager, S. Harb, P. Cremer, D. Phelan, S. E. Nissen a W. Jaber, „Association of Cardiorespiratory Fitness With Long-term Mortality Among Adults Undergoing Exercise Treadmill Testing“, *JAMA Netw Open*, vol. 1, č. 6, s. e183605, říjen 2018, doi: 10.1001/jamanetworkopen.2018.3605.

[64] F. Beignon, N. Gueguen, H. Tricoire-Leignel, C. Mattei a G. Lenaers, „The multiple facets of mitochondrial regulations controlling cellular thermogenesis“, *Cellular and Molecular Life Sciences*, vol. 79, č. 10, s. 525, říjen 2022, doi: 10.1007/s00018-022-04523-8.

[65] S. Cadenas, „Mitochondrial uncoupling, ROS generation and cardioprotection“, *Biochimica et Biophysica Acta (BBA) – Bioenergetics*, vol. 1859, č. 9, s. 940–950, září 2018, doi: 10.1016/j.bbabio.2018.05.019.

[66] M.-G. Kang et al., „Mitochondrial Thermogenesis Can Trigger Heat Shock Response in the Nucleus“, *ACS Cent Sci*, vol. 10, č. 6, s. 1231–1241, červen 2024, doi: 10.1021/acscentsci.3c01589.

[67] S. A. Dodonova et al., „Synephrine and Its Derivative Compound A: Common and Specific Biological Effects“, *Int J Mol Sci*, vol. 24, č. 24, s. 17537, prosinec 2023, doi: 10.3390/ijms242417537.

[68] B. Nielsen a L. Nybo, „Cerebral Changes During Exercise in the Heat“, *Sports Medicine*, vol. 33, č. 1, s. 1–11, 2003, doi: 10.2165/00007256-200333010-00001.

[69] D. Gagnon a C. G. Crandall, „Sweating as a heat loss thermoeffector“, 2018, s. 211–232. doi: 10.1016/B978-0-444-63912-7.00013-8.

[70] S. K. Riebl a B. M. Davy, „The Hydration Equation“, *ACSMs Health Fit J*, vol. 17, č. 6, s. 21–28, listopad 2013, doi: 10.1249/FIT.0b013e3182a9570f.

[71] S. N. Cheuvront, K. J. Sollanek a R. W. Kenefick, „Forecasting individual exercise sweat losses from forecast air temperature and energy expenditure“, *Front Sports Act Living*, vol. 5, prosinec 2023, doi: 10.3389/fspor.2023.1277070.

[72] J. D. Périard, T. M. H. Eijsvogels a H. A. M. Daanen, „Exercise under heat stress: thermoregulation, hydration, performance implications, and mitigation strategies“, *Physiol Rev*, vol. 101, č. 4, s. 1873–1979, říjen 2021, doi: 10.1152/physrev.00038.2020.

[73] R. J. Maughan, P. Watson a S. M. Shirreffs, „Implications of active lifestyles and environmental factors for water needs and consequences of failure to meet those needs“, *Nutr Rev*, vol. 73, č. suppl 2, s. 130–140, září 2015, doi: 10.1093/nutrit/nuv051.

[74] A. P. Jung, P. A. Bishop, A. Al-Nawwas a R. B. Dale, „Influence of Hydration and Electrolyte Supplementation on Incidence and Time to Onset of Exercise-Associated Muscle Cramps“, *J Athl Train*, vol. 40, č. 2, s. 71–75, červen 2005.

[75] A. J. Siegel, „Fatal Water Intoxication and Cardiac Arrest in Runners During Marathons: Prevention and Treatment Based on Validated Clinical Paradigms“, *Am J Med*, vol. 128, č. 10, s. 1070–1075, říjen 2015, doi: 10.1016/j.amjmed.2015.03.031.

[76] C. S. D. Almond et al., „Hyponatremia among Runners in the Boston Marathon“, *New England Journal of Medicine*, vol. 352, č. 15, s. 1550–1556, duben 2005, doi: 10.1056/NEJMoa043901.

[77] A. Arab, N. Rafie, R. Amani a F. Shirani, „The Role of Magnesium in Sleep Health: a Systematic Review of Available Literature“, *Biol Trace Elem Res*, vol. 201, č. 1, s. 121–128, leden 2023, doi: 10.1007/s12011-022-03162-1.

[78] B. Murray a C. Rosenbloom, „Fundamentals of glycogen metabolism for coaches and athletes“, *Nutr Rev*, vol. 76, č. 4, s. 243–259, duben 2018, doi: 10.1093/nutrit/nuy001.

[79] C. M. Kerksick et al., „International society of sports nutrition position stand: nutrient timing“, *J Int Soc Sports Nutr*, vol. 14, č. 1, leden 2017, doi: 10.1186/s12970-017-0189-4.

[80] R. F. G. J. King, B. Jones a J. P. O'Hara, „The availability of water associated with glycogen during dehydration: a reservoir or raindrop?“, *Eur J Appl Physiol*, vol. 118, č. 2, s. 283–290, únor 2018, doi: 10.1007/s00421-017-3768-9.

[81] N. T. Bendsen et al., „Is beer consumption related to measures of abdominal and general obesity? A systematic review and meta-analysis“, *Nutr Rev*, vol. 71, č. 2, s. 67–87, únor 2013, doi: 10.1111/j.1753--4887.2012.00548.x.

[82] A. Kwok, A. L. Dordevic, G. Paton, M. J. Page a H. Truby, „Effect of alcohol consumption on food energy intake: a systematic review and meta-analysis“, *British Journal of Nutrition*, vol. 121, č. 5, s. 481–495, květen 2019, doi: 10.1017/S0007114518003677.

[83] T. W. Elffers et al., „Body fat distribution, in particular visceral fat, is associated with cardiometabolic risk factors in obese women", *PLOS One*, vol. 12, č. 9, s. e0185403, září 2017, doi: 10.1371/journal.pone.0185403.

[84] S. Dhokte a K. Czaja, „Visceral Adipose Tissue: The Hidden Culprit for Type 2 Diabetes", *Nutrients*, vol. 16, č. 7, s. 1015, březen 2024, doi: 10.3390/nu16071015.

[85] S. H. Jung, K. H. Ha a D. J. Kim, „Visceral Fat Mass Has Stronger Associations with Diabetes and Prediabetes than Other Anthropometric Obesity Indicators among Korean Adults", *Yonsei Med J*, vol. 57, č. 3, s. 674, 2016, doi: 10.3349/ymj.2016.57.3.674.

[86] A. Topiwala, K. P. Ebmeier, T. Maullin-Sapey a T. E. Nichols, „Alcohol consumption and MRI markers of brain structure and function: Cohort study of 25,378 UK Biobank participants", *Neuroimage Clin*, vol. 35, s. 103066, 2022, doi: 10.1016/j.nicl.2022.103066.

[87] R. Burton a N. Sheron, „No level of alcohol consumption improves health", *The Lancet*, vol. 392, č. 10152, s. 987–988, září 2018, doi: 10.1016/S0140-6736(18)31571-X.

[88] L. Jophlin, T.-Y. Liu a C. J. McClain, „Nutritional deficiencies in alcohol use disorder/alcohol-associated liver disease", *Curr Opin Gastroenterol*, vol. 40, č. 2, s. 112–117, březen 2024, doi: 10.1097/MOG.0000000000000999.

[89] E. B. Flink, „Mineral Metabolism in Alcoholism", in *The Biology of Alcoholism*, Boston, MA: Springer US, 1971, s. 377–395. doi: 10.1007/978-1-4615-6525-3_12.

[90] A. S. Bell, E. M. O'Connell a F. W. Lohoff, „Cholesterol and alcohol", in *Cholesterol*, Elsevier, 2022, s. 747–767. doi: 10.1016/B978-0-323-85857-1.00036-5.

[91] M. L. Muiesan, C. Agabiti-Rosei, A. Paini a M. Salvetti, „Uric Acid and Cardiovascular Disease: An Update", *Eur Cardiol*, vol. 11, č. 1, s. 54–59, srpen 2016, doi: 10.15420/ecr.2016:4:2.

[92] P. Memar a F. Faradji, „A Novel Multi-Class EEG-Based Sleep Stage Classification System", *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, vol. 26, č. 1, s. 84–95, leden 2018, doi: 10.1109/TNSRE.2017.2776149.

[93] M. G. Frank a H. C. Heller, „The Function(s) of Sleep", 2018, s. 3–34. doi: 10.1007/164_2018_140.

[94] E. C. Lingas, „A Narrative Review of the Carcinogenic Effect of Night Shift and the Potential Protective Role of Melatonin", *Cureus*, vol. 15, č. 23, srpen 2023, doi: 10.7759/cureus.43326.

[117] X. Yang et al., „Association of plant-based diet and type 2 diabetes mellitus in Chinese rural adults: The Henan Rural Cohort Study", *J Diabetes Investig*, vol. 12, č. 9, s. 1569–1576, září 2021, doi: 10.1111/jdi.13522.

[118] H. Peña-Jorquera et al., „Plant-Based Nutrition: Exploring Health Benefits for Atherosclerosis, Chronic Diseases, and Metabolic Syndrome—A Comprehensive Review", *Nutrients*, vol. 15, č. 14, s. 3244, červenec 2023, doi: 10.3390/nu15143244.

[119] T. L. Halton a F. B. Hu, „The Effects of High Protein Diets on Thermogenesis, Satiety and Weight Loss: A Critical Review", *J Am Coll Nutr*, vol. 23, č. 5, s. 373–385, říjen 2004, doi: 10.1080/07315724.2004.10719381.

[120] S. Sieri et al., „Dietary Intakes of Animal and Plant Proteins and Risk of Colorectal Cancer: The EPIC-Italy Cohort", *Cancers (Basel)*, vol. 14, č. 12, s. 2917, červen 2022, doi: 10.3390/cancers14122917.

[121] N. Fleming, „Protein mania", *New Sci (1956)*, vol. 242, č. 3226, s. 38–41, duben 2019, doi: 10.1016/S0262-4079(19)30695-5.

[122] X. Zhang et al., „Identification of a leucine-mediated threshold effect governing macrophage mTOR signalling and cardiovascular risk", *Nat Metab*, vol. 6, č. 2, s. 359–377, únor 2024, doi: 10.1038/s42255-024-00984-2.

[123] M. Beltrá, F. Borrás a A. B. Ropero, „Are Foods with Protein Claims Healthy? A Study of the Spanish Market", *Nutrients*, vol. 16, č. 24, s. 4281, prosinec 2024, doi: 10.3390/nu16244281.

[124] A. Y. Coqueiro, M. M. Rogero a J. Tirapegui, „Glutamine as an Anti-Fatigue Amino Acid in Sports Nutrition", *Nutrients*, vol. 11, č. 4, s. 863, duben 2019, doi: 10.3390/nu11040863.

[125] A. Kurz, „Physiology of Thermoregulation", *Best Pract Res Clin Anaesthesiol*, vol. 22, č. 4, s. 627–644, prosinec 2008, doi: 10.1016/j.bpa.2008.06.004.

[126] S. D. R. Galloway a R. J. Maughan, „Effects of ambient temperature on the capacity to perform prolonged cycle exercise in man", *Medicine & Science in Sports & Exercise*, vol. 29, č. 9, s. 1240–1249, září 1997, doi: 10.1097/00005768-199709000-00018.

[127] N. El Helou et al., „Impact of Environmental Parameters on Marathon Running Performance", *PLOS One*, vol. 7, č. 5, s. e37407, květen 2012, doi: 10.1371/journal.pone.0037407.

[128] H. Gatterer et al., „Practicing Sport in Cold Environments: Practical Recommendations to Improve Sport Performance and Reduce Negative Health Outcomes", *Int J Environ Res Public Health*, vol. 18, č. 18, s. 9700, září 2021, doi: 10.3390/ijerph18189700.

[129] U. Bergh a B. Ekblom, „Influence of muscle temperature on maximal muscle strength and power output in human skeletal muscles", *Acta*

Physiol Scand, vol. 107, č. 1, s. 33–37, září 1979, doi: 10.1111/j.1748-1716.1979.tb06439.x.

[130] J. W. Castellani, A. J. Young, M. B. Ducharme, G. G. Giesbrecht, E. Glickman a R. E. Sallis, „Prevention of Cold Injuries during Exercise", *Med Sci Sports Exerc*, vol. 38, č. 11, s. 2012–2029, listopad 2006, doi: 10.1249/01.mss.0000241641.75101.64.

[131] H. Pohjantähti, J. Laitinen a J. Parkkari, „Exercise-induced bronchospasm among healthy elite cross country skiers and non-athletic students", *Scand J Med Sci Sports*, vol. 15, č. 5, s. 324–328, říjen 2005, doi: 10.1111/j.1600-0838.2004.00423.x.

[132] D. R. Pendergast, „The effect of body cooling on oxygen transport during exercise", *Med Sci Sports Exerc*, vol. 20, č. Sup 1, s. S171–S176, říjen 1988, doi: 10.1249/00005768-198810001-00015.

[133] T. Stensrud, S. Berntsen a K.-H. Carlsen, „Exercise capacity and exercise-induced bronchoconstriction (EIB) in a cold environment", *Respir Med*, vol. 101, č. 7, s. 1529–1536, červenec 2007, doi: 10.1016/j.rmed.2006.12.011.

[134] J. W. Castellani a M. J. Tipton, „Cold Stress Effects on Exposure Tolerance and Exercise Performance", in *Comprehensive Physiology*, Wiley, 2015, s. 443–469. doi: 10.1002/cphy.c140081.

[135] M. Nimmo, „Exercise in the cold", *J Sports Sci*, vol. 22, č. 10, s. 898–916, říjen 2004, doi: 10.1080/02640414000058883.

[136] J. D. Layden, M. J. Patterson a M. A. Nimmo, „Effects of reduced ambient temperature on fat utilization during submaximal exercise", *Med Sci Sports Exerc*, vol. 34, č. 5, s. 774–779, květen 2002, doi: 10.1097/00005768-200205000-00008.

[137] B. Kingma, „The thermoneutral zone implications for metabolic studies", *Frontiers in Bioscience*, vol. E4, č. 5, s. 518, 2012, doi: 10.2741/e518.

[138] D. D. Gagnon et al., „Cold exposure enhances fat utilization but not non-esterified fatty acids, glycerol or catecholamines availability during submaximal walking and running", *Front Physiol*, vol. 4, 2013, doi: 10.3389/fphys.2013.00099.

[139] K. Charlot, C. Faure a S. Antoine-Jonville, „Influence of Hot and Cold Environments on the Regulation of Energy Balance Following a Single Exercise Session: A Mini-Review", *Nutrients*, vol. 9, č. 6, s. 592, červen 2017, doi: 10.3390/nu9060592.

[140] E. F. Foxman et al., „Temperature-dependent innate defense against the common cold virus limits viral replication at warm temperature in mouse airway cells", *Proceedings of the National Academy of Sciences*, vol. 112, č. 3, s. 827–832, leden 2015, doi: 10.1073/pnas.1411030112.

[141] N. V. Margaritelis et al., „Muscle damage and inflammation after eccentric exercise: can the repeated bout effect be removed?“, *Physiol Rep*, vol. 3, č. 12, s. e12648, prosinec 2015, doi: 10.14814/phy2.12648.

[142] L. Forcina, M. Cosentino a A. Musarò, „Mechanisms Regulating Muscle Regeneration: Insights into the Interrelated and Time-Dependent Phases of Tissue Healing“, *Cells*, vol. 9, č. 5, s. 1297, květen 2020, doi: 10.3390/cells9051297.

[143] T. Citherlet, F. Crettaz von Roten, B. Kayser a K. Guex, „Acute Effects of the Wim Hof Breathing Method on Repeated Sprint Ability: A Pilot Study“, *Front Sports Act Living*, vol. 3, srpen 2021, doi: 10.3389/fspor.2021.700757.

[144] O. Almahayni a L. Hammond, „Does the Wim Hof Method have a beneficial impact on physiological and psychological outcomes in healthy and non-healthy participants? A systematic review“, *PLOS One*, vol. 19, č. 3, s. e0286933, březen 2024, doi: 10.1371/journal.pone.0286933.

[145] E. L. Reed et al., „Cardiovascular and mood responses to an acute bout of cold water immersion“, *J Therm Biol*, vol. 118, s. 103727, prosinec 2023, doi: 10.1016/j.jtherbio.2023.103727.

[146] N. A. Shevchuk, „Adapted cold shower as a potential treatment for depression“, *Med Hypotheses*, vol. 70, č. 5, s. 995–1001, leden 2008, doi: 10.1016/j.mehy.2007.04.052.

[147] I. Kralová Lesná, J. Rychlíková, L. Vávrová a S. Vybíral, „Could human cold adaptation decrease the risk of cardiovascular disease?“, *J Therm Biol*, vol. 52, s. 192–198, srpen 2015, doi: 10.1016/j.jtherbio.2015.07.007.

[148] V. Tibenská et al., „Gradual cold acclimation induces cardioprotection without affecting β-adrenergic receptor-mediated adenylyl cyclase signaling“, *J Appl Physiol*, vol. 128, č. 4, s. 1023–1032, duben 2020, doi: 10.1152/japplphysiol.00511.2019.

[149] J. Leeder, C. Gissane, K. van Someren, W. Gregson a G. Howatson, „Cold water immersion and recovery from strenuous exercise: a meta-analysis“, *Br J Sports Med*, vol. 46, č. 4, s. 233–240, březen 2012, doi: 10.1136/bjsports-2011-090061.

[150] E. S. Malta, Y. M. Dutra, J. R. Broatch, D. J. Bishop a A. M. Zagatto, „*The Effects of Regular Cold-Water Immersion* Use on Training-Induced Changes in Strength and Endurance Performance: *A Systematic Review* with Meta-Analysis“, *Sports Medicine*, vol. 51, č. 1, s. 161–174, leden 2021, doi: 10.1007/s40279-020-01362-0.

[151] K. M. Slattery, L. K. Wallace, A. J. Murphy a A. J. Coutts, „Physiological *Determinants of Three-Kilometer Running Performance in Experienced*

Triathletes", *The Journal of Strength and Conditioning Research*, vol. 20, č. 1, s. 47, 2006, doi: 10.1519/R-16724.1.

[152] R. A. Jacobs et al., „Determinants of time trial performance and maximal incremental exercise in highly trained endurance athletes", *J Appl Physiol*, vol. 111, č. 5, s. 1422–1430, listopad 2011, doi: 10.1152/japplphysiol.00625.2011.

[153] D. R. Bassett, „Limiting factors for maximum oxygen uptake and determinants of endurance performance", *Med Sci Sports Exerc*, s. 70, leden 2000, doi: 10.1097/00005768-200001000-00012.

[154] C. Lundby a R. A. Jacobs, „Adaptations of skeletal muscle mitochondria to exercise training", *Exp Physiol*, vol. 101, č. 1, s. 17–22, leden 2016, doi: 10.1113/EP085319.

[155] A. -K. Meinild Lundby et al., „Exercise training increases skeletal muscle mitochondrial volume density by enlargement of existing mitochondria and not de novo biogenesis", *Acta Physiologica*, vol. 222, č. 1, leden 2018, doi: 10.1111/apha.12905.

[156] A. Murray a M. Cardinale, „Cold applications for recovery in adolescent athletes: a systematic review and meta analysis", *Extrem Physiol Med*, vol. 4, č. 1, s. 17, prosinec 2015, doi: 10.1186/s13728-015-0035-8.

[157] R. Allan et al., „Postexercise cold water immersion modulates skeletal muscle PGC-1α mRNA expression in immersed and nonimmersed limbs: evidence of systemic regulation", *J Appl Physiol*, vol. 123, č. 2, s. 451–459, srpen 2017, doi: 10.1152/japplphysiol.00096.2017.

[158] C. H. Joo et al., „Passive and post-exercise cold-water immersion augments PGC-1α and VEGF expression in human skeletal muscle", *Eur J Appl Physiol*, vol. 116, č. 11–12, s. 2315–2326, prosinec 2016, doi: 10.1007/s00421-016-3480-1.

[159] F. Lateef, „Post exercise ice water immersion: Is it a form of active recovery?", *J Emerg Trauma Shock*, vol. 3, č. 3, s. 302, 2010, doi: 10.4103/0974-2700.66570.

[160] L. L. Smith, „Cytokine hypothesis of overtraining: a physiological adaptation to excessive stress?", *Med Sci Sports Exerc*, vol. 32, č. 2, s. 317, únor 2000, doi: 10.1097/00005768-200002000-00011.

[161] G. Banfi, G. Lombardi, A. Colombini a G. Melegati, „Whole-Body Cryotherapy in Athletes", *Sports Medicine*, vol. 40, č. 6, s. 509–517, červen 2010, doi: 10.2165/11531940-000000000-00000.

[162] J. M. Peake et al., „The effects of cold water immersion and active recovery on inflammation and cell stress responses in human skeletal muscle after resistance exercise", *J Physiol*, vol. 595, č. 3, s. 695–711, únor 2017, doi: 10.1113/JP272881.

[163] E. Heyman, B. De Geus, I. Mertens a R. Meeusen, „Effects of Four Recovery Methods on Repeated Maximal Rock Climbing Performance“, *Med Sci Sports Exerc*, vol. 41, č. 6, s. 1303–1310, červen 2009, doi: 10.1249/MSS.0b013e318195107d.

[164] H. Wang, L. Wang a Y. Pan, „Impact of different doses of cold water immersion (duration and temperature variations) on recovery from acute exercise-induced muscle damage: a network meta-analysis“, *Front Physiol*, vol. 16, únor 2025, doi: 10.3389/fphys.2025.1525726.

[165] A. J. Peacock, „ABC of oxygen: Oxygen at high altitude“, *BMJ*, vol. 317, č. 7165, s. 1063–1066, říjen 1998, doi: 10.1136/bmj.317.7165.1063.

[166] J. Rojas-Camayo et al., „Reference values for oxygen saturation from sea level to the highest human habitation in the Andes in acclimatised persons“, *Thorax*, vol. 73, č. 8, s. 776–778, srpen 2018, doi: 10.1136/thoraxjnl-2017-210598.

[167] L. A. Garvican-Lewis, I. Halliday, C. R. Abbiss, P. U. Saunders a C. J. Gore, „Altitude Exposure at 1800 m Increases Haemoglobin Mass in Distance Runners“, *J Sports Sci Med*, vol. 14, č. 2, s. 413–7, červen 2015.

[168] R. A. Jacobs et al., „Twenty-eight days of exposure to 3454 m increases mitochondrial volume density in human skeletal muscle“, *J Physiol*, vol. 594, č. 5, s. 1151–1166, Mar. 2016, doi: 10.1113/JP271118.

[169] R. A. Jacobs et al., „Twenty-eight days of exposure to 3454 m increases mitochondrial volume density in human skeletal muscle“, *J Physiol*, vol. 594, č. 5, s. 1151–1166, březen 2016, doi: 10.1113/JP271118.

[170] A. J. Murray a J. A. Horscroft, „Mitochondrial function at extreme high altitude“, *J Physiol*, vol. 594, č. 5, s. 1137–1149, březen 2016, doi: 10.1113/JP270079.

[171] M. Limmer, M. de Marées a P. Platen, „Alterations in acid–base balance and high-intensity exercise performance after short-term and long-term exposure to acute normobaric hypoxic conditions“, *Sci Rep*, vol. 10, č. 1, s. 13732, srpen 2020, doi: 10.1038/s41598-020-70762-z.

[172] G. van Hall, J. A. L. Calbet, H. Søndergaard a B. Saltin, „The re-establishment of the normal blood lactate response to exercise in humans after prolonged acclimatization to altitude“, *J Physiol*, vol. 536, č. 3, s. 963–975, listopad 2001, doi: 10.1111/j.1469-7793.2001.00963.x.

[173] A. B. Zepeda, C. A. Figueroa, A. Pessoa a J. G. Farías, „Free fatty acids reduce metabolic stress and favor a stable production of heterologous proteins in Pichia pastoris“, *Brazilian Journal of Microbiology*, vol. 49, č. 4, s. 856–864, říjen 2018, doi: 10.1016/j.bjm.2018.03.008.

[174] P. D. Wagner, „Skeletal muscle angiogenesis. A possible role for hypoxia“, *Adv Exp Med Biol*, vol. 502, s. 21–38, 2001.

[175] H. Hoppeler, „Vascular Growth in Hypoxic Skeletal Muscle", *Advances in Experimental Medicine and Biology*, 1999, s. 277–286. doi: 10.1007/978-1-4615-4711-2_21.

[176] M. Vogt a H. Hoppeler, „Is Hypoxia Training Good for Muscles and Exercise Performance?", *Prog Cardiovasc Dis*, vol. 52, č. 6, s. 525–533, květen 2010, doi: 10.1016/j.pcad.2010.02.013.

[177] J. A. Sinex a R. F. Chapman, „Hypoxic training methods for improving endurance exercise performance", *J Sport Health Sci*, vol. 4, č. 4, s. 325–332, prosinec 2015, doi: 10.1016/j.jshs.2015.07.005.

[178] B. D. Levine a J. Stray-Gundersen, „'Living high-training low': effect of moderate-altitude acclimatization with low-altitude training on performance", *J Appl Physiol*, vol. 83, č. 1, s. 102–112, červenec 1997, doi: 10.1152/jappl.1997.83.1.102.

[179] D. L. Bonetti a W. G. Hopkins, „Sea-Level Exercise Performance Following Adaptation to Hypoxia", *Sports Medicine*, vol. 39, č. 2, s. 107–127, 2009, doi: 10.2165/00007256-200939020-00002.

[180] J. T. Costello, P. R. Baker, G. M. Minett, F. Bieuzen, I. B. Stewart a C. Bleakley, „Whole-body cryotherapy (extreme cold air exposure) for preventing and treating muscle soreness after exercise in adults", *Cochrane Database of Systematic Reviews*, vol. 2015, č. 9, září 2015, doi: 10.1002/14651858.CD010789.pub2.

[181] K. Mizumura a T. Taguchi, „Delayed onset muscle soreness: Involvement of neurotrophic factors", *The Journal of Physiological Sciences*, vol. 66, č. 1, s. 43–52, červen 2016, doi: 10.1007/s12576-015-0397-0.

[182] K. Cheung, P. A. Hume a L. Maxwell, „Delayed Onset Muscle Soreness", *Sports Medicine*, vol. 33, č. 2, s. 145–164, 2003, doi: 10.2165/00007256-200333020-00005.

[183] M. M. Hall, S. Rajasekaran, T. W. Thomsen a A. R. Peterson, „Lactate: Friend or Foe", *PM&R*, vol. 8, č. 3S, březen 2016, doi: 10.1016/j.pmrj.2015.10.018.

[184] S. P. Cairns, „Lactic Acid and Exercise Performance", *Sports Medicine*, vol. 36, č. 4, s. 279–291, 2006, doi: 10.2165/00007256-200636040-00001.

[185] E. Hultman a P. L. Greenhaff, „Skeletal muscle energy metabolism and fatigue during intense exercise in man", *Sci Prog*, vol. 75, č. 298 Pt 3–4, s. 361–70, 1991.

[186] M. Hargreaves a L. L. Spriet, „Skeletal muscle energy metabolism during exercise", *Nat Metab*, vol. 2, č. 9, s. 817–828, srpen 2020, doi: 10.1038/s42255-020-0251-4.

[187] W. T. Staublr, „Eccentric Action of Muscles", *Exerc Sport Sci Rev*, vol. 16, s. 157–186, 1989, doi: 10.1249/00003677-198900170-00008.

[188] E. Ferraro, A. M. Giammarioli, S. Chiandotto, I. Spoletini a G. Rosano, „Exercise-induced skeletal muscle remodeling and metabolic adaptation: redox signaling and role of autophagy“, *Antioxid Redox Signal*, vol. 21, č. 1, s. 154–76, červenec 2014, doi: 10.1089/ars.2013.5773.

[189] T. Yoshioka a A. P. Somlyo, „Calcium and magnesium contents and volume of the terminal cisternae in caffeine-treated skeletal muscle“, *J Cell Biol*, vol. 99, č. 2, s. 558–568, srpen 1984, doi: 10.1083/jcb.99.2.558.

[190] H.-Y. Chen, F.-C. Cheng, H.-C. Pan, J.-C. Hsu a M.-F. Wang, „Magnesium Enhances Exercise Performance via Increasing Glucose Availability in the Blood, Muscle, and Brain during Exercise“, *PLOS One*, vol. 9, č. 1, s. e85486, leden 2014, doi: 10.1371/journal.pone.0085486.

[191] J. C. Weavil a M. Amann, „Neuromuscular fatigue during whole body exercise“, *Curr Opin Physiol*, vol. 10, s. 128–136, srpen 2019, doi: 10.1016/j.cophys.2019.05.008.

[192] R. Meeusen, P. Watson, H. Hasegawa, B. Roelands a M. F. Piacentini, „Brain neurotransmitters in fatigue and overtraining“, *Applied Physiology, Nutrition, and Metabolism*, vol. 32, č. 5, s. 857–864, říjen 2007, doi: 10.1139/H07-080.

[193] K. Cheung, P. A. Hume a L. Maxwell, „Delayed Onset Muscle Soreness“, *Sports Medicine*, vol. 33, č. 2, s. 145–164, 2003, doi: 10.2165/00007256-200333020-00005.

[194] B. Sonkodi, „Delayed Onset Muscle Soreness and Critical Neural Microdamage-Derived Neuroinflammation“, *Biomolecules*, vol. 12, č. 9, s. 1207, srpen 2022, doi: 10.3390/biom12091207.

[195] L. S. Chow et al., „Exerkines in health, resilience and disease“, *Nat Rev Endocrinol*, vol. 18, č. 5, s. 273–289, květen 2022, doi: 10.1038/s41574-022-00641-2.

[196] G. Novelli, G. Calcaterra, F. Casciani, S. Pecorelli a J. L. Mehta, „'Exerkines': A Comprehensive Term for the Factors Produced in Response to Exercise“, *Biomedicines*, vol. 12, č. 9, s. 1975, září 2024, doi: 10.3390/biomedicines12091975.

[197] M. Mussawar, A. A. Balsom, J. O. Totosy de Zepetnek a J. L. Gordon, „The effect of physical activity on fertility: a mini-review“, *F S Rep*, vol. 4, č. 2, s. 150–158, červen 2023, doi: 10.1016/j.xfre.2023.04.005.

[198] X. Lu et al., „Exercise and exerkines: Mechanisms and roles in anti-aging and disease prevention“, *Exp Gerontol*, vol. 200, s. 112685, únor 2025, doi: 10.1016/j.exger.2025.112685.

[199] G. A. Roth et al., „Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a systematic analysis for the Global Burden of Disease Study 2017“, *The*

Lancet, vol. 392, č. 10159, s. 1736–1788, listopad 2018, doi: 10.1016/S0140-6736(18)32203-7.

[200] D. Furman et al., „Chronic inflammation in the etiology of disease across the life span", *Nat Med*, vol. 25, č. 12, s. 1822–1832, prosinec 2019, doi: 10.1038/s41591-019-0675-0.

[201] D. Morris, „Effects of Oral Lactate Consumption on Metabolism and Exercise Performance", *Curr Sports Med Rep*, vol. 11, č. 4, s. 185–188, 2012, doi: 10.1249/JSR.0b013e31825da992.

[202] J. M. Robbins a R. E. Gerszten, „Exercise, exerkines, and cardiometabolic health: from individual players to a team sport", *Journal of Clinical Investigation*, vol. 133, č. 11, červen 2023, doi: 10.1172/JCI168121.

[203] R. P. Patrick a T. L. Johnson, „Sauna use as a lifestyle practice to extend healthspan", *Exp Gerontol*, vol. 154, s. 111509, říjen 2021, doi: 10.1016/j.exger.2021.111509.

[204] T. Laukkanen, H. Khan, F. Zaccardi a J. A. Laukkanen, „Association Between Sauna Bathing and Fatal Cardiovascular and All-Cause Mortality Events", *JAMA Intern Med*, vol. 175, č. 4, s. 542, srpen 2015, doi: 10.1001/jamainternmed.2014.8187.

[205] W. Pilch et al., „Comparison of the Physiological Reactions and Physiological Strain in Healthy Men under Heat Stress in Dry and Steam Heat Saunas", *Biol Sport*, vol. 31, č. 2, s. 145–149, duben 2014, doi: 10.5604/20831862.1099045.

[206] M. Iguchi, A. E. Littmann, S.-H. Chang, L. A. Wester, J. S. Knipper a R. K. Shields, „Heat Stress and Cardiovascular, Hormonal, and Heat Shock Proteins in Humans", *J Athl Train*, vol. 47, č. 2, s. 184–190, březen 2012, doi: 10.4085/1062-6050-47.2.184.

[207] G. Pawelec, D. Goldeck a E. Derhovanessian, „Inflammation, ageing and chronic disease", *Curr Opin Immunol*, vol. 29, s. 23–28, srpen 2014, doi: 10.1016/j.coi.2014.03.007.

[208] D. Gryka, W. Pilch, M. Szarek, Z. Szygula a Ł. Tota, „The effect of sauna bathing on lipid profile in young, physically active, male subjects", *Int J Occup Med Environ Health*, vol. 27, č. 4, leden 2014, doi: 10.2478/s13382-014-0281-9.

[209] F. Zaccardi, T. Laukkanen, P. Willeit, S. K. Kunutsor, J. Kauhanen a J. A. Laukkanen, „Sauna Bathing and Incident Hypertension: A Prospective Cohort Study", *Am J Hypertens*, vol. 30, č. 11, s. 1120–1125, listopad 2017, doi: 10.1093/ajh/hpx102.

[210] S. Hillebrand et al., „Heart rate variability and first cardiovascular event in populations without known cardiovascular disease: meta-analysis and dose–response meta-regression", *EP Europace*, vol. 15, č. 5, s. 742–749, květen 2013, doi: 10.1093/europace/eus341.

[211] G. S. M. Scoon, W. G. Hopkins, S. Mayhew a J. D. Cotter, „Effect of post-exercise sauna bathing on the endurance performance of competitive male runners“, *J Sci Med Sport*, vol. 10, č. 4, s. 259–262, srpen 2007, doi: 10.1016/j.jsams.2006.06.009.

[212] A. Garolla et al., „Seminal and molecular evidence that sauna exposure affects human spermatogenesis“, *Human Reproduction*, vol. 28, č. 4, s. 877–885, srpen 2013, doi: 10.1093/humrep/det020.

[213] C. Hu et al., „Heat shock proteins: Biological functions, pathological roles, and therapeutic opportunities“, *MedComm (Beijing)*, vol. 3, č. 3, září 2022, doi: 10.1002/mco2.161.

[214] M. Mavroudi, A. Kabasakalis, A. Petridou a V. Mougios, „Blood Lactate and Maximal Lactate Accumulation Rate at Three Sprint Swimming Distances in Highly Trained and Elite Swimmers“, *Sports*, vol. 11, č. 4, s. 87, srpen 2023, doi: 10.3390/sports11040087.

[215] H. M. Piper, R. Spahr a I. Probst, „Glucose, lactate, and palmitate as substrates for the resting cardiac myocyte“, *Basic Res Cardiol*, vol. 80 Suppl 2, s. 97–101, 1985.

[216] R. Bellomo, „Bench-to-bedside review: lactate and the kidney“, *Crit Care*, vol. 6, č. 4, s. 322–6, srpen 2002, doi: 10.1186/cc1518.

[217] O. Faude, W. Kindermann a T. Meyer, „Lactate Threshold Concepts“, *Sports Medicine*, vol. 39, č. 6, s. 469–490, květen 2009, doi: 10.2165/00007256-200939060-00003.

[218] J. Mikkola, V. Vesterinen, R. Taipale, B. Capostagno, K. Häkkinen a A. Nummela, „Effect of resistance training regimens on treadmill running and neuromuscular performance in recreational endurance runners“, *J Sports Sci*, vol. 29, č. 13, s. 1359–1371, říjen 2011, doi: 10.1080/02640414.2011.589467.

[219] A. K. Ghosh, „Anaerobic threshold: its concept and role in endurance sport“, *Malays J Med Sci*, vol. 11, č. 1, s. 24–36, leden 2004.

[220] A. K. Ghosh, „Anaerobic threshold: its concept and role in endurance sport“, *Malays J Med Sci*, vol. 11, č. 1, s. 24–36, leden 2004.

[221] I. Garcia-Tabar a E. M. Gorostiaga, „A 'Blood Relationship' Between the Overlooked Minimum Lactate Equivalent and Maximal Lactate Steady State in Trained Runners. Back to the Old Days?“, *Front Physiol*, vol. 9, červenec 2018, doi: 10.3389/fphys.2018.01034.

[222] C. Foster, D. J. Fitzgerald a P. Spatz, „Stability of the blood lactate-heart rate relationship in competitive athletes“, *Med Sci Sports Exerc*, vol. 31, č. 4, s. 578–582, duben 1999, doi: 10.1097/00005768-199904000-00014.

[223] S. P. Keith, I. Jacobs a T. M. McLellan, „Adaptations to training at the individual anaerobic threshold“, *Eur J Appl Physiol Occup Physiol*, vol. 65, č. 4, s. 316–323, 1992, doi: 10.1007/BF00868134.

[224] M. J. Joyner, „Modeling: optimal marathon performance on the basis of physiological factors", *J Appl Physiol*, vol. 70, č. 2, s. 683–687, únor 1991, doi: 10.1152/jappl.1991.70.2.683.

[225] S. K. Powers a E. T. Howley, „Bez názvu", in *The Physiology of Training: Effect on $\dot{V}O_2$ Max, Performance, and Strength*, McGraw-Hill Education, 2007, ch. Exercise P.

[226] J. Zelenka, A. Dvořák a L. Alán, „L-Lactate protects skin fibroblasts against aging-Associated mitochondrial dysfunction via mitohormesis", *Oxid Med Cell Longev*, vol. 2015, 2015, doi: 10.1155/2015/351698.

[227] S.-T. Luo et al., „The Promotion of Erythropoiesis via the Regulation of Reactive Oxygen Species by Lactic Acid", *Sci Rep*, vol. 7, č. 1, s. 38105, únor 2017, doi: 10.1038/srep38105.

[228] X. Cai et al., „Lactate activates the mitochondrial electron transport chain independently of its metabolism", *Mol Cell*, vol. 83, č. 21, s. 3904-3920. e7, listopad 2023, doi: 10.1016/j.molcel.2023.09.034.

[229] L. Zhang et al., „Lactate transported by MCT1 plays an active role in promoting mitochondrial biogenesis and enhancing TCA flux in skeletal muscle", *Sci Adv*, vol. 10, č. 26, červen 2024, doi: 10.1126/sciadv.adn4508.

[230] C. Morland et al., „Exercise induces cerebral VEGF and angiogenesis via the lactate receptor HCAR1", *Nat Commun*, vol. 8, č. 1, s. 15557, květen 2017, doi: 10.1038/ncomms15557.

[231] E. Benarroch, „What Is the Role of Lactate in Brain Metabolism, Plasticity, and Neurodegeneration?", *Neurology*, vol. 102, č. 9, květen 2024, doi: 10.1212/WNL.0000000000209378.

[232] X. Xue, B. Liu, J. Hu, X. Bian a S. Lou, „The potential mechanisms of lactate in mediating exercise-enhanced cognitive function: a dual role as an energy supply substrate and a signaling molecule", *Nutr Metab (Lond)*, vol. 19, č. 1, s. 52, prosinec 2022, doi: 10.1186/s12986-022-00687-z.

[233] H. Z. Ji et al., „Relationships between athletic ability and academic performance in primary school students: A 3-year follow-up study", *Front Public Health*, vol. 10, leden 2023, doi: 10.3389/fpubh.2022.1012757.

[234] R. L. Hughes a H. D. Holscher, „Fueling Gut Microbes: A Review of the Interaction between Diet, Exercise, and the Gut Microbiota in Athletes", *Advances in Nutrition*, vol. 12, č. 6, s. 2190–2215, listopad 2021, doi: 10.1093/advances/nmab077.

[235] K. Deighton, R. Barry, C. E. Connon a D. J. Stensel, „Appetite, gut hormone and energy intake responses to low volume sprint interval and traditional endurance exercise", *Eur J Appl Physiol*, vol. 113, č. 5, s. 1147–1156, květen 2013, doi: 10.1007/s00421-012-2535-1.

[236] M. S. Farvid et al., „Dairy Food Intake and All-Cause, Cardiovascular Disease, and Cancer Mortality", *Am J Epidemiol*, vol. 185, č. 8, s. 697–711, duben 2017, doi: 10.1093/aje/kww139.

[237] V. Pala et al., „Yogurt consumption and risk of colorectal cancer in the Italian European prospective investigation into cancer and nutrition cohort", *Int J Cancer*, vol. 129, č. 11, s. 2712–2719, prosinec 2011, doi: 10.1002/ijc.26193.

[238] F. Magne et al., „The Firmicutes/Bacteroidetes Ratio: A Relevant Marker of Gut Dysbiosis in Obese Patients?", *Nutrients*, vol. 12, č. 5, s. 1474, květen 2020, doi: 10.3390/nu12051474.

[239] Y. Feng a D. Xu, „Short-chain fatty acids are potential goalkeepers of atherosclerosis", *Front Pharmacol*, vol. 14, listopad 2023, doi: 10.3389/fphar.2023.1271001.

[240] J. Scheiman et al., „Meta-omics analysis of elite athletes identifies a performance-enhancing microbe that functions via lactate metabolism", *Nat Med*, vol. 25, č. 7, s. 1104–1109, červenec 2019, doi: 10.1038/s41591-019-0485-4.

[241] R. J. Maughan et al., „IOC consensus statement: dietary supplements and the high-performance athlete", *Br J Sports Med*, vol. 52, č. 7, s. 439–455, duben 2018, doi: 10.1136/bjsports-2018-099027.

[242] R. Jäger et al., „International Society of Sports Nutrition Position Stand: Probiotics", *J Int Soc Sports Nutr*, vol. 16, č. 1, leden 2019, doi: 10.1186/s12970-019-0329-0.

[243] M. Aldunate et al., „Vaginal concentrations of lactic acid potently inactivate HIV", *Journal of Antimicrobial Chemotherapy*, vol. 68, č. 9, s. 2015–2025, září 2013, doi: 10.1093/jac/dkt156.

[244] R. F. Baggaley et al., „Heterosexual Anal Intercourse: A Neglected Risk Factor for HIV?", *American Journal of Reproductive Immunology*, vol. 69, č. s1, s. 95–105, únor 2013, doi: 10.1111/aji.12064.

[245] P. Łaniewski, K. A. Owen, M. Khnanisho, R. M. Brotman a M. M. Herbst-Kralovetz, „Clinical and Personal Lubricants Impact the Growth of Vaginal Lactobacillus Species and Colonization of Vaginal Epithelial Cells: An in Vitro Study", *Sex Transm Dis*, vol. 48, č. 1, s. 63–70, leden 2021, doi: 10.1097/OLQ.0000000000001272.

[246] M. Ristow et al., „Antioxidants prevent health-promoting effects of physical exercise in humans", *Proceedings of the National Academy of Sciences*, vol. 106, č. 21, s. 8665–8670, květen 2009, doi: 10.1073/pnas.0903485106.

[247] G. Bellinghieri, V. Savica a D. Santoro, „Renal Alterations During Exercise", *Journal of Renal Nutrition*, vol. 18, č. 1, s. 158–164, leden 2008, doi: 10.1053/j.jrn.2007.10.031.

[248] W. Kindermann, „Creatine Kinase Levels After Exercise“, *Dtsch Arztebl Int*, květen 2016, doi: 10.3238/arztebl.2016.0344a.

[249] E. Stenner, E. Gianoli, B. Biasioli, C. Piccinini, G. Delbello a A. Bussani, „Muscular damage and intravascular haemolysis during an 18 hour subterranean exploration in a cave of 700 m depth“, *Br J Sports Med*, vol. 40, č. 3, s. 235–238, březen 2006, doi: 10.1136/bjsm.2005.021402.

[250] E. Marques, R. Kramer a D. G. Ryan, „Multifaceted mitochondria in innate immunity“, *npj Metabolic Health and Disease*, vol. 2, č. 1, s. 6, květen 2024, doi: 10.1038/s44324-024-00008-3.

[251] S. Khan, S. Basu, D. Raj a A. Lahiri, „Role of mitochondria in regulating immune response during bacterial infection“, *International Review of Cell and Molecular Biology*, listopad 2023, s. 159–200. doi: 10.1016/bs.ircmb.2022.10.004.

[252] K. Witek, J. Ścisłowska, D. Turowski, K. Lerczak, S. Lewandowska-Pachecka a A. Pokrywka, „Total bilirubin in athletes, determination of reference range“, *Biol Sport*, vol. 1, s. 45–48, 2017, doi: 10.5114/biolsport.2017.63732.

[253] Maren. S. Fragala, C. Bi, M. Chaump, H. W. Kaufman a M. H. Kroll, „Associations of aerobic and strength exercise with clinical laboratory test values“, *PLOS One*, vol. 12, č. 10, s. e0180840, říjen 2017, doi: 10.1371/journal.pone.0180840.

[254] L. Vítek a C. Tiribelli, „Gilbert's syndrome revisited“, *J Hepatol*, vol. 79, č. 4, s. 1049–1055, říjen 2023, doi: 10.1016/j.jhep.2023.06.004.

[255] J. Woronyczová et al., „Serum Bilirubin Concentrations and the Prevalence of Gilbert Syndrome in Elite Athletes“, *Sports Med Open*, vol. 8, č. 1, s. 84, prosinec 2022, doi: 10.1186/s40798-022-00463-6.

[256] M. Cantinotti et al., „Cardiac Troponin-T Release After Sport and Differences by Age, Sex, Training Type, Volume, and Intensity: A Critical Review“, *Clinical Journal of Sport Medicine*, vol. 32, č. 3, s. e230–e242, květen 2022, doi: 10.1097/JSM.0000000000000940.

[257] C. Malm, J. Jakobsson a A. Isaksson, „Physical Activity and Sports—Real Health Benefits: A Review with Insight into the Public Health of Sweden“, *Sports*, vol. 7, č. 5, s. 127, květen 2019, doi: 10.3390/sports7050127.

[258] F. Sanchis-Gomar, G. Olaso-Gonzalez, D. Corella, M. C. Gomez-Cabrera a J. Vina, „Increased Average Longevity among the 'Tour de France' Cyclists“, *Int J Sports Med*, vol. 32, č. 08, s. 644–647, srpen 2011, doi: 10.1055/s-0031-1271711.

[259] B. Y. Farahmand et al., „Mortality amongst participants in Vasaloppet: a classical long-distance ski race in Sweden“, *J Intern Med*, vol. 253, č. 3, s. 276–283, březen 2003, doi: 10.1046/j.1365-2796.2003.01122.x.

[260] U. M. Kujala, „Essay: Does training adversely affect long-term health?“, *The Lancet*, vol. 366, s. S55–S56, prosinec 2005, doi: 10.1016/S0140-6736(05)67853-1.

[261] M. D. Bernat-Adell et al., „Recovery of Inflammation, Cardiac, and Muscle Damage Biomarkers After Running a Marathon“, *J Strength Cond Res*, vol. 35, č. 3, s. 626–632, březen 2021, doi: 10.1519/JSC.0000000000003167.

[262] N. R. Rodriguez, N. M. DiMarco a S. Langley, „Position of the American Dietetic Association, Dietitians of Canada, and the American College of Sports Medicine: Nutrition and Athletic Performance“, *J Am Diet Assoc*, vol. 109, č. 3, s. 509–527, březen 2009, doi: 10.1016/j.jada.2009.01.005.

[263] D. T. Thomas, L. M. Burke a K. A. Erdman, „Nutrition and Athletic Performance“, *Med Sci Sports Exerc*, vol. 48, č. 3, s. 543–568, březen 2016, doi: 10.1249/MSS.0000000000000852.

[264] C. Kerksick et al., „International Society of Sports Nutrition position stand: Nutrient timing“, *J Int Soc Sports Nutr*, vol. 5, č. 1, leden 2008, doi: 10.1186/1550-2783-5-17.

[265] M. Kanter, „High-Quality Carbohydrates and Physical Performance“, *Nutr Today*, vol. 53, č. 1, s. 35–39, leden 2018, doi: 10.1097/NT.0000000000000238.

[266] Y. Wang, Z. Liu, Y. Han, J. Xu, W. Huang a Z. Li, „Medium Chain Triglycerides enhances exercise endurance through the increased mitochondrial biogenesis and metabolism“, *PLOS One*, vol. 13, č. 2, s. e0191182, únor 2018, doi: 10.1371/journal.pone.0191182.

[267] T. J. Chapman-Lopez a Y. Koh, „The Effects of Medium-Chain Triglyceride Oil Supplementation on Endurance Performance and Substrate Utilization in Healthy Populations: A Systematic Review“, *J Obes Metab Syndr*, vol. 31, č. 3, s. 217–229, září 2022, doi: 10.7570/jomes22028.

[268] T. J. Chapman-Lopez a Y. Koh, „The Effects of Medium-Chain Triglyceride Oil Supplementation on Endurance Performance and Substrate Utilization in Healthy Populations: A Systematic Review“, *J Obes Metab Syndr*, vol. 31, č. 3, s. 217–229, září 2022, doi: 10.7570/jomes22028.

[269] N. S. Guest et al., „International society of sports nutrition position stand: caffeine and exercise performance“, *J Int Soc Sports Nutr*, vol. 18, č. 1, Jan. 2021, doi: 10.1186/s12970-020-00383-4.

[270] T. M. McLellan, J. A. Caldwell a H. R. Lieberman, „A review of caffeine's effects on cognitive, physical and occupational performance“, *Neurosci Biobehav Rev*, vol. 71, s. 294–312, prosinec 2016, doi: 10.1016/j.neubiorev.2016.09.001.

[271] T. E. Graham, „Caffeine and Exercise“, *Sports Medicine*, vol. 31, č. 11, s. 785–807, 2001, doi: 10.2165/00007256-200131110-00002.

[272] H. T. Bilondi, H. Valipour, S. Khoshro, P. Jamilian, A. Ostadrahimi a M. Zarezadeh, „The effect of caffeine supplementation on muscular strength and endurance: A meta-analysis of meta-analyses", *Heliyon*, vol. 10, č. 15, s. e35025, srpen 2024, doi: 10.1016/j.heliyon.2024.e35025.

[273] L. M. R. Loureiro, C. E. G. Reis a T. H. M. da Costa, „Effects of Coffee Components on Muscle Glycogen Recovery: A Systematic Review", *Int J Sport Nutr Exerc Metab*, vol. 28, č. 3, s. 284–293, květen 2018, doi: 10.1123/ijsnem.2017-0342.

[274] J. A. Muljadi, P. Kaewphongsri, K. Chaijenkij a J. Kongtharvonskul, „Effect of caffeine on delayed-onset muscle soreness: a meta-analysis of RCT", *Bull Natl Res Cent*, vol. 45, č. 1, s. 197, prosinec 2021, doi: 10.1186/s42269-021-00660-5.

[275] C. Pickering a J. Grgic, „Caffeine and Exercise: What Next?", *Sports Medicine*, vol. 49, č. 7, s. 1007–1030, červenec 2019, doi: 10.1007/s40279-019-01101-0.

[276] H. Turcotte, J.-B. Langdeau, G. Thibault a L.-P. Boulet, „Prevalence of respiratory symptoms in an athlete population", *Respir Med*, vol. 97, č. 8, s. 955–963, srpen 2003, doi: 10.1016/S0954-6111(03)00123-9.

[277] K. W. Rundell, J. Im, L. B. Mayers, R. L. Wilber, L. Szmedra a H. R. Schmitz, „Self-reported symptoms and exercise-induced asthma in the elite athlete", *Med Sci Sports Exerc*, s. 208–213, únor 2001, doi: 10.1097/00005768-200102000-00006.

[278] C. A. Jaworski a V. Rygiel, „Acute Illness in the Athlete", *Clin Sports Med*, vol. 38, č. 4, s. 577–595, říjen 2019, doi: 10.1016/j.csm.2019.05.001.

[279] D. C. Nieman a L. M. Wentz, „The compelling link between physical activity and the body's defense system", *J Sport Health Sci*, vol. 8, č. 3, s. 201–217, květen 2019, doi: 10.1016/j.jshs.2018.09.009.

[280] C. Malm, „Susceptibility to infections in elite athletes: the S-curve", *Scand J Med Sci Sports*, vol. 16, č. 1, s. 4–6, únor 2006, doi: 10.1111/j.1600-0838.2005.00499.x.

[281] E. M. Peters a E. D. Bateman, „Ultramarathon running and upper respiratory tract infections. An epidemiological survey", *S Afr Med J*, vol. 64, č. 15, s. 582–4, říjen 1983.

[282] D. C. Nieman, L. M. Johanssen, J. W. Lee a K. Arabatzis, „Infectious episodes in runners before and after the Los Angeles Marathon", *J Sports Med Phys Fitness*, vol. 30, č. 3, s. 316–28, září 1990.

[283] D. König, D. Grathwohl, C. Weinstock, H. Northoff a A. Berg, „Upper respiratory tract infection in athletes: influence of lifestyle, type of sport, training effort, and immunostimulant intake", *Exerc Immunol Rev*, vol. 6, s. 102–20, 2000.

[284] A. Van Tonder, M. Schwellnus, S. Swanevelder, E. Jordaan, W. Derman a D. C. Janse van Rensburg, „A prospective cohort study of 7031 distance runners shows that 1 in 13 report systemic symptoms of an acute illness in the 8–12 day period before a race, increasing their risk of not finishing the race 1.9 times for those runners who started the race: SAFER study IV", *Br J Sports Med*, vol. 50, č. 15, s. 939–945, srpen 2016, doi: 10.1136/bjsports-2016-096190.

[285] L. SPENCE et al., „Incidence, Etiology, and Symptomatology of Upper Respiratory Illness in Elite Athletes", *Med Sci Sports Exerc*, vol. 39, č. 4, s. 577–586, duben 2007, doi: 10.1249/mss.0b013e31802e851a.

[286] L. Rama et al., „Changes in natural killer cell subpopulations over a winter training season in elite swimmers", *Eur J Appl Physiol*, vol. 113, č. 4, s. 859–868, duben 2013, doi: 10.1007/s00421-012-2490-x.

[287] I. S. Svendsen, M. Gleeson, T. A. Haugen a E. Tønnessen, „Effect of an intense period of competition on race performance and self-reported illness in elite cross-country skiers", *Scand J Med Sci Sports*, vol. 25, č. 6, s. 846–853, prosinec 2015, doi: 10.1111/sms.12452.

[288] M. Drew et al., „Prevalence of illness, poor mental health and sleep quality and low energy availability prior to the 2016 Summer Olympic Games", *Br J Sports Med*, vol. 52, č. 1, s. 47–53, leden 2018, doi: 10.1136/bjsports-2017-098208.

[289] T. Timpka et al., „Preparticipation predictors for championship injury and illness: cohort study at the Beijing 2015 International Association of Athletics Federations World Championships", *Br J Sports Med*, vol. 51, č. 4, s. 271–276, únor 2017, doi: 10.1136/bjsports-2016-096580.

[290] R. Baskerville, L. Castell a S. Bermon, „Sports and Immunity, from the recreational to the elite athlete", *Infect Dis Now*, vol. 54, č. 4, s. 104893, červen 2024, doi: 10.1016/j.idnow.2024.104893.

[291] Y. Park, A. F. Subar, A. Hollenbeck a A. Schatzkin, „Dietary Fiber Intake and Mortality in the NIH-AARP Diet and Health Study", *Arch Intern Med*, vol. 171, č. 12, červen 2011, doi: 10.1001/archinternmed.2011.18.

[292] D. R. Axon, J. Vanova, C. Edel a M. Slack, „Dietary Supplement Use, Knowledge, and Perceptions Among Student Pharmacists", *Am J Pharm Educ*, vol. 81, č. 5, s. 92, červen 2017, doi: 10.5688/ajpe81592.

[293] D. K. Arnett et al., „2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines", *Circulation*, vol. 140, č. 11, září 2019, doi: 10.1161/CIR.0000000000000678.

[294] D. Aune et al., „Fruit and vegetable intake and the risk of cardiovascular disease, total cancer and all-cause mortality—a systematic review and

dose-response meta-analysis of prospective studies", *Int J Epidemiol*, vol. 46, č. 3, s. 1029–1056, červen 2017, doi: 10.1093/ije/dyw319.

[295] E. A. O'Connor et al., „Vitamin and Mineral Supplements for the Primary Prevention of Cardiovascular Disease and Cancer", *JAMA*, vol. 327, č. 23, s. 2334, červen 2022, doi: 10.1001/jama.2021.15650.

[296] P. Peeling, M. Sim a A. K. A. McKay, „Considerations for the Consumption of Vitamin and Mineral Supplements in Athlete Populations", *Sports Medicine*, vol. 53, č. S1, s. 15–24, prosinec 2023, doi: 10.1007/s40279-023-01875-4.

[297] F. Farrokhyar et al., „Effects of Vitamin D Supplementation on Serum 25-Hydroxyvitamin D Concentrations and Physical Performance in Athletes: A Systematic Review and Meta-analysis of Randomized Controlled Trials", *Sports Medicine*, vol. 47, č. 11, s. 2323–2339, listopad 2017, doi: 10.1007/s40279-017-0749-4.

[298] D. Holmannova et al., „High prevalence of low vitamin D status in the Czech Republic: a retrospective study of 119,925 participants", *Eur J Clin Nutr*, březen 2025, doi: 10.1038/s41430-025-01587-0.

[299] J. F. Creeden, D. M. Gordon, D. E. Stec a T. D. Hinds, „Bilirubin as a metabolic hormone: the physiological relevance of low levels", *American Journal of Physiology-Endocrinology and Metabolism*, vol. 320, č. 2, s. E191–E207, únor 2021, doi: 10.1152/ajpendo.00405.2020.

[300] M. B. Sintzel, M. Rametta a A. T. Reder, „Vitamin D and Multiple Sclerosis: A Comprehensive Review", *Neurol Ther*, vol. 7, č. 1, s. 59–85, červen 2018, doi: 10.1007/s40120-017-0086-4.

[301] T. M. Abugoukh et al., „Does Vitamin D Have a Role in Diabetes?", *Cureus*, říjen 2022, doi: 10.7759/cureus.30432.

[302] T. Podlogar, N. Cooper-Smith, J. T. Gonzalez a G. A. Wallis, „Personalised carbohydrate feeding during exercise based on exogenous glucose oxidation: a proof-of-concept study", *Performance Nutrition*, vol. 1, č. 1, s. 2, duben 2025, doi: 10.1186/s44410-025-00003-9.

[303] A. E. Jeukendrup, L. Moseley, G. I. Mainwaring, S. Samuels, S. Perry a C. H. Mann, „Exogenous carbohydrate oxidation during ultraendurance exercise", *J Appl Physiol*, vol. 100, č. 4, s. 1134–1141, duben 2006, doi: 10.1152/japplphysiol.00981.2004.

[304] R. A. Fielding, D. L. Costill, W. J. Fink, D. S. King, M. Hargreaves a J. E. Kovaleski, „Effect of carbohydrate feeding frequencies and dosage on muscle glycogen use during exercise", *Med Sci Sports Exerc*, vol. 17, č. 4, s. 472–476, srpen 1985, doi: 10.1249/00005768-198508000-00012.

[305] A. Jeukendrup, „A Step Towards Personalized Sports Nutrition: Carbohydrate Intake During Exercise", *Sports Medicine*, vol. 44, č. S1, s. 25–33, květen 2014, doi: 10.1007/s40279-014-0148-z.

[306] K. J. Kolnes et al., „Effects of seven days' fasting on physical performance and metabolic adaptation during exercise in humans", *Nat Commun*, vol. 16, č. 1, s. 122, leden 2025, doi: 10.1038/s41467-024-55418-0.

[307] J. Conde-Pipó et al., „Intermittent Fasting: Does It Affect Sports Performance? A Systematic Review", *Nutrients*, vol. 16, č. 1, s. 168, leden 2024, doi: 10.3390/nu16010168.

[308] H. Zouhal et al., „Exercise Training and Fasting: Current Insights", *Open Access J Sports Med*, vol. 11, s. 1–28, leden 2020, doi: 10.2147/OAJSM.S224919.

[309] L. Bi a G. Triadafilopoulos, „Exercise and gastrointestinal function and disease: an evidence-based review of risks and benefits", *Clinical Gastroenterology and Hepatology*, vol. 1, č. 5, s. 345–355, září 2003, doi: 10.1053/S1542-3565(03)00178-2.

[310] H. Jarrett, S. Medlin a J. Morehen, „The Role of the Gut Microbiome and Probiotics in Sports Performance: A Narrative Review Update", *Nutrients*, vol. 17, č. 4, s. 690, Feb. 2025, doi: 10.3390/nu17040690.

[311] L. Mancin et al., „Standardization of gut microbiome analysis in sports", *Cell Rep Med*, vol. 5, č. 10, s. 101759, říjen 2024, doi: 10.1016/j.xcrm.2024.101759.

[312] A. Pradhan, A. Srinivasan, S. Raj, D. Howard a R. S. Aujla, „A systematic review and meta-analysis of preoperative carbohydrate drink loading prior to elective hip and knee arthroplasty", *J Clin Orthop Trauma*, vol. 63, s. 102928, duben 2025, doi: 10.1016/j.jcot.2025.102928.

[313] C. Ricci et al., „Preoperative carbohydrate loading before elective abdominal surgery: A systematic review and network meta-analysis of phase II/III randomized controlled trials", *Clinical Nutrition*, vol. 41, č. 2, s. 313–320, únor 2022, doi: 10.1016/j.clnu.2021.12.016.

[314] M. Pöchmüller, L. Schwingshackl, P. C. Colombani a G. Hoffmann, „A systematic review and meta-analysis of carbohydrate benefits associated with randomized controlled competition-based performance trials", *J Int Soc Sports Nutr*, vol. 13, č. 1, leden 2016, doi: 10.1186/s12970-016-0139-6.

[315] L. M. Burke, J. A. Hawley, E. J. Schabort, A. S. C. Gibson, I. Mujika a T. D. Noakes, „Carbohydrate loading failed to improve 100-km cycling performance in a placebo-controlled trial", *J Appl Physiol*, vol. 88, č. 4, s. 1284–1290, duben 2000, doi: 10.1152/jappl.2000.88.4.1284.

[316] L. M. Burke et al., „Adaptations to short-term high-fat diet persist during exercise despite high carbohydrate availability", *Med Sci Sports Exerc*, vol. 34, č. 1, s. 83–91, leden 2002, doi: 10.1097/00005768-200201000-00014.

[317] A. Kazemi, G. Racil, A. H. Ahmadi Hekmatikar, M. Behnam Moghadam, P. Karami a M. Henselmans, „Improved physical performance of elite soccer players based on GPS results after 4 days of carbohydrate loading followed by 3 days of low carbohydrate diet", *J Int Soc Sports Nutr*, vol. 20, č. 1, prosinec 2023, doi: 10.1080/15502783.2023.2258837.

[318] L. M. BURKE et al., „Adaptations to short-term high-fat diet persist during exercise despite high carbohydrate availability", *Med Sci Sports Exerc*, vol. 34, č. 1, s. 83–91, leden 2002, doi: 10.1097/00005768-200201000-00014.

[319] T. D. Noakes, „What Is the Evidence That Dietary Macronutrient Composition Influences Exercise Performance? A Narrative Review", *Nutrients*, vol. 14, č. 4, s. 862, Feb. 2022, doi: 10.3390/nu14040862.

[320] J. Karlsson a B. Saltin, „Diet, muscle glycogen, and endurance performance", *J Appl Physiol*, vol. 31, č. 2, s. 203–206, srpen 1971, doi: 10.1152/jappl.1971.31.2.203.

[321] M. Henselmans, T. Bjørnsen, R. Hedderman, and F. T. Vårvik, „The Effect of Carbohydrate Intake on Strength and Resistance Training Performance: A Systematic Review", *Nutrients*, vol. 14, č. 4, s. 856, únor 2022, doi: 10.3390/nu14040856.

[322] S. J. M. Osendarp, K. H. Brown, L. M. Neufeld, E. Udomkesmalee a S. E. Moore, „The double burden of malnutrition—further perspective", *The Lancet*, vol. 396, č. 10254, s. 813, září 2020, doi: 10.1016/S0140-6736(20)31364-7.

[323] T. Gödecke, A. J. Stein a M. Qaim, „The global burden of chronic and hidden hunger: Trends and determinants", *Glob Food Sec*, vol. 17, s. 21–29, červen 2018, doi: 10.1016/j.gfs.2018.03.004.

[324] A. Kraus, L. Zita, O. Krunt, H. Härtlová a E. Chmelíková, „Determination of selected biochemical parameters in blood serum and egg quality of Czech and Slovak native hens depending on the housing system and hen age", *Poult Sci*, vol. 100, č. 2, s. 1142–1153, únor 2021, doi: 10.1016/j.psj.2020.10.039.

[325] C. Shekhar, „Hidden Hunger: Addressing Micronutrient Deficiencies Using Improved Crop Varieties", *Chem Biol*, vol. 20, č. 11, s. 1305–1306, listopad 2013, doi: 10.1016/j.chembiol.2013.11.003.

[326] G. B. M. Mensink et al., „Mapping low intake of micronutrients across Europe", *British Journal of Nutrition*, vol. 110, č. 4, s. 755–773, srpen 2013, doi: 10.1017/S000711451200565X.

[327] K. D. Stark, M. E. Van Elswyk, M. R. Higgins, C. A. Weatherford a N. Salem, „Global survey of the omega-3 fatty acids, docosahexaenoic acid and eicosapentaenoic acid in the blood stream of healthy adults", *Prog*

Lipid Res, vol. 63, s. 132–152, červenec 2016, doi: 10.1016/j.plipres.2016.05.001.

[328] K. Albracht-Schulte et al., „Omega-3 fatty acids in obesity and metabolic syndrome: a mechanistic update“, *J Nutr Biochem*, vol. 58, s. 1–16, srpen 2018, doi: 10.1016/j.jnutbio.2018.02.012.

[329] V. Šmíd et al., „Effect of Omega-3 Polyunsaturated Fatty Acids on Lipid Metabolism in Patients With Metabolic Syndrome and NAFLD“, *Hepatol Commun*, vol. 6, č. 6, s. 1336–1349, červen 2022, doi: 10.1002/hep4.1906.

[330] M. Bishehkolaei a Y. Pathak, „Influence of omega n-6/n-3 ratio on cardiovascular disease and nutritional interventions“, *Human Nutrition & Metabolism*, vol. 37, s. 200275, září 2024, doi: 10.1016/j.hnm.2024.200275.

[331] J. M. Hands, M. L. Anderson, T. Cooperman a L. A. Frame, „A Multi-Year Rancidity Analysis of 72 Marine and Microalgal Oil Omega-3 Supplements“, *J Diet Suppl*, vol. 21, č. 2, s. 195–206, březen 2024, doi: 10.1080/19390211.2023.2252064.

[332] Z. Veprikova et al., „Mycotoxins in Plant-Based Dietary Supplements: Hidden Health Risk for Consumers“, *J Agric Food Chem*, vol. 63, č. 29, s. 6633–6643, červenec 2015, doi: 10.1021/acs.jafc.5b02105.

[333] X. Zhang, L. Zhang, T. Zhou a Y. Zhou, „Fungal flora and mycotoxin contamination in tea: Current status, detection methods and dietary risk assessment – A comprehensive review“, *Trends Food Sci Technol*, vol. 127, s. 207–220, září 2022, doi: 10.1016/j.tifs.2022.05.019.

[334] J. H. Lee, J. H. Han, A. Y. Min, H. Kim a D. Shin, „Screening for twenty-eight target anabolic-androgenic steroids in protein supplements using QuEChERS extraction followed by liquid chromatography-tandem mass spektrometry“, *Food Additives & Contaminants: Part A*, vol. 37, č. 9, s. 1425–1436, září 2020, doi: 10.1080/19440049.2020.1773543.

[335] S. B. Bandara, K. M. Towle a A. D. Monnot, „A human health risk assessment of heavy metal ingestion among consumers of protein powder supplements“, *Toxicol Rep*, vol. 7, s. 1255–1262, 2020, doi: 10.1016/j.toxrep.2020.08.001.

[336] D. Fitzpatrick, T. Leckie, G. Heine a L. Hodgson, „The use of pain killers (NSAIDs) in athletes: How large is the risk?“, *J Sci Med Sport*, vol. 28, č. 3, s. 198–205, březen 2025, doi: 10.1016/j.jsams.2024.11.010.

[337] C. Cornu et al., „Effect of Non-Steroidal Anti-Inflammatory Drugs on Sport Performance Indices in Healthy People: a Meta-Analysis of Randomized Controlled Trials“, *Sports Med Open*, vol. 6, č. 1, s. 20, prosinec 2020, doi: 10.1186/s40798-020-00247-w.

[338] R. L. Nahon, J. S. S. Lopes, A. M. de Magalhães Neto, A. de S. Machado a L. C. Cameron, „Anti-Inflammatories for Delayed Onset Muscle

Soreness: Systematic Review and Meta-Analysis", *Revista Brasileira de Medicina do Esporte*, vol. 27, č. 6, s. 646–654, prosinec 2021, doi: 10.1590/1517-8692202127062021_0072.

[339] G. S. Lipman et al., „Ibuprofen versus placebo effect on acute kidney injury in ultramarathons: a randomised controlled trial", *Emergency Medicine Journal*, vol. 34, č. 10, s. 637–642, říjen 2017, doi: 10.1136/emermed-2016-206353.

[340] B. V. Scheer a E. Valero Burgos, „The Hidden Danger of Endurance Races: Analgesic Use among Ultramarathon Runners", *Br J Sports Med*, vol. 47, č. 10, s. e3.13–e3, červenec 2013, doi: 10.1136/bjsports-2013-092558.20.