

17 Literatura a prohlubující literatura

- Akrali O**, Tiner M, Ozaksoy D: Effects of lower extremity on prognosis of flexible flatfoot in children. *Foot Ankle Int* 2000; č. 21(9): 772-774.
- Alexander RM**: The Human Machine. Natural History Museum Publication, London 1992; a) 74-87.
- Ali-Gombe A**, Croft PR, Silman AJ: Osteoarthritis of the hip and acetabular dysplasia in Nigerian men. *J Rheumatol* 1996; 23(3): 512-515.
- Allison SC**, Abraham LD: Sensitivity of qualitative and quantitative spasticity measures to clinical treatment with cryotherapy. *Int J Rehabil Res* 2001; 24(1): 15-24.
- Allgöwer M**: Allgemeine und spezielle Chirurgie. Springer 1976: a) 284.
- Anderson FA Jr**, Wheeler HB, Goldberg RJ, Hosmer DW, Patwardhan NA, Jovanović B, Forcier A, Dalen JE: A population-based perspective of the hospital incidence and case-fatality rates of deep vein thrombosis and pulmonary embolism. The Worcester DVT Study. *Arch Intern Med* 1991; 151(5): 933-8.
- Anderson ST**, Charlesworth RW: Rheumatic disease among Air force recruits: a multimillions dollar epidemic. *Semin Arthritis Rheum* 1993; 22(4): 275-9.
- Antonicelli R**, Sardina M, Scotti A, Bonizzoni E, Paciaroni E: Randomized trial of the effects of low-dose calcium-heparin in patients with peripheral arterial disease and claudication. Italian CAP Study Group. *Am J Med* 1999; 107(3): 234-239.
- Attia J**, Ray JG, Cook DJ, Douketis J, Ginsberg JS, Geerts WH: Deep vein thrombosis and its prevention in critically ill adults. *Arch Intern Med* 2001; 161(10): 1268-79.
- Auerswald W**: Ganong Lehrbuch der medizinischen Physiologie. Springer 1974(3): a) 514-16; b) 521-23.
- Bazner H**, Oster M, Daffertshoffer M, Hennerici M: Assessment of gait in subcortical vascular encephalopathy by computerized analysis: a cross-sectional and longitudinal study. *J Neurol* 2000; 247(11): 841-49.
- Beck W**, citován v: Die Füße sind die Dummen. *Bunte* 2000; 35: 78-81.
- Belchamber TL**, van den Bogert AJ: Contribution of proximal and distal moments to axial tibia rotation during walking and running. *J Biomech* 2000; 33(11): 1397-1403.
- Bencardino J**, Rosenberg ZS, Bertran J, Liu X, Marty-Delfaut E: Morton's neuroma: is it always symptomatic? *AJR Am J Roentgenol* 2000; 175(3): 649-53.
- Benninghoff A**, vybral z: Anatomie – Band 1 Bewegungsapparat. Urban & Schwarzenberg 1985, München. A) 198 obr. 3.3-10; b) 338-39.
- Benson ER**, Schutzer SF: Posttraumatic piriformis syndrome: diagnosis and results of operative treatment. *J Bone Joint Surg Am* 1999; 81(7): 941-49.
- Bergmann C**, Kniggeordorf H, Graichen F, Rohlmann A: Influence of shoes and heel strike on the loading of the hip joint. *J Biomech* 1995; 28(7): 817-27.
- Biasca N**, Zanetti M, Zollinger H: Outcomes after partial neurectomy of Morton's neuroma related to preoperative case histories, clinical findings, and findings of magnetic resonance imaging scans. *Foot Ankle Int* 1999; 20(9): 568-75.
- Birbamer G**: Das Gehirn denkt nicht in Muskelkontraktionen; Gespräch über kognitiv-therapeutische Übungen nach Perfetti, Teil II. *Ars Medici* 2001; 7: 340-42.
- Bonnel F**, Canovas F, Poiree G, Duserre F, Vergnes C: Evaluation of the Scarf osteotomy in hallux valgus related to distal metatarsal articular angle: a prospective study of 79 operated cases. *Rev Chir Orthop Reparatrice Appar Mot* 1999; 85(4): 381-86.
- Bradbury A**, Evans C, Allan P, Lee A, Ruckley CV, Fowkes FG: What are the symptoms of varicose veins? Edinburgh vein study cross sectional population survey. *BMJ* 1999; 318(7180): 353-56.
- Brandt KG**, Heiman DK, Slemenda C, Katz BP, Mazuca S, Braunstein EM, Byrd D: A comparison of lower extremity muscle strength, obesity, and depression scores in elderly subjects with knee pain with and without radiographic evidence of knee osteoarthritis. *J Rheumatol* 2000; 27(8): 1937-46.

- Broderston** A, Pederson B, Reimers J: Foot deformities and relation to the length of leg muscles in Danish children aged 3-17 years. *Ugeskr Laeger* 1993; 155(48): 3914-16.
- Bruckner** L, Rosler P: Orthopaedic findings of the foot as related to age and body weight among women. *ZFA* 1981; 36(1): 29-35.
- Camasta** CA: Hallux limitus and hallux rigidus. Clinical examination, radiographic findings, and natural history. *Clin Podiatr Med Surg* 1996; 13(3): 423-48.
- Capitão** LM, Menezes JD, Gouveia-Oliveira A: The epidemiology of chronic venous insufficiency in Portugal. *Acta Med Port* 1995; 8(9): 485-91.
- Carmelli** D, Kelly-Hayes M, Wolf PA, Swan GE, Jack LM, Reed t, Guralnik JM: The contribution of genetic influences to measures of lower-extremity function in older male twins. *J Gerontol A Biol Sci Med Sci* 2000; 55(1): B49-53.
- Chalmers** N: Man's place in evolution. *Natural History Museum Publications*, Cambridge University Press 1991; a) 44-45; b) 53; c) 96.
- Chappuis** C: Ein geriatrisches Zentrum – sein Beitrag zur Bearbeitung der Problemliste des Hausarztes am Beispiel des Problems „Sturz“. *Moderne Geriatrie* 1994: 118-124.
- Cheng** SW, Ting AC, Lau H, Wong J: Epidemiology of atherosclerotic peripheral arterial occlusive disease in Hong Kong. *World J Surg* 1999; 23(2): 202-206.
- Cicchinelli** LD, Camasta CA, McGlamry ED: Iatrogenic metatarsus primus elevatus. Etiology, evaluation, and surgical management. *J Am Podiatr Med Assoc* 1997; 87(4): 165-77.
- Coleman** KJ, Raynor HR, Mueller DM, Černý FJ, Dorn JM, Epstein LH: Providing sedentary adults with choices for meeting their walking goals. *Prev Med Assoc* 1999; 28(5): 509-10.
- Connolly** KJ: Mechanisms of Motor Skill Development. Academic Press London, 1970: 161-88.
- Connolly** KJ, Manoel EJ: Variability and stability in the development of skilled actions; v: Neurophysiology & Neuropsychology of Motor Development. Mac Keith Press, Cambridge, 1977: 286-318; a) 325-27.
- Cooper** KH: Bewegungstraining („Aerobics“). Fischer Bücherei, Frankfurt 1970.
- Coughlin** MJ: Roger A. Mann Award. Juvenile hallux valgus: aetiology and treatment. *Foot Ankle Int* 1995; 16(11): 682-97.
- Coupland** CA, Cliffe CJ, Bassey EJ, Grainge MJ, Hosking DJ, Chilvers CE: Habitual physical activity and bone mineral density in post-menopausal women in England. *Int J Epidemiol* 1999; 28(2): 241-46.
- Coutts** F: Gait analysis in the therapeutic environment. *Man Ther* 1999; 4(1): 2-10.
- Craik** KJW: Theory of the human operator in control systems. I. The operator as an engineering system. *British Journal of Psychology*. 1947; (38): 56-61.
- Cummings** SR, Ling X, Stone K: Consequences of foot binding among older women in Beijing, China. *American Journal Public Health*. 1997; 87(10): 1677-79.
- Dahlke** R: Krankheit als Symbol – Handbuch der Psychosomatik. Bertelsmann; 1996 (3. vydání): 14-17.
- Davis** JW, Nevitt MC, Wasnich RD, Ross PD: A cross-cultural comparison of neuromuscular performance, functional status, and falls between Japanese and white women. *Journal Gerontol a Biol Sci Med Sci* 1999; 54(6): M288-92.
- Debrunner** HU: Biomechanik des Fußes. Bücherei des Orthopäden, Band 49, Enke; 1985: a) 7; b) 76.
- Didia** BC, Omu ET, Obvuoforibor AA: The use of footprint contact index II for classification of flat feet in Nigerian population. *Foot Ankle*. 1987; 7(5): 285-89.
- Diener** M: Zeigt her Eure Schuhe! Jeder Laufschuh hat seine spezifischen Eigenschaften. *K-Tip* 2000: 16: 35-37.
- Dücker** M: Antiker Sport und Olympische Spiele. Pädagogischer Dienst der Staatlichen Museen Preussischer Kulturbesitz. 1981, Berlin: Umschlag Stadionlauf der Männer, Panathenäische Preisamphore z obdobi okolo roku 480 před našim letopočtem.
- Dujardin** F, Aucouturier T, Bocquet G, Duparc F, Weber J, Thomine JM: Kinematics of the healthy and arthritic hip joint during walking. A study of 136 subjects. *Rev Chir Orthop Reparatrice Appar Mot* 1998; 84(8): 689-99.

- Duncan** JJ, Gordon NF, Scott DJ: Women walking for health and fitness. How much is enough? *JAMA* 1991; 266(23): 3 295-99.
- Durkin** MT, Turton EP, Wijesinghe LD, Scott DJ, Berridge DC: Long saphenous vein stripping and quality of life – a randomized trial. *Eur J Vasc Endovasc Surg* 2001; 21(6): 545-49.
- Dwyer** FC: The present status of the problem of pes cavus. *Clinical Orthopaedic* 1975; (106): 254-275.
- Eastwod** DM, Menelaus MB, Dickens DR, Broughton NS, Cole WG: Idiopathic toe-walking: does treatment alter the natural history? *Journal Pediatr Orthopaedic* 2000; 9(1): 47-49.
- Einarsdottir** H, Troell S, Wykman A: Hallux valgus in ballet dancers: a myth? *Foot ankle Int* 1995; 16(2): 92-94.
- Evans** CJ, Fowkes FG, Ruckley CV, Lee AJ: Prevalence of varicose veins and chronic venous insufficiency in men and women in the general population: Edinburgh vein study. *J Epidemiol Community Health* 1999; 53(3): 149-153.
- Feneis** H: *Anatomisches Bilderwörterbuch*. Thieme, 1974.
- Ferrell** BA, Josephson KR, Pollan AM, Loy S, Ferrell BR: A randomized trial of walking versus physical methods for chronic pain management. *Aging (Milano)* 1997; 9(1-2): 99-105.
- Fowkes** BA, Josephson KR, Pollan AM, Loy S, Ferrell BR: A randomized trial of walking versus physical methods for chronic pain management. *Aging (Milano)* 1997; 9(1-2): 99-105.
- Friden** T, Erlandsson T, Zetterstrom R, Lindstrand A, Moritz U: Compression or distraction of the anterior cruciate injured knee. Variations in injury pattern in contact sports and downhill skiing. *Knee Surg Sports Traumatologic Arthroscopy* 1995; 3(3): 144-147.
- Fritz** GR, Prieskorn D: First metatarsocuneiform motion: a radiographic and statistical analysis. *Foot Ankle Int* 1995; 16(3): 117-123.
- Fuchs** R, Staheli LT: Sprinting and intoeing. *J Pediatr Orthop* 1996; 16(4): 489-91.
- Fuller** B: *Synergetics*. Macmillan, New York, 1982. 382-430 (405).
- Garcia-Rodrigues** A, Martin Jimenez F: Flexible flat feet in children: a real problem? *Pediatrics* 1999; 103(6): e84.
- Gerhardt** JJ, Rippstein JR: Gelenk und Bewegung. Huber, 1992, Bern. a) 110-111; b) 100-101; c).
- Gesell** A: *The Embryology of Behaviour*. 1946. Přetištěno pod názvem: *Classics in Developmental Medicine* No. 3 Mac Keith, London, 1988.
- Gesell** A: Maturation and infant behaviour pattern. *Psychological review*, 1929; (36): 307-319.
- Gibellini** R, Fanello M, Bardile AF, Salerno M, Alois T: Exercise training in intermittent claudication. *Int Angiol* 2000; 19(1): 8-13.
- Godlee** F: *Clinical Evidence – die besten Studien für die klinische Praxis*, Ausgabe 2000. Hand Huber, 2000: a) 271-272; b) 302-303.
- Gonzales** de Aledo Linos A, Rollan A, Bonilla Miera C, Montes Conde A, Diego Santamaria MC, Obeso García M: Results of podoscope screening in 948 non-selected children, with special reference to cavus foot. *An Esp Pediatr* 1996; 45(6): 579-82.
- Gottschalk** FA, Solomon L, Beighton PH: The prevalence of hallux valgus in South African males. *South Africa Medicine Journal* 1984; 65(18): 725-26.
- Gould** N, Schneider W, Ashikaga T: Epidemiological survey of foot problems in the continental United States: 1978-1979. *Foot Ankle* 1980; 1(1): 8-11.
- Goutallier** D, Garabedian JM, Allain J, Bernageau J: Effect of osseous torsions of the lower limb on the development of lateral femorotibial knee arthrosis. *Rev Chir Orthop Reparatrice Appar Mot* 1997; 83(7): 613-621.
- Grossner** M, Neumaier A: *Technik-Trainink*. BLV Sportwissen. 1982: 9-10.
- Gunther** KP, Sturmer T, Sauerland S, Zeissig I, Sun Y, Kessler S, Scharf HP, Brenner H, Puhl W: Prevalence of generalized osteoarthritis in patients with advanced hip and knee osteoarthritis: the Ulm Osteoarthritis Study. *Ann Rheum Dis* 1998; 57(12): 717-723.
- Gutzwiller** F: *Savoir vivre – Gesundheit ist beeinflussbar*, by PAX, Basel, 2000.
- Guyton** GP, Mann RA, Krieger LE, Mendel T, Kahan J: Cumulative industrial trauma as an aetiology of seven common disorders in the foot and ankle: what is the evidence? *Foot Ankle Int* 2000; 21(12): 1047-1056.

- Hafner J:** Conservative therapy in varicose symptom complex. *Schweiz Rundsch Med Prax* 2001; 90(6): 197-204.
- Hamdorf PA, Penhall RK:** Walking with its training effects on the fitness and activity patterns of 79-91 year old females. *Aust N Z J Med* 1999; 29(1): 22-28.
- Haskell WL, Alderman EL, Fair JM, Maron DJ, Mackey SF, Superko HR, Williams PT, Johnstone IM, Champagne MA, Krauss RM:** Effects of intensive multiple risk factor reduction on coronary atherosclerosis and clinical cardiac events in men and women with coronary artery disease. The Stanford Coronary Risk Intervention Project (SCRIP). *Circulation* 1994; 89(3): 975-990.
- Hawker GA, Wright JG, Coyte PC, Williams JI, Harvey B, Glazier R, Badley EM:** Differences between men and women in the rate of use of hip and knee arthroplasty. *N Engl J Med* 2000; 342(14): 1016-22.
- Hawker GA, Wright JG, Coyte PC, Williams JI, Harvey B, Glazier R, Wilkins A, Badley EM:** Determining the need for hip and knee arthroplasty: the role of clinical severity and patients' preferences. *Med Care* 2001; 39(3): 06-16.
- Hayafune N, Hayafune H, Jacob AC:** Pressure and force distribution characteristics under the normal foot during the push-off phase in gait. *The Foot* 1999 (9): 88-92.
- Horton GA, Park YW, Myerson MS:** Role of metatarsus primus elevatus in the pathogenesis of hallux rigidus. *Foot Ankle Int* 1999; 20(12): 777-780.
- Helliwell TR, Tynan M, Hayward M, Klenerman L, Whitehouse G, Edwards RH:** The pathology of the lower leg muscles in pure forefoot pes cavus. *Acta Neuropathol (Berl)* 1995; 89(6): 552-559.
- Herzog G:** Der diabetische Fuß. *Ars Medici* 2001; 15: 448-451 a 17: 816-817.
- Holden MK, Gill KM, Maglioni MR, Nathan J, Piehl-Barker L:** Clinical gait assessment in the neurologically impaired. Reliability and meaningfulness. *Phys Ther* 1984; 64(1): 35-40.
- Hooi JD, Stoffers HE, Kester AD, Rinkens PR, Kaiser V, van Ree JW, Knottnerus JA:** Risk factors and cardiovascular diseases associated with asymptomatic peripheral arterial occlusive disease. The Limburg PAOD Study. *Peripheral arterial occlusive disease. Scand J Prim Health Care* 1998; 16(3): 177-182.
- Hu FB, Manson JE, Stampfer MJ, Colditz G, Liu S, Solomon CG, Willett WC:** Diet, lifestyle, and the risk of type 2 diabetes mellitus in women. *N Engl J Med* 2001; 345(11): 790-797.
- Hulet C, Hurwitz DE, Andriacchi TP, Galante JO, Vielpeau C:** Function gait adaptations in patients with painful hip. *Rev Chir Orthop Reparatrice Appar Mot* 2000; 86(6): 581-589.
- Hullin MG, Robb JE, Loudon JR:** Gait patterns in children with hemiplegic spastic cerebral palsy. *J Pediatr Orthop B* 1996; 5(4): 247-251.
- Hung LK, Ho YF, Leung PC:** Survey of foot deformities among 166 geriatric inpatients. *Foot Ankle* 1985; 5(4): 156-164.
- Huwylar J:** Der Tänzer und sein Körper; Aspekte des Tanzes aus ärztlicher Sicht. *Perimed, Erlangen*, 1992; a) 125-139; b) 60-61; c) 134.
- Ingvarsson T, Hagglund G, Lohmander LS:** Prevalence of hip osteoarthritis in Iceland. *Ann Rheum Dis* 1999; 58(4): 201-207.
- Ito H, Shimizu A, Miyamoto T, Katsura Y, Tanaka K:** Clinical significance of increased mobility in the sagittal plane in patients with hallux valgus. *Foot Ankle Int* 1999; 20(1): 29-32.
- Jacob HA:** Zur Belastung des Fußes beim Gehen und Stehen – Hat die „Dreipunktetheorie“ noch Bestand? *Deutsche Assoziation für orthopädische Fußchirurgie. D.A.F News* 2000; 6(2): 4-10.
- Jacobi G, Riepert T, Kieslich M, Bohl J:** Todesfall während der Physiotherapie nach Vojta. *Z. f. Physiotherapeuten* 2001; 53(4): 573-576.
- Jerosch J, Mamsch H:** Deformities and misalignment of feet in children – a field study of 345 students. *Z Orthop Ihre Grenzgeb* 1998; 136(3): 215-220.
- Joseph B, Chacko V, Abraham T, Jacob M:** Pathomechanics of congenital and acquired hallux varus: a clinical and anatomical study. *Foot Ankle* 1987; 8(3): 137-143.
- Kahn CR, Solymoss S, Lamping DL, Abenhaim L:** Long-term outcomes after deep vein thrombosis: postphlebotic syndrome and quality of life. *J Gen Intern Med* 2000; 15(6): 425-429.

- Kapandji** IA: Funktionelle Anatomie des Menschen; Band 2, Untere Extremitäten; Bücherei des Orthopäden. Enke, Stuttgart, 1985. a) 168; b) 152; c) 194-195; d) 126-127; e) 182-183; f) 74, obr. 23-4.
- Katsouyanni** K, Skalkidis Y, Petridou E, Polychronopoulou-Trichopoulou A, Willett W, Trichopoulos D: Diet and peripheral arterial occlusive disease: the role of poly-, mono-, and saturated fatty acids. *Am J Epidemiol* 1991; 133(1): 24-31.
- Kawate** K, Ohneda Y, Kakihana T, Tamai S: Assessment of limp in patients with coxarthrosis-three-dimensional measurement of the pelvic rotation. *Nippon Seikeigeka Gakkai Zashi* 1992; 66(7): 643-656.
- Kelly** AM: Does the clinically significant difference in visual analogue scale pain vary with gender, age or cause of pain? *Acad Emerg Med* 1998; 5(11): 1086-90.
- Kieser** W: Die Seele der Muskeln – Krafttraining jenseits von Sport und Show. Walter Verlag, Zürich, 1998: 44-50.
- Kouvalchouk** JF, Bonnet JM, de Mondenard JP: Pyramidal syndrome. Apropos of 4 cases treated by surgery and review of the literature. *Rev Chir Orthop Reparatrice Appar Mot* 1996; 82(7): 647-657.
- Krawetz** P, Nance P: Gait analysis of spinal cord injured subjects: effects of injury level and spasticity. *Arch Phys Med Rehabil* 1996; 77(7): 635-638.
- Kusumoto** A, Suzuki T, Kumakura C, Ashizawa K: A comparative study of foot morphology between Filipino and Japanese women, with reference to the significance of a deformity like hallux valgus as a normal variation. *Ann Hum Biol* 1996; 23(51): 373-385.
- Lambert** C, Thomann B, Brunner R: Tibial torsion deformities. *Orthopäde* 2000; 29(9): 802-807.
- Lamont** EB, Christakis NA: Prognostic disclosure to patients with cancer near the end of life. *Ann Intern Med* 2001; 134(12): 1096-1105.
- Lang** J, Wachsmuth W: Praktische Anatomie – Band I, Teil IV, druhé vydání. Springer Verlag, Berlin, 1972: a) 355; b) 383-386; c) 396; d) 397; e) 394; f) 384-391; g) 152-159; h) 62-63; i) 382-383; j) 419 a 440.
- Langman** J: Medizinische Embryologie. Thieme, 1970: 139.
- Lanyon** P, Muir K, Doherty S, Doherty M: Assessment of a genetic contribution to osteoarthritis of the hip: sibling study. *BMJ* 2000; 321(7270): 1179-1183.
- Lariviere** LY, Miladi L, Dubousset JF, Seringe R: Failure of Dwyer's procedure in internal pes cavus in children. Physiopathological considerations and therapeutic deductions. *Rev Chir Orthop Reparatrice Appar Mot* 1985; 71(8): 563-573.
- Larsen** C: Funktionelle Frührehabilitation der Füße gemäß Spiraldynamikkonzept. *Trauma Berufskrankh*, 2001-3; Suppl 2: 242-247.
- Larsen** C: Der koordinierte Fuß. Neuer Merkur, München, 1996.
- Larsen** C: Die zwölf Grade der Einheit. Via Nova, 1995. A) 50-52; b) 202-249; c) 141; obr. 6.4 a-b.
- Larsen** C: Gut zu Fuß ein Leben lang. 4. Auflage. Trias Verlag, Stuttgart, 2013.
- Larsen** C: Prävention von Fußdeformitäten Krankengymnastik. Plaum, München, 1998; 50(9): 1534-1544.
- Larsen** C: Gesunde Füße für Ihr Kind. Trias Verlag, Stuttgart, 2002.
- Lau** EC, Cooper C, Lam D, Chan VN, Tsang KK, Sham A: Factors associated with osteoarthritis of the hip and knee in Hong Kong Chinese: obesity, joint injury, and occupational activities. *Am J Epidemiol* 2000; 152(9): 855-862.
- Laughlin** TJ: Complications of distal first metatarsal osteotomies. *J Foot Ankle Surg* 1995; 34(6): 524-531; Diskuse 593-594.
- Lee** RE, Goldberg JH, Sallis JF, Hickmann SA, Castro CN, Chen AH: A prospective analysis of the relationship between walking and mood in sedentary ethnic minority women. *Women Health* 2001; 32(4): 1-15.
- Leng** GC, Lee AJ, Fowkes FG, Whiteman M, Dunbar J, Housley E, Ruckley CV: Incidence, natural history and cardiovascular events in symptomatic and asymptomatic peripheral arterial disease in the general population. *Int J Epidemiol* 1996; 25(6): 1172-1181.
- Leon** AS, Casal D, Jacobs D Jr: Effect of 2,000 kcal per week of walking and stair climbing on physical fitness and risk factors for coronary heart

- disease. *J Cardiopulm Rehabil* 1996; 16(3): 183-192.
- Lillich JS, Baxter DE:** Common forefoot problems in runners. *Foot Ankle* 1986; 7: 145-151.
- Lord SE, Halligan PW, Wade DT:** Visual gait analysis: the development of a clinical assessment and scale. *Clin Rehabil* 1988; 12(2): 107-119.
- Lorei MP, Hershman EB:** Peripheral nerve injuries in athletes. Treatment and prevention. *Sports Med* 1993; 16(2): 130-147.
- Magee DJ:** Orthopaedic physical assessment. Saunders, Philadelphia, 1992. a) 451; b) 464; c) 484; d) 420-423.
- Maier E:** Der beste Schuh nützt nichts am falschen Fuß! *Podologie Mae* 2000: 8-10.
- Maier E:** Die Abwicklung des Fußes zum Schritt! *Podologie Apr* 1999: 12-21.
- Mantha S, Thisted R, Foss J, Ellis JE, Roizen MF:** *Anesth Analg* 1993; 77(5): 1041-1047.
- Marti B, Martin BW:** Sportliches Training oder Bewegung im Alltag zur Optimierung von Gesundheit und Lebensqualität. *Therapeutische Umschau Wellness – Fitness – Doping*. 2001; 58(4): 189-195.
- Marti B:** Health effects of recreational running in women. Some epidemiological and preventive aspects. *Sports Med* 1991; 11(1): 20-51.
- Marti B:** Benefits and risks of running among women: an epidemiologic study. *Int J Sports Med* 1988; 9(2): 92-98.
- Mathias S, Nayak US, Isaacs B:** Balance in elderly patients: the „get up and go“ test. *Arch Phys Med Rehabil* 1986; 67(6): 387-389.
- Mattalino AJ, Deese JM, Campbell ED:** Office evaluation and treatment of lower extremity injuries in the runner. *Clin Sports Med* 1989; 8(3): 461-475.
- Maxwell AJ, Anderson BE, Cooke JP:** Nutrition therapy for peripheral arterial disease: a double-blind, placebo-controlled, randomized trial of HeartBar. *Vasc Med* 2000; 5(1): 11-19.
- Menz HB, Lord SR:** Foot problems, function impairment, and falls in older people. *J Am Podiatr Med Assoc* 1999; 89(9): 458-467.
- Merkle B:** Das Achilles Projekt. *Ars Medici* 2000: 135-136.
- Min RJ, Navarro L:** Transcatheter duplex ultrasound-guided sclerotherapy for treatment of greater saphenous vein reflux: preliminary report. *Dermatol Surg* 2000; 26(5): 410-414; diskuse 413-414.
- Mondelli M, Giannini F, Reale F:** Clinical and electrophysiological findings and follow-up in tarsal tunnel syndrome. *Electroencephalogr Clin Neurophysiol* 1998; 109(5): 418-425.
- Morlock M:** Fuß. Und Sprunggelenkmodelle zur Belastungsanalyse – der Fuß im Sport. Kongressband 1. Wiesbadener Symposium 1997: 11-16.
- Morris D:** Bodywatching. Grafton books – Collins Publishing Group, 1987: a) 238; b) 241; c) 244.
- Morscher E, Ulrich J, Dick W:** Morton's intermetatarsal neuroma: Morphology and histological substrate. *Foot Ankle Int* 1998; 21(7): 558-562.
- Muller-Limmroth W, Bierlein HR, Diebschlag W:** Pressure distribution under the human footsole: qualitative and quantitative results. *Z Orthop Ihre Grenzgeb* 1977; 115(6): 929-936.
- Myles PS, Troedel S, Boquest M, Reeves M:** The pain visual analog scale: is it linear or nonlinear? *Anesth Analg* 1999; 89(6): 1517-1520.
- Myerson MS, Badekas A:** Hypermobility of the first ray. *Foot Ankle Clin* 2000; 5(3): 469-484.
- Napolitano C, Walsh S, Mahoney L, McCrea J:** Risk factors that may adversely modify the natural history of the pediatric pronated foot. *Clin Podiatr Med Surg* 2000; 17(3): 397-417.
- Nadkar MY, Samant RS, Vaidya SS, Borges NE:** Relationship between osteoarthritis of knee and menopause. *J Assoc Physicians India* 1999; 47(12): 1161-1163.
- Navarro L, Min RJ, Bone C:** Endovenous laser: a new minimally invasive method of treatment for varicose veins-preliminary observations using an 810 nm diode laser. *Dermatol Surg* 2001; 27(2): 117-122.
- Nielsen JF, Sinkjaer T:** A comparison of clinical and laboratory measures of spasticity. *Mult Scler* 1996; 1(5): 296-301.
- Niethard FU, Pfeil J:** Orthopädie – Duale Reihe, Hippokrates, Stuttgart 1992: a) 492; b) 500.
- Noszvai-Nagy M:** Zunehmend krankhafte Befunde an Kinderfüßen – Studie der Technischen Universität Karlsruhe. *Podologie* 1999; L(12): 4.
- Nurse MA, Nigg BM:** Quantifying a relationship between tactile and vibration sensitivity of the

- human foot with plantar pressure distributions during gait. *Clin Biomech* (Bristol, Avon) 1999; 14(9): 667-672.
- Nygaard IE:** Relationship between foot flexibility and urinary incontinence in nulliparous varsity athletes. *Obstet Gynecol* 1996; 87(6): 1049-1051.
- O'Byrne JM, Jenkinson A, O'Brien TM:** Quantitative analysis and classification of gait patterns in cerebral palsy using a three-dimensional motion analyzer. *J Child Neurol* 1998; 13(3): 101-108.
- Oglivie-Harris DJ, Carr MM, Fleming PJ:** The foot in ballet dancers: the importance of second toe length. *Foot Ankle Int* 1995; 16(3): 144-147.
- Olerud CM, Rosendahl Y:** Torsion-transmitting Properties of the hind foot. *Clin Orthop* 1998; (214): 285-294.
- Omey ML, Micheli LJ:** Foot and ankle problems in young athlete. *Med Sci Sports Exerc* 1999; (7 Suppl): 470-486.
- Ornish D, Scherwitz LW, Billings JH, Brown SE, Gould KL, Merritt TA, Sparler S, Armstrong WT, Ports TA, Kirkeeide RL, Hogeboom C, Brand RJ:** Intensive lifestyle changes for reversal of coronary heart disease. *JAMA* 1998; 280(23): 2001-2007.
- Ornish D, Brown SE, Scherwitz LW, Billings JH, Armstrong WT, Ports TA, McLanahan SM, Kirkeeide RL, Brand RJ, Gould KL:** Can lifestyle changes reverse coronary heart disease? The Lifestyle Heart Trial, *Lancet* 1990; 336(8708): 129-133.
- Orzechowski W, Wall A:** Selected anatomical features of the foot with hallux valgus. *Chir Narzadow Ruchu Ortop Pol* 1999; 64(3): 311-317.
- Parkkari J, Natri A, Kannus P, Manttari A, Laukkanen R, Haapasalo H, Nenonen A, Pasanen M, Oja P, Vuori I:** A controlled trial of the health benefits of regular walking on a golf course. *Am J Med* 2000; 109(2): 102-108.
- Pascarella EM, Estrada RJ:** Pes cavo-valgus foot. *J Foot Surg* 1991; 30(6): 553-557.
- Patla AE, Adkin A, Ballard T:** Online steering: coordination and control of body centre of mass, head and body reorientation. *Ex Brain Res* 1999; 129(4): 629-634.
- Peat G, McCarney R, Croft P:** Knee pain and osteoarthritis in older adults: a review of community burden and current use of primary health care. *Ann Rheum Dis* 2001; 60(2): 91-97.
- Pereira MA, Kriska AM, Day RD, Cauley JA, LaPorte RE, Kuller RH:** A randomized walking trial in postmenopausal women: effects on physical activity and health 10 years later. *Arch Intern Med* 1998; 158(15): 1695-1701.
- Perfetti C:** Der hemiplegische Patient – Kognitiv-therapeutische Übungen. Pflaum, München. 1997 a) 80-90; b) 34-35.
- Piccioli A, Prandoni P, Goldhaber SZ:** Epidemiologic characteristics, management, and outcome of deep venous thrombosis in a tertiary-care hospital: the Brigham and Women's Hospital DVT registry. *Am Heart J* 1996; 132(5): 1010-1014.
- Pisano F, Miscio G, Del Conte C, Pianca D, Candeloro E, Colombo R:** Quantitative measures of spasticity in post-stroke patients. *Clin Neurophysiol* 2000; 111(6): 1015-1022.
- Powell JT:** Vascular damage from smoking: disease mechanisms at the arterial wall. *Vasc Med* 1998; 3(1): 21-28.
- Price AE, Maisel R, Drennan JC:** Computed tomographic analysis of pes cavus. *J Pediatr Orthop* 1993; 13(5): 646-653.
- Priebe MM, Sherwood AM, Thornby JI, Kharas NF, Markowski J:** Clinical assessment of spasticity in spinal cord injury: a multidimensional problem. *Arch Phys Med Rehabil* 1996; 77(7): 713-716.
- Rao UB, Joseph B:** The influence of footwear on the prevalence of flat foot. *J Bone Joint Surg Br* 1992; 74(4): 525-527.
- Reimers J, Pedersen B, Brodersen A:** Foot deformity and the length of triceps surae in Danish children between 3 and 17 years old. *J Pediatr Orthop B* 1995; 4(1): 71-73.
- Renström PA:** Sportverletzungen und Überlastungsschäden – Prävention, Therapie, Rehabilitation. Enzyklopädie der Sportmedizin, medizinische Kommission des IOC. Deutscher Ärzte-Verlag, Köln, 1997 a) 213; b) 234; c) 232; d) 144; e) 350; f) 351-359; g) 352.
- Robbins S:** Foot position awareness in younger and older men: the influence of footwear sole properties. *J Am Geriatr Soc* 1997; 45(1): 61-66.

- Roche** Lexikon: Medizin. Urban Schwarzenberg, 1999 a) 367.
- Ronconi P**, Monachino P, Baleanu PM, Favilli G: Distal oblique osteotomy of the first metatarsal for the correction of hallux limitus and rigidus deformity. *J Foot Angle Surg* 2000; 39(3): 154-160.
- Rutherford OM**: Is there a role for exercise in the prevention of osteoporotic fractures? *Br J Sports Med* 1999; 33(6): 377-386.
- Sachithanandem V**, Joseph B: The influence of footwear on the prevalence of flat foot. A study of 1846 skeletally mature persons. *J Bone Joint Surg Br* 1995; 77(2): 254-257.
- Samama MM**: An epidemiologic study of risk factors for deep vein thrombosis in medical outpatients: the Sirius study. *Arch Intern Med* 2000; 160(22): 3415-3420.
- Savvidis E**, von der Decken CB: Forces acting on foot soles during stair climbing in healthy probands and in patients with coxarthrosis. *Bio-med Tech (Berlin)* 1999; 44(4): 98-103.
- Schad W**: Der Fuß des Menschen – ein lange verkanntes Detail seiner Evolution. Deutsche Assoziation für orthopädische Fußchirurgie. *D.A.F. News* 2000 5(2): 10-13.
- Schalow G**, Zäch GA: Reorganisation of the human CNS; theoretical basis for modern Neurorehabilitation. *Gen Physiol Biophys* 2000 (Suppl 1): 1-244.
- Schwarz T**, Schmidt B, Hohlein U, Beyer J, Schroeder HE, Schellong SM: Eligibility for home treatment of deep vein thrombosis: prospective study. *BMJ* 2001; 322(7296): 1192-1193.
- Scurr JH**, Machin SJ, Bailey-King S, Mackie IJ, McDonald S, Smith PD: Frequency and prevention of symptomless deep-vein thrombosis in long-haul flights: a randomized trial. *Lancet* 2001; 357(9267): 1485-1489.
- Segesser B**, Morscher E, Goesele A: Lesions of the growth plate caused by sports stress. *Orthopaedie* 1995; 24(5): 446-456.
- Seichert N**: Funktionelle instrumentierte Ganganalyse (FIGA) – neuartiges Verfahren und klinische Relevanz. Symposium *SUVA Klinik* Bellikon, Schweiz ze dne 4. 12. 1997; Persönliche Mitteilung, 2000.
- Shanahan MD**, Douglas DL, Sharrard WJ, Duckworth T, Betts R: The long-term results of the surgical management of paralytic pes cavus by soft tissue release and tendon transfer. *Z Kinderchir* 1985; 40 Suppl 1: 37-41.
- Sherwood AM**, Graves DE, Priebe MM: Altered motor control and spasticity after spinal cord injury: subjective and objective assessment. *J Rehabil Res Dev* 2000; 37(1): 41-52.
- Snow RE** (1994): High heeled shoes: their effect on centre of mass position, posture, three-dimensional kinematics, rearfoot motion and ground reaction forces. *Arch Phys Med Rehabil* 1994; 75(5): 568-576.
- Sobel E**, Levitz SJ, Caselli MA, Tran M, Lepore F, Lilja E, Sinaie M, Wain E: Reevaluation of relaxed calcaneal stance position. Reliability and normal values in children and adults. *J Am Podiatr Med Assoc* 1999; 89(5): 258-264.
- Sobel E**, Levitz SJ, Caselli MA, Brentnall Z, Tran MQ: Natural history of the rearfoot angle: preliminary values in 150 children. *Foot Ankle Int* 1999; 20(2): 119-125.
- Sobel E**, Caselli MA, Velz Z: Effect of persistent toe walking on ankle equines. Analysis of 60 idiopathic toe walkers. *J Am Podiatr Med Assoc* 1997; 87(1): 17-22.
- Sohier R**, Sohier J: Das analytische Konzept. Edition Kiné-Sciences, 1991 (1): 149-150.
- Staheli LT**: Torsion – treatment indications. *Clin Orthop* 1989; (247): 61-66.
- Staheli LT**: In-toeing and out-toeing in children. *J Fam Pract* 1983; 16(5): 1005-1011.
- Strecker W**, Keppler P, Gebhard F, Kinzl L: Length and torsion to the lower limb. *J Bone Joint Surg Br* 1997; 79(6): 1019-1023.
- Sun Tsu**: Wahrhaft siegt, wer nicht kämpft. Der Chinesische Klassiker bearbeitet von Thomas Cleary. Bauer, 1992: 11-15.
- Suter E**, Marti B, Gutzwiller F: Jogging or walking – comparison of health effects. *Ann Epidemiol* 1994; 4(5): 375-381.
- Tanaka T**, Takeda H, Izumi T, Ino S, Ifukube T: Effects on the location of the centre of gravity and the foot pressure contribution to standing balance associated with ageing. *Ergonomics* 1999; 42(7): 997-1010.
- Taylor J**, Tucker LA: Comparison of the CardioGlide, CrossWalk, and treadmill walking in development of cardiovascular endurance, dynamic strength, and flexibility in middle-aged

- men and women. *Percept Mot Skills* 1996; 82(3 Pt 1): 875-882.
- Thelen E**, Ulrich BD, Niles D: Bilateral coordination in human infants: stepping on a spilt-belt treadmill. *Journal of Experimental Psychology Med* 1987; 13: 405-410.
- Thune I**, Furberg AS: Physical activity and cancer risk: dose-response and cancer, all sites a site-specific. *Med Sci Sports Exerc* 2001; 33(6 Suppl): S530-S550; diskuse S609-610.
- Treiman GS**, Copland S, McNamara RM, Yellin AE, Schneider PA, Treiman RL: Factor influencing ulcer healing in patients with combined arterial and venous insufficiency. *J Vasc Surg* 2001; 33(6): 1158-1164.
- Trnka HJ**, Zettl R, Hungerford M, Muhlbauer M, Ritschl P: Acquired hallux varus and clinical tolerability. *Foot Ankle Int* 1997; 18(9): 593-597.
- Turnbull LA**, Wood N, Kester G: Controlled trial of the subjective patient benefits of accompanied walking to the operating theatre. *Int J Clin Pract* 1998; 52(2): 81-83.
- Turner AP**, Barlow JH, Heathcote-Elliott C: Long term health impact of playing professional football in the United Kingdom. *Br J Sports Med* 2000; 34(5): 332-336.
- US Department of Health and Human Services**: Physical Activity and Health – a Report of the Surgeon General. Atlanta 1996, www.cdc.gov.
- Van den Oever R**, Hepp B, Debbaut B, Simon I: Socio-economic impact of chronic venous insufficiency. An underestimated public health problem. *Int Angiol* 1998; 17(3): 161-167.
- Van Gils CC**, Streed RH, Page JC: Torsion of the human Achilles tendon. *J Foot Angle Surg* 1996; 35(1): 41-48.
- Van Lith R**: Pathogenese und Pathomechanik des Vorfußes. *Podologie* 1998; XLIX(7): 18-24.
- Verhaegen MJ**: Aquatic ape theory and fossil hominids. *Med Hypotheses* 1991; 35(2): 108-114.
- Verhaegen MJ**: The Aquatic Ape Theory: evidence and a possible scenario. *Med Hypotheses* 1985; 16(1): 17-32.
- Vingard E**, Alfredsson L, Malchau H: Lifestyle factors and hip arthrosis. A case referent study of body mass index, smoking and hormone therapy in 530 Swedish women. *Acta Orthop Scand* 1997; 68(3): 216-220.
- Vingard E**, Alfredsson L, Goldie I, Hgstedt C: Sports and osteoarthritis of the hip. An epidemiologic study. *Am J Sports Med* 1993; 21(2): 195-200.
- Waibel P**: Differences in the prognosis of arterial reconstruction in stage II and IV peripheral arterial disease. Results of a minimum of 15 to 25 years follow-up. *Vasa* 1992; 21(1): 46-51.
- Waidelich HA**, Strecker W, Schneider E: Computed torsion-angle and length management of the lower extremity. The methods, normal values and radiation load. *Rofo Fortschr Gen Röntgenstr Neuen Bildgeb Verfahr* 1992; 157(3): 245-251.
- Waldecker U**: Plantar fat pad atrophy: a cause of metatarsalgia? *J Foot Angle Surg* 2001; 40(1): 21-27.
- Walker RD**, Nawaz S, Wilkinson CH, Saxton JM, Pockley AG, Wood RF: Influence of upper- and lower-limb exercise training on cardiovascular function and walking distances in patients with intermittent claudication. *J Vasc Surg* 2000; 31(4): 662-669.
- Weineck J**: Optimales Training – leistungsphysiologische Trainingslehre. Demeter Spitta, Balingen, 1997. a) 187-191.
- Weineck J**: Sportanatomie – Beiträge zur Sportmedizin Band 9. Perimed, Erlangen, 1998. a) 168.
- WHO** World Health Organisation: The World Health Report 1997, Genf.
- Widhe T**: Foot deformities at birth: a longitudinal prospective study over a 16-year period. *J Pediatr Orthop* 1997; 17(1): 20-24.
- Widmer LK**, Waibel P: Arterielle Durchblutungsstörungen in der Praxis. Roche – wissenschaftlicher Dienst. Hand Huber, 1972. a) 47; b) 40.
- Wiedmer L**, Langer T, Knusel O: Gait pattern of patients with hip arthritis. *Orthopade* 1992; 21(1): 35-40.
- Wigger P**: Surgical therapy of primary varicose veins. *Schweiz Med Wochenschr* 1998; 128(45): 1781-1788.
- Williams KR**, Cavanagh PR: Biomechanical correlates with running economy in elite distance runners. *Proceedings of the North American Congress on Biomechanics; American Society*

of Biomechanics, Montreal; 1986, Vol 2: 287-288.

Windhager R, Lack W, Kutschera H, Wimberger W, Mayr M: Clinical and radiologic comparison of „idiopathic“ and „neurogenic“ pes cavus. *Z Orthop Ihre Grenzgeb* 1989; 127(2): 169-173.

Wolfson L, Whipple R, Derby C, Judge J, King M, Amerman P, Smith J, Smyers D: Balance and strength training in older adults: intervention gains and Tai Chi maintenance. *J Am Geriatr Soc* 1996; 44(5): 498-506.

Woolf-May K, Bird S, Owen A: Effects of an 18 week walking programme on cardiac function in previously sedentary or relatively inactive adults. *Br J Sports Med* 1997; 31(1): 48-53.

Wu KK: Morton neuroma and metatarsalgia. *Curr Opin Rheumatol* 2000; 12(2): 131-142.

Wyss JM, Carlson SH: The role of the central nervous system in hypertension. *Curr Hypertens Rep* 1999; 1(3): 246-253.

Yoshimura N, Sasaki S, Iwasaki K, Danjoh S, Kinoshita H, Yasuda T, Tamaki T, Hashimoto T, Kellingray S, Croft P, Coggon D, Cooper C: Occupational lifting is associated with hip osteoarthritis: a Japanese case – control study. *J Rheumatol* 2000; 27(2): 434-440.

Yu GV, Sellers CS, Shook JE, Karlock LG: Iatrogenic deformities of the first ray. *Clin Podiatr Med Surg* 1996; 13(3): 367-422.

Zanetti M, Strehle JK, Zollinger H, Hodler J: Morton neuroma and fluid in intermetatarsal bursae on MR images of 70 asymptomatic volunteers. *Radiology* 1997; 203(2): 516-520.

Zatsiorsky VM: Biomechanics in Sport, Volume IX of the Encyclopaedia of Sports Medicine. IOC Medical Commission Publication. Blackwell Science, Oxford, 2000. a) 164-174; b) 224; c) 578-579; d) 580; obr. 27. 3.

Zollinger H, Fellmann J: Natural course of juvenile foot deformities. *Orthopäde* 1994; 23(3): 206-210.

17.1 Důležité adresy a odkazy

BIMS® Posuvné měřítko k určení délky dětského chodidla a vnitřní délky stélky www.bims.ch.

EMED® EMED-System von Novel, München. Přístroj je k dispozici ve formě Orthopädie Bähler, Zürich.

FiGA® Funkční instrumentální analýza chůze www.rehabellikon.ch.

ISD Institut medicínsko-terapeutické Spiraldynamik® Zürich, isd@hin.ch.

MBT® Technologie bosonohých Masajů www.masai.ch.

SDI Spiraldynamik® International Zürich, www.spiraldynamik.com.