VII SEZNAM POUŽITÉ SOUVISEJÍCÍ LITERATURY

Bodénès C., CHancerel E., Gailing O., Vendramin G. G., Bagnoli F., Durand J., Goicoechea P. G., Soliani C., Villani F., Mattioni C., Koelewin H. P., Murat F., Salse J., Roussel G., Boury C., Alberto F., Kremera A., Plomion C. 2012. Comparative mapping in the Fagaceae and beyond with EST-SSRs. BMC Plant Biology 12: 153. https://doi.org/10.1186/1471-2229-12-153

Bordács S., Zhelev P., Schirone B. 2019. EUFORGEN Technical Guidelines for genetic conservation and use of pubescent oak (*Quercus pubescens*). European Forest Genetic Resources Programme (EUFOR GEN), European Forest Institute: 6 s.

Bruschi P., Vendramin G. G., Bussotti F., Grossoni P. 2000. Morphological and Molecular Differentiation between *Quercus petraea* (Matt.) Liebl. and *Quercus pubescens* Willd. *(Fagaceae)* in Northern and Central Italy. Annals of Botany 85 (3): 325–333. http://www.jstor.org/stable/42770646

Burger K., Gailing O. 2022. Genetic variability of indigenous (*Quercus robur* L.) and late flushing oak (*Quercus robur* L. subsp. *slavonica* (Gáyer) Mátyás) in adult stands compared with their natural regeneration. European Journal of Forest Research, 141: 1073–1088. https://doi.org/10.1007/s10342-022-01491-3

Di Pietro R., Di Marzio P., Antonecchia G., Conte A. L., Fortini P. 2020. Preliminary characterization of the *Quercus pubescens* complex in southern Italy using molecular markers. Acta Botanica Croatia, 79 (1): 15–25.

Dow B. D., Ashley M. V., Howe H. F. 1995. Characterization of highly variable (GA/CT) n microsatellites in the bur oak, *Quercus macrocarpa*. TAG. Theoretical and applied genetics. Theoretische und angewandte Genetik, 91: 137–141.

Durand J., Bodénès C., Chancerel E. et al. 2010. A fast and cost-effective approach to develop and map EST-SSR markers: oak as a case study. BMC Genomics, 11: 570. https://doi.org/10.1186/1471-2164-11-570

Dzialuk A., Chybicki I., Burczyk J. 2005. PCR multiplexing of nuclear microsatellite loci in *Quercus* species. Plant Molecular Biology Reporter, 23: 121–128. https://doi.org/10.1007/BF02772702

Falush D., Stephens M., Pritchard J. K. 2003. Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. Genetics, 164(4): 1567–1587.

Falush D., Stephens M., Pritchard J. K. 2007. Inference of population structure using multilocus genotype data: dominant markers and null alleles. Molecular Ecology, 7(4): 574–578.

Grulich V. 2017. Červený seznam cévnatých rostlin ČR. In: Grulich V., Chobot K. (eds.): Červený seznam ohrožených druhů České republiky. Cévnaté rostliny. Příroda, 35: 75–132.

Guichoux E., Lagache L., Wagner S., Léger P., Petit R. J. 2011. Two highly validated multiplexes (12-plex and 8-plex) for species delimitation and parentage analysis in oaks (*Quercus spp.*). Molecular Ecology Resources, 11: 578–585.

Hejný S., Slavík B. (eds.) 1990. Květena České republiky 2. Praha, Academia: 540 s.

Hubisz M. J., Falush D., Stephens M., Pritchard J. K. 2009. Inferring weak population structure with the assistance of sample group information. Molecular Ecology Resources, 9 (5): 1322–1332.

Isagi Y., Suhandono S. 1997. PCR primers amplifying microsatellite loci of *Quercus myrsinifolia* Blume and their conservation between oak species. Molecular Ecology, 6: 897–899.

Kalinowski S. T., Taper M. L., Marshall T. C. 2007. Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. Molecular Ecology, 16: 1099–1106.

Kampfer S., Lexer C., Glössl J., Steinkellner H. 1998. Characterization of (GA)n Microsatellite Loci from *Quercus Robur*. Hereditas, 129: 183–186.

Koontz A. C., Schumacher E. K., Spence E. S., Hoban S. M. 2024. Ex situ conservation of two rare oak species using microsatellite and SNP markers. Evolutionary Applications, 17: e13650. https://doi.org/10.1111/eva.13650

Lexer C., Heinze B., Steinkellner H., Kampfer S., Ziegenhagen B., Glössl J. 1999. Microsatellite analysis of maternal half-sib families of *Quercus robur*, pedunculate oak: detection of seed contaminations and inference of the seed parents from the offspring. Theoretical and Applied Genetics, 99: 185–191.

Li Y. L., Liu J. X. 2018. Structure Selector: A web based software to select and visualize the optimal number of clusters using multiple methods. Molecular Ecology Resources, 18: 176–177.

Máchová P., Trčková O., Cvrčková H. 2018. Use of nuclear microsatellite loci for evaluating genetic diversity of selected populations of *Picea abies* (L.) Karsten in the Czech Republic. Forests, 9 (92): 1–15. doi:10.3390/f9020092

Maghuly F., Pinsker W., Praznik W., Fluch S. 2006. Genetic diversity in managed subpopulations of Norway spruce [*Picea abies* (L). Karst.]. Forest Ecology and Management, 222: 266–271.

Mullis K. B. 1990. The Unusual Origin of the Polymerase Chain Reaction. Scientific American, 262: 56–65.

Nei M. 1972. Genetic distance between populations. American Naturalist, 106: 283–392.

Peakall R., Smouse P. E. 2006. GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. Molecular Ecology Notes, 6: 288–295.

Peakall R., Smouse P. E. 2012. GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research – an update. Bioinformatics, 28: 2537–2539.

Pluess A. R., Määttänen K. 2013. Characterization of eighteen novel microsatellite markers and multiplex PCR protocol for *Fagus sylvatica*. Conservation Genetics Resources, 5: 311–314.

Pritchard J. K., Stephens M., Donnelly P. 2000. Inference of population structure using multilocus genotype data. Genetics, 155(2): 945–959.

Raymond M., Rousset F. 1995. GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. Journal of Heredity, 86: 248–249.

Rousset F. 2008. Genepop'007: a complete reimplementation of the Genepop software for Windows and Linux. Molecular Ecology Resources, 8: 103–106.

Schueler S., Tusch A., Schuster M., Ziegenhagen B. 2003. Characterization of microsatellites in wild and sweet cherry (*Prunus avium* L.) – markers for individual identification and reproductive processes. Genome, 46: 95–102.

Steinkellner H., Lexer C., Turetschek E., Glössl J. 1997. Conservation of (GA) n microsatellite loci between *Quercus species*. Molecular Ecology, 6: 1189–1194.

Svoboda P. 1955. Lesní dřeviny a jejich porosty. Část II. Státní zemědělské nakladatelství, Praha.

Thunder R., Conrad A. O., Burdine Ch., Yang J., Lhotka J. M., Abbott A. G., Nelson C. D. 2022. White oak (*Quercus alba* L.) microsatellite markers for genetic diversity studies. Research Note: 6 s. https://doi.org/10.2737/SRS-RN-26

Unger G. M., Konrad H., Geburek T. 2011. Does spatial genetic structure increase with altitude? An answer from *Picea abies* in Tyrol, Austria. Plant Systematics and Evolution, 292: 133–141.

Úradníček L., Maděra P., Tichá S., Koblížek J. 2009. Dřeviny České republiky. Kostelec nad Černými lesy, Lesnická práce: 367 s.

Van Oosterhout C., Hutchinson W. F., Wills D. P. M., Shipley P. 2004. Micro-Checker: software for identifying and correcting genotyping errors in microsatellite data. Molecular Ecology Notes, 4: 535–538.

Vornam B., Decarli N., Gailing O. 2004. Spatial distribution of genetic variation in a natural beech stand (*Fagus sylvatica* L.) based on microsatellite markers. Conservation Genetics, 5: 561–570.

Wagner S., Gerber S., Petit R. J. 2012. Two highly informative dinucleotide SSR multiplexes for the conifer *Larix decidua* (European larch). Molecular Ecology Resources, 12 (4): 717–725. DOI 10.1111/j.1755-0998.2012.03139.x

Wójkiewicz B., Lewandowski A., Żukowska W. B., Litkowiec M., Wachowiak W. 2021. Low effective population size and high spatial genetic structure of black poplar populations from Oder valley in Poland. Annals of Forest Science, 78: 37. DOI 10.1007/s13595-021-01055-2