Literatura

Albright, AL. Spastic cerebral palsy: Approaches to Drug Treatment, 1995, *CNS Drugs*, p. 17–27.

Albright, AL., Cervi, A., Singletary, J. Intrathecal baclofen for spasticity in cerebral palsy. *JAMA*, 1991, p. 1418–1422.

Alexander, RD., Davids, JR., Peace, LC., Gidewall, MA. Wrist arthrodesis in childern with cerebral palsy. *J. Pediatr. Orthop.*, 2000, 20, 4, p. 490–495.

Anderson, JP., Snow, B., Dorey, J., Kabo, JM. Efficacy of soft splints in reducing severe knee – flexion contractures. *Dev. Med. Child Neurol.*, 1988, 30, 4, p. 502–508.

Arens, LJ., Peacock, WJ., Peter, J. Selective posterior rhizotomy; A long-term follow-up study. *Childs Neur. Syst.*, 1989, 5, p. 148–152.

Atar, D., Grant, AD., Bash, J., Lehman, WB. Combined hip surgery in cerebral palsy patients. *Am. J. Orthop.*, 1995, 24, 1, p. 52–55.

Baker, R., Jasinski, M., Maciag-Tymecka, I., Michalowska-Mrozek, J., Bonikowski, M., et al. Botulinum toxin (Dysport®) in the treatment of dynamic equinus spasticity in patients with diplegic cerebral palsy: results of a randomised, double-blind, placebo-controlled, dose-ranging study. *Dev. Med. Child Neurol.*, 2002, in print.

Bakheit, AMO., Severa, p., Cosgrove, A., Morton, R., Roussonis, SH., Doderlein, L., Lin, JP. Safety profile and efficacy of botulinum toxin A (Dysport®) in children with muscle spasticity. *Dev. Med. Child Neurol.*, 2001, 43, p. 234–238.

Ballard, RA. Resuscitation in the Delivery Room. In Taeusch, H.W., Ballard, RA. (Eds.) *Avery´s Diseases of the Newborn*, 7th ed. W.B.Saunders Comp., 1998, p. 319–333.

Barkovich, AJ., Hallam, D. Neuroimaging in perinatal hypoxic-ischemic injury. *Ment. Retard. Dev. Disablil. Res. Rev.*, 1997, 3, p. 28–41.

Barric, JL., Galasko, CS. Surgery for unstable hips in cerebral palsy. *J. Pediatr. Orthop. B.*, 1996, 5, 3, p. 225–231.

Baud, O., Foix-L´Helias, L., Kaminski, M., et al. Antenatal glucosteroid treatment and cystic periventricular leukomalacia in very premature infants. *N. Engl. J. Med.*, 1999, 341, 16, p. 1190–1196.

Baumann, JU. *Operative Behandlung der infantilen Zerebralparesen*. Stuttgart: G. Thieme, 1970, p. 104.

Blair, E, Stanley, F. Causal pathways to cerebral palsy. *Curr. Pediatr.*, 2002, 12, p. 179–185.

Bobath, B., Bobath, K. *Motor Development in the Different Types of Cerebral Palsy*. London : Heinemann Medical books, 1975.

Bobath, K. *Neurophysiological Basis for the treatment of Cerebral palsy*. Oxford : Spastic International Medical Publications, 1980.

Bobath, B. *Abnormal postural reflex activity caused by brain lesions*. London : Heinemann, 1965.

Course Notes to the Bobath course, London : Bobath Centre, 1997.

Bobath, B. The very early treatment od cerebral palsy. *Dev. Med. Child Neurol.*, 1967, 9, p. 373–390.

KNIHOVNA FAKULTY TV UP - OLOMOUC

Boylan, GB., Young, K., Panerai, RB., et al. Dynamic Cerebral Autoregulation in Sick Newborn Infants. *Pediat. Res.*, 2000, 48, 1, p. 12–17.

Brázdilová, N. *Organicky podmíněné dysfunkce CNS*. Brno: Institut pro další vzdělávání pracovníků ve zdravotnictví, 1996, ISBN 80-7013-221-3.

Brereton, A. le Gay, Sattler, J. *Cerebral Palsy, Basic Abilities*. New South Wales Spactic Society, 1975.

Broom, MJ., Banta, JV., Renshaw, TS. Spinal fusion augmented by Luque –rod segmental instrumentation for neuromuscular scoliosis. *J. Bone Jt. Surg.*, 1989, 71A, 1, p. 32–34.

Brunner, R. Which procedure gives best resuslts in reconstructing dislocated hip joints in cerebral palsy. *Acta Orthop. Belg.*, 1998, 64, 1, p. 7–16.

Brunner, R., Baumann, JU. Clinical benefit of reconstruction of dislocated or subluxated hip joints in patients with spastic cerebral palsy. *J. Pediatr. Orthop.*, 1994, 14, 2, p. 290–294.

Brunner, R., Baumann, JU. Long- term effects of intertrochanteric varus-derotation osteotomy on femur and acetabulum in spastic cerebral palsy an 11- to 18- year follow-up study. *J. Pediatr. Orthop.*, 1997, 17, 5, p. 585–591.

Calderon-Gonzalez, R., Calderon-Sepulveda, R., Rincon-Reyes, M., Garcia-Ramirez, J., Mino-Arango, E. Botulinum toxin A in management of cerebral palsy. *Pediatric Neurology*, 1994, 10, p. 284–288.

Capko, J. *Základy fyziatrické léčby*. Praha : Grada Publishing, 1998, p. 65, s. 365–371.

Carpenter, EB., Seitz, DG. Intramuscular alcohol as an aid in the management of spastic cerebral palsy. *Dev. Med. Child Neurol.*, 1980, 22, p. 497–501.

Cendrowski, W., Sobczyk, W. Clonazepam, baclofen, and placebo in the treatment of spasticity. *Eur. Neurol.*, 1977, 16, p. 257–62.

Cesari, B., Touzet, P., Journeau, P., Padovani, J.P., Rigault, P., Pouliquen, M. Value of pelvic osteotomy in the management of the hip in childern with cerebral palsy. *Rev. Chir. Orthop. Reparatrice. Appar. Mot.*, 1995, 81, 4, p. 310–316.

Cmunt, E. *Ortopedická obuv*. Praha : J. Cendelín, 1996, s. 62.

Committee on Fetus and Newborn, American Academy of Peddiatrics, and Committee on Obstetric Practice, American College of Obstetricians and Gynecologists: Use and Abuse of the Apgar Score. *Pediatrics*, 1996, 98, 1, p. 141–143.

Corbett, M., Frankel, HL., Michaelis, L. A double-blind cross-over trial of valium in the treatment of spasticity. *Paraplegia*, 1972, 10, p. 19–22.

Cornell, MS. The hip in cerebral palsy. *Dev. Med. Child Neurol.*, 1995, 37, 1, p. 3–18.

Corry, IS., Cosgrove, AP., Walsch, EG., McClean, D., Graham, HK. Botulinum toxin A in the hemiplegic upper limb: a double-blind trial. *Dev. Med. Child Neurol.*, 1997, 39, p. 185–193.

Cosgrove, AP., Covry, IS., et al. Botulinum toxin in management of lower limb in cerebral palsy. *Dev. Med. Child Neurol.*, 1994, 36, p. 386–396.

Cosgrove, AP., Graham, HK. Botulinum toxin A prevents the development of contractures in the hereditary spastic mouse. *Dev. Med. Child Neurol.*, 1994, 36, p. 379–385.

Cosgrove, AP., Corry, IS., Graham, HK. Botulinum Toxin in the management of the lower limb in cerebral palsy. *Dev. Med. Child Neurol*, 1994, 36, p. 386–396.

Cosgrove, AP., Graham, HK. Cerebral Palsy. In: Moore AP. (Ed.) *Handbook of Botulinum Toxin*. Oxford: Blackwell Science, 1997, p. 222–247.

Cottalorda, J., Gautheron, V., Metton, G., Charmet, E., Chavrier, Y. Toe-walking in children younger than six years with cerebral palsy. The contribution of seriál corrective casts. *J. Bone Joint Surg Br.*, 2000, 82, p. 541–544.

Cséfalvay, Z. Dyzartria. In Lechta,V., a kol. *Logopedické repetitórium*. Bratislava : SPN, 1990.

Cséfalvay, Z. Diagnostika dyzartrie. In Lechta, V., a kol.: *Diagnostika narušenej komunikačnej dovednosti*. Martin : Osveta, 1995.

Cunningham,C. What is an Adequate Blood Pressure in the Preterm Newborn? In Hansen, TN., McIntosh, N.(Eds.) *Current Topics in Neonatology 3*. W.B.Saunders, 1999, p. 62–92.

Delivoria-Papadopoulos, M., Mishra, OP. Mechanisms of cerebral injury in perinatal asphyxia and strategies for prevention. *J. Pediatr.*, 1998, 132, 3, Pt 2, p. S30–S34.

Denislic, M., Meh, D. Botulinum toxin in the treatment of cerebral palsy. *Neuropediatrics*, 1995, 26, p. 249–52.

Dietz, F.R., Knutson, L.M. Chiari pelvic osteotomy in cerebral palsy. *J. Pediatr. Orthop.* 1995, 15, 3, p. 372–380.

Dittrich, J., Jirout, J., Vlach, V. Pneumoencephalographic findigs in various forms of cerebral palsy. *Cerebral palsy bulletin 4*, 1960, p. 286–289.

Dobeš, M., Dobešová, P. *Cvičíme na velkém míči*. Havířov – město : Domiga 1996, s. 4–5.

Duffy, CM., Cosgrove, APOD. The foot in cerebral palsy. *Curr. Orthop.*, 2002, 16, 2, p. 104–113.

Dvořák, J. *Logopedický slovník*. Žďár nad Sázavou : Logopedické centrum, 1998.

Dyke, CG., Davidoff, LM., Masson, CB. Cerebral hemiatrophy with homolateral hypertrophy of the skull and sinuses. *Surg. Gynecol. Obstet.*, 1933, 57, p. 588–600.

Edwards, AD., Azzopardi, DV. Perinatal Hypoxia-Ischemia and Brain Injury. Commentary. *Pediatr. Res.*, 2000, 47, 4, p. 431–432.

Ekblom, B.: Effects of the hip abduction orthosis on muscle activity in children with cerebral palsy. *Physiother. Theory Pract.*, 2002, 18, 1, p. 55–63.

Fasano, VA., Barolat-Romana, G., Zeme, p., Sguazzi, A. Surgical treatment of spasticity in cerebral palsy. *Child's Brain*, 1978, 4, p. 289–305.

Fasano, VA., Barolat-Romana, G., Zeme, p., Sguazzi, A. Electrophysiological assessment of spinal circuits in spasticity by direct dorsal root stimulation. *Neurosurgery*, 1979, 4, p. 146–151.

Feet, BA., Gilland, E., Groenendaal, F., et al. Cerebral excitatory amino acids and Na+, K+-ATPase during resuscitation of severely hypoxic newborn piglets. *Acta Paediatr.*, 1998, 87, p. 889–895.

Fellman, V., Raivio, KO. Reperfusion Injury as the Mechanism of Brain Damage after Perinatal Asphyxia. Review Article. *Pediatr. Res.*, 1997, 41, 5, p. 599–606.

Fetters L. Cerebral palsy: Comtemporary treatment concepts. In Lister MJ. (Ed.): *Contemporary Management of Motor Control Problems*. Proceedings of the II Step Conference. Alexandria : Foudation for Physical Therapy, 1991, p. 219–224.

Filipková, M. *Komplexní fyzioterapie u rizikových dětí*. (Absolventská práce) Brno, SZŠ a VOŠ Merhautova, 2000, s. 38–42.

Finer, NN., Horbar, JD., Carpenter, JH. Cardiopulmonary resuscitation in the very low birth weight infant: the Vermont Oxford Network experience. *Pediatrics*, 1999, 104, 3, Pt 1, p. 428–434.

Finnstrom, O., Otterbald, O., Sedin, G., et al. Neurosensory outcome and growth at three years in extremely low birthweight infants: follow-up results from the Swedish national prospective study. *Acta Paediatr.*, 1998, 87, 10, p. 1055–1060.

Flett, PJ., Stern, LM., Waddy, H., Connel, TM., Seeger, JD., Gibson, SK. Botulinum toxin A versus fixed cast stretching for dynamic calf tightness in cerebral palsy. *Journal of Pediatrics and Child Health*, 1999, 35, p. 71–77

Foerster, O. On the indications and results of the excision of posterior spinal roots in men. *Surg. Gynecol. Ostet.* 1913, 16, 463–464.

Foulon, W., Van Lidekerke, D., Demanet, C. Markers of infection and their relationship to preterm delivery. *Am. J. Perinatol.*, 1995, 12, 3, p. 208–211.

Franz, DN., Perry, RS. Mechanisms for differential blocks among single myelinated and non-myelinated axons by procaine. *J. Physiol.* (Lond.), 1974, 236, p. 193–210.

Gabos, PG., Miller, F., Galban, MA., Gupta, GG., Dabney, K. Prosthetic interposition arthroplasty for the palliative treatment of end-stage spastic hip disease in nonambulatory patients with cerebral palsy. *J. Pediatr. Ortop.*, 1999, 19, 6, p. 796–804.

Gage, JR, Novacheck, TF. An update on the treatment of gait problems in cerebral palsy. *J. Pediatr. Orthop. B.*, 2001, 10, p. 265–274

Gangale, DC. *The source for Oral-Facial Exercises*. East Moline, USA : Lingui Systems, 2001.

Gerstmann, DR., Wood, M., Miller, A., et al. Childhood outcome after early high-frequency oscillatory ventilation for neonatal respiratory distress syndrome. *Pediatrics*, 2001, 108, 3, p. 617–623.

Givon, U., Miller, F. Shortening of a unit rod protruding into the hip joint: Case report and description of a surgical technique. *J. Spinal Disord.*, 1999, 12, 1, p. 74–76.

Goddard-Finegold, J. Introduction and Overview of Antenatal Central Nervous System Insults. In Taeusch, HW. and Ballard RA. (Eds.) *Avery´s Diseases of the Newborn*, VIIth ed., W.B.Saunders Comp., 1998, p. 797–801.

Graham, HK., Aoki, KR., Autti-Ramo, I., et al. Recommendations for the use of botulinum toxin type A in the management of cerebral palsy. *Gait Posture*, 2000, 11, p. 67–79.

Grimby, G., Andren, E. Progress of disability in daily activities in adults with cerebral palsy or spina bifida. Abstracts of 3rd World Congress in Neurological Rehabilitation, 2002, p. 77.

Gros, C., Ourknine, G., Vlahovitch, B., Frerebeau, P. La radicotomie sélective postérioeure dans le traitment neuro-chirurgical de l'hypertonie pyramidale. *Neurochirurgie,* 1967, 13, p. 505–518.

Gros, C. Spasticity-Clinical classification and surgical treatment. *Adv. Tech. Stand. Neurosurg.*, 1979, 6, p. 55–97.

Hadraba, I.: *Ortopedická protetika*. Praha : Státní pedagogické nakladatelství, 1986, s. 62.

Hagberg, G., Sanner, G., Steen, M. The dysequilibrium syndrome in cerebral palsy. Clinical aspects and treatment. *Acta Paediatr. Scand.*, 1972, Suppl. 226.

Hagberg, B., Hagberg, G., Olow, I. The changing panorama of cerebral palsy in Sweden 1954–1970. I. Analysis of general changes. *Acta Paediatr. Scand.*, 1975, 64, p. 187–192.

Hagberg, B., Hagberg, G., Olow, I. The changing panorama of cerebral palsy in Sweden 1954–1970. II. Analysis of various syndromes. *Acta Paediatr. Scand.*, 1975, 64, p. 193–200.

Hagberg, B., Hagberg, G., Olow, I. The changing panorama of cerebral palsy in Sweden. IV. Epidemiological trends 1959–1978. *Acta Paediatr. Scand.*, 1984, 73, p. 433–440.

Hagberg, B., Hagberg, G., Olow, I., Von Wendt, L. The changing panorama of cerebral palsy in Sweden. V. The birth year period 1979–1982. *Acta Paediatr. Scand.*, 1989, 78, p. 283–290.

Hagberg, B., Hagberg, G., Olow, I. The changing panorama of cerebral palsy in Sweden. VI. Prevalence and origin during the birth period 1983–1986. *Acta Paediatr. Scand.*, 1993, 82, p. 387–393.

Hagberg, B., Hagberg, G., Beckung, E., Uvebrant, P. Changing panorama of cerebral palsy in Sweden. VIII. Prevalence and origin in the birth period 1991–1994. *Acta Paediatr. Scand.*, 2001, 90, p. 271–277.

Hargreaves, DG., Warrvick, DJ., Tonkin, MA. Changes in hand function following wrist arthrodesis in cerebral palsy. *J. Hand. Surg. Br.*, 2000, 25, 2, p. 193–194.

Harris, p., Riffle, K. Effects of inhibitive ankle foot orthoses on standing balance in a child with cerebral palsy. *Phys. Ther. J.*, 1986, 66, p. 663–667.

Hesse, p. Locomotor therapy in neurorehabilitation. *NeuroRehab.*, 2001, 16, p. 133 –139.

Holling, EE., Leviton, A. Characteristics of cranial ultrasound white matter echolucencies the predict disability: a review. *Dev. Med. Child Neurol.*, 1999, 41, p. 136–139.

Horbar, JD., Badger, GJ., Carpenter, JH., et al. Trends in Mortality and Morbidity for Very Low Birth Weight Infants, 1991–1999. *Pediatrics*, 2002, 110, 1, p. 143–151.

Hromádková, J., a kolektiv. *Fyzioterapie*. 1. vydání, Jinočany: H + H, s.r.o., 1999, p. 400–426, ISBN 80-86022-45-5.

Hupka, J. *Fyzikálna terapia*. Martin : Osveta, 1993, s. 48–49.

Hur, JJ. Review of research on therapeutic interventions for children with cerebral palsy. *Acta Neurologica Scandinavica*, 1995, 91, p. 423–432.

Hurvitz, EA., Leonard, Ch., Ayyangar, R., Nelson, VS. Complematary and alternative medicine use in families of children with cerebral palsy. *Dev. Med. Child Neurol.*, 2003, 45, p. 364–370.

Chauvel, P., Trottier, p., Wignal, JP., Bancaud, J. Somatomotor seizures of frontal lobe origin. In Chauvel P., Delgado-Escueta AV., Halgren E., Bancaud J. (eds.) *Frontal lobe epilepsies*. New York : Raven Press, 1992, p. 185–232.

Chvojka, J. *Magnetoterapie v teorii a praxi*. Professional Publishing, 2000, s. 50–53.

Ingram, TTS. *Paediatric aspects of cerebral palsy*. Edinburgh : E.p. Livingstone, 1964.

Ingram, TTS. The neurology of cerebral palsy. *Arch. Dis. Child.*, 1966, 41, p. 337–357.

Javůrek, J. *Fototerapie biolaserem*. Praha : Grada Publishing, 1995, s. 91–92.

Javůrek, J., Zounková, I. *Ošetřování a pohybový režim dětí s centrální koordinační poruchou*. 1. vyd., Praha : Národní centrum podpory zdraví, 1994.

Jenter, M., Lipton, GE., Miller, F. Operative treatment for hallux valgus in childern with cerebral palsy. *Foot Ankle. Int.*, 1998, 19, 12, p. 830–835.

Jeray, KJ., Rentz, J., Ferguson, RL. Local bone-graft technique for subtalar extraarticular arthrodesis in cerebral palsy. *J Pediatr. Orthop.*, 1998, 18, 1, p. 75–80.

Jessen, Ch., Mackie, P., Jarvis, p. Epidemiology of cerebral palsy. Letters to the editor. *Arch. Dis. Child. Fetal Neonatal Ed.*, 1999, 80, 2, p. F158–F160.

Jones, RF., Burke, D., Marosszeky, JE., Gillies, DJ. A new agent for the control of spasticity. *J. Neurol. Neurosurg. Psychiat.*, 1970, 33, p. 464–468.

Jozwiak, M., Marciniak, W., Piontek, T., Pietrzak, p. Dega´s transiliac osteotomy in the treatment of spastic hip subluxation and dislocation in cerebral palsy. *J. Pediatr. Orthop. Br.*, 2000, 9, 4, p. 257–264.

Kábele, F. *Rozvíjení hybnosti a řeči u dětí s DMO*. Praha : SPN, 1988.

Kaňovský, P., Streitová, H., Lněnička, J., Kubová, D., Bareš, M., Hortová, H. Indukce změny pohybového vzorce u dětí postižených dětskou mozkovou obrnou navozená současnou léčbou rehabilitační a botulotoxinem A. *Česká a slovenská neurologie a neurochirurgie*, 1999, 62, s. 203–208.

Kaňovský, P., Streitová, H., Lněnička, J., Hortová, H., Bareš, M., Kubová, D. Botulinum toxin A treatment of spasticity in non-ambulatory pediatric cerebral palsy. *Neurology*, 1999, 46 (suppl. 1), p. S86.

Keith, LG., Oleszczuk, JJ., Keith, DM. Multiple gestation: reflections on epidemiology, causes, and concequencep. *Int. J. Fertil. Womens Med.*, 2000, 45, 3, p. 206–214.

Kennedy, CR., Ayers, p., Campbell, MJ., et al. Randomized, Controlled Trial of Acetazolamide and Furosemide in Posthemorrhagic Ventricular Dilation in Infancy: Follow-Up at 1 year. *Pediatrics*, 2001, 108, 3, p. 597–607.

Kim, DS., Choi, JU., Yang, KH., Park, CI. Selective posterior rhizotomy in children with cerebral palsy: a 10-year experience. *Child's Nerv. Syst.*, 2001, 17, p. 556–562.

Klenková, J. *Možnosti stimulace preverbálních a verbálních schopností vývojově postižených dětí*. Brno : Paido, 2000.

Kolář, P. Posturální aktivita a DMO. Zdravotnické noviny, *Lékařské listy*, 2000, XLIX, 29, s. 1–2.

Kolář, P. Systematizace svalových dysbalancí z pohledu vývojové kineziologie. *Rehabilitace a fyzikální lékařství*, 2001, 4, s. 152–164.

Kolář, P. Význam posturální aktivity pro včasný záchyt pacientů s dětskou mozkovou obrnou. *Pediatrie pro praxi,* 2001, 4, s. 190–194.

Koman, LA., Mooney, JF., Smith, B., Goodman, A., Mulvaney, T. Management of cerebral palsy with botulinum-A toxin: Preliminary investigation. *Journal of Pediatric Orthopaedics*, 1993, 13, p. 489–495.

Koman, LA., Mooney, JF., Smith, B., Goodman, A., Mulvaney, T. Management of spasticity in cerebral palsy with botulinum-A toxin: report of a preliminary, randomised, double-blind trial. *Journal of Pediatrics Orthopaedics*, 1994, 14, p. 299–303.

Koukola, T., Satyaraj, E., Patel, DD., Tchernev, VT., Grimwade, BG., Kingsmore, SF., et al. Cerebral palsy is characterized by protein mediators in cord serum. *Ann. Neurol.*, 2004, 55, p. 186–194.

Kovačiková,V., Beranová, B. Postavení Vojtovy metody ve fyzioterapii hybných poruch. *Rehabilitácia*, 1998, XXXI, 2, s. 82–85.

Kovačiková,V., Beranová, B. Využití neuroplasticity v terapii pohybových poruch. *Rehabilitácia*, 1998, XXXI, 2, s. 81.

Kovačiková,V., Beranová, B. Kontakt fyzioterapeuta s pacientem. *Rehabilitácia*, 1998, XXXI, 2, s. 185–187.

Krageloh-Mann I., et al. Bilateral spastic cerebral palsy – MRI pathology and origin. Analysis from a representative series of 56 cases. *Dev. Med. Child Neurol.*, 1995, 38, p. 379–97.

Kuban, K., Sanocka, U., Leviton, A., et al. White matter disorders of prematurity: Association with intraventricular hemorrhage and ventriculomegaly. *J. Pediatr.*, 1999, 134, 5, p. 539–546.

Kulichová, J. et al. *Hipporehabilitace*. Praha : Nadace OF, 1995, s. 10–11; 26–27; 38.

Kumazaki, K., Nakayama, M., Sumida, Y., et al. Placental features in preterm infants with periventricular leukomalacia. *Pediatrics*, 2002, 109, 4, p. 650–655.

Kyllerman, M. Dyskinetic cerebral palsy. An analysis of 115 Swedish cases. *Neuropädiatrie*, 1977, 8, (Suppl.), p. S28–S32.

Kyllerman, M. Reduced optimality in pre- and perinatal conditions in dyskinetic cerebral palsy – distribution and comparison to control. *Neuropediatrics*, 1983, 14, p. 29–36

Kyllerman, M. The epidemiology of chronic neurologic diseases of children in Sweden. In French, JH., Harel, p., Casaer, P. (Eds.) *Child Neurology and Developmental Disabilities*. Baltimore : Paul Brookes, 1989, p. 137–143.

Inder, TE., Anderson, NJ., Spencer, C., et al. White matter injury in the premature infant: a comparison between serial cranial sonographic and MR findings at term. *Am. J. Neuroradiol.*, 2003, 24, p. 805–809.

Lehovský, M., Kraus, J. Monosynaptic H reflexes in Clinical Child Neurology. *Acta Univ. Carol.*, 1981, 27, 7/8, p. 399–466.

Lehovský, M., a kol. *Pečujeme o nervově nemocné dítě*. Praha : Avicenum, Zdravotnické nakladatelství, 08/31,08-040-89, vydání 1., 1989, s. 30–40

Lechta,V. *Symptomatické poruchy řeči u dětí*. Portál : Praha, 2002.

Letts, M., Rathbone, D., Yamashita, D., et al. Soft Boston orthosis in management of neuromuscular scoliosis: a preliminary report. *J. Pediatr. Orthop.*, 1992, 12, 4, p. 470–474.

Lubec, B., Dell´Anna, E., Fang-Kircher, p., et al. Decrease of Brain Protein Kinase C, Protein Kinase A, and Cyclin-Dependent Kinase Correlating with pH Precedes Neuronal Death in Neonatal Asphyxia. *J. Invest. Med.*, 1997, 45, p. 284–294.

Ludwig, G. Der Innenschuh, ein Hilfsmittel bei der Behandlung der Infantilen Zerebralparese. *Z. Krankengymn.*, 1977, 29, 1, p. 52–53.

Lundy, D.W., Ganey, T.M., Ogden, J.A., Guidera, K.J. Pathologic morphology of the dislocated proximal femur in children with cerebral palsy. *J. Pediatr. Orthop.*, 1998, 18, 4, p. 528–534.

MacLennan, A. A template for defining a causal relation between acute itrapartum events and cerebral palsy: international consensus statement. *BMJ*, 1999, 319, 7216, p. 1054–1059.

Mall, V., Heinen, F., Kirschner, J., Linder, et al. Evaluation of botulinum toxin A therapy in children with adductor spasm by gross motor function measure. *Journal of Child Neurology*, 2000, 15, p. 214–217.

Manning, J. Facilitation of movement – the Bobath approach. *Physiotherapy,* 1972, 58, 403–408.

Mayston, MJ., Barber, ChE., Stern, G., Bryce, J. The Bobath Concept – Will it stand the test of time?, *The Newsletter of British Association of Bobath trained Therapists*, 1997–1998, 28, p. 3–9.

Mayston, MJ. The Bobath Concept – Evolution and Application. *Med. Sport Sci.*, 1992, 36, p. 1–6.

Mayston, MJ. People with cerebral palsy: effects of and perspectives for therapy. *Neural. Plast.*, 2001, 8, p. 51–69.

Metaxiotis, D., Siebel, A., Doederlein, L. Repeated botulinum toxin A injections in the treatment of spastic equinus foot. *Clin Orthop.*, 2002, 394, p. 177–185.

Moreau, M., Cook, DC., Ashton, B. Adductor and psoas release for subluxation of the hip in children with spastic cerebral palsy. *J. Pediatr. Orthop.*, 1995, 15, 5, p. 672–676.

Mulier, T., Moens, P., Molenaers, G., Spaepen, D., Dereymaeker, G., Fabry, G. Split posterior tibial tendon transfer trough the interosseus membrane in spastic equinovarus deformity. *Foot Ankle. Int.*, 1995, 16, 12, p. 754–759.

Nelson, J., Ellenberg, J. Antecendens of cerebral palsy: multivariate analysis of risk. *N. Engl. J. Med.*, 1986, 315, p. 81–86.

Nelson, KB., Dambrosia, JM., Gerther, JK., Phillips, TM. Neonatal cytokines and coagulation factors in children with cerebral palsy. *Ann. Neurol.*, 1998, 44, p. 665–675.

Neubauer, K. Terapie dysartrie. In: Lechta,V., a kol.: *Terapia narušenej komunikačnej schopnosti.* Martin : Osveta, 2002.

Neville, B. Botulinum toxin in the cerebral palsies, grounds for cautious optimism. *British Medical Journal*, 1994, 309, p. 1526–1527.

Noorani, PA., Bodensteiner, JB., Barnes, PD. Colpocephaly. Frequency and associated findings. *Child. Neurol.*, 1988, 3, p.100–104.

O´Connor, AR., Stephenson, T., Johnson, A., et al. Long-term ophthalmic outcome of low birth weight children with and without retinopathy of prematurity. *Pediatrics*, 2002, 109, 1, p. 12–18.

Ohyu, J., Endo, A., Itoh, M., Takashima, p. Hypocapnia Under Hypotension Induces Apoptotic Neuronal Cell Death in Hippocampus of Newborn Rabbits. *Pediatr. Res.*, 2000, 48, 1, p. 24–29.

Oppenheim, WL. Selective posterior rhizotomy. *Clin. Orthop. Rel. Res.*, 1990; 253, p. 20–29.

O´Shea, MT., Klinepeter, KL., Goldstein, DJ., et al. Survival and Developmental Disability in Infants With Birth Weights of 501 to 800 Grams, Born Between 1979 and 1994. *Pediatrics*, 1997, 100, 6, p. 982–986.

O´Shea, TM., Preisser, JS., Klinepeter, KL., Dillard, RG. Trends in mortality and cerebral palsy in a geographically based cohort of very low birth weight neonates born between 1982 to 1994. *Pediatrics*, 1998, 101, 4, Pt 1, p. 642–647.

Paine, RP. Minimal Chronic Brain Syndromes. *Dev. Med. Child Neurol.*, 21, 4, 1962, cit. Brázdilová (1996).

Park, TS., Gaffney, PE., Kaufman, BA., Molleston, MC. Selective lumbosacral dorsal rhizotomy immediately caudal to the conus medullaris for cerebral palsy spasticity. *Neurosurgery*, 1993, 33, p. 929–934.

Park, TS., Owen, JH. Surgical management of spastic diplegia in cerebral palsy. *N. Engl. J. Med.*, 1992, 326, p. 745–749.

Pavlů, D. *Speciální fyzioterapeutické koncepty a metody.* Akademické nakladatelství CERM, s.r.o., Brno, 2002, ISBN 80-7204-266-1.

Pavlů, D., Véle, F., Havlíčková, L. Elektromyografická a kineziologická analýza Vojtova terapeutického principu. *Rehabilitace a fyzikální lékařství*, Česká lékařská společnost J. E. Purkyně, 2000, 7, 2, p.74–77.

Peacock, WJ., Arens, LJ. Selective dorsal rhizotomy for relief of spasticity in cerebral palsy.p. *Afr. Med. J.*, 1982, 62, p. 119–124.

Peacock, WJ., Arens, LJ., Berman, B. Cerebral palsy spasticity. Selective posterior rhizotomy. *Pediatr. Neurosci.*, 1987, 13, p. 61–66.

Peacock, WJ., Staudt, LA. Functional outcomes following selective posterior rhizotomy in children with cerebral palsy. *J. Neurosurg.*, 1991, 74, p. 380–385.

Pellicer, A., Gaya, F., Quero, MR., Cabans, F. Noninvasive continuous monitoring of the effects of head position on brain hemodynamics in ventilated infants. *Pediatrics*, 2002, 109, 3, p. 434–440.

Penn, RD., Kroin, JS. Intrathecal baclofen alleviates spinal cord spasticity. *Lancet* 1984, p. 1078.

Perlman, JM. White matter injury in the preterm infant: an important determination of abnormal neurodevelopment outcome. *Early Hum. Dev.*, 1998, 53, 2, p. 99–120.

Petäjä, J., Hiltunen, L., Fellman, V. Increased Risk of Intraventricular Hemorrhage in Preterm Infants with Thrombophilia. *Pediatr. Res.*, 2001, 49, 5, p. 643–646.

Pharoah, POD., Platt, MJ., Cooke, T. The changing epidemiology of cerebral palsy. *Arch. Dis. Child Fetal Neonatal Ed,* 1996, 75, p. F169–F173.

Polin, RA., Spitzer, AR. *Fetal and Neonatal Secrets.* Hanley and Belfus Inc., 2001, p. 324–353.

Prechtl, HF., et. al. An early marker for neurological deficits after perinatal brain lesions. *Lancet*, 1997, 349, p. 1361–1363.

Rayan, GM.,Young, BT. Arthrodesis of the spastic wrist. *J. Hand. Surg.*, 1999, 24, 5, p. 944–952.

Rennie, JM., Roberton, NRC. *Neurological Problems. A Manual of Neonatal Intensive Care*, 4th ed., Arnold, 2002, p. 311–357.

Robert, R., Young, MD. Spasticity: a review. *Neurology*, 1994, 44, 9, p. S12-S20.

Russel, DJ., Rosenbaum, PL., Cadman, DT., Gowland, C., Hardy, p., Jarvis, p. The gross motor function measure: a means to evaluate the effects of physical therapy. *Dev. Med. Child Neurol.*, 1989, 31, p. 341–352 .

Sala, DA., Grant, AD. Prognosis for ambulation in cerebral palsy. *Dev. Med. Child Neurol.*, 1995, 37, p. 1020–1026.

Salokorpi, T., Rajantie, I., Viitala, J., et al. Does perinatal hypocarbia play a role in the pathogenesis of cerebral palsy? *Acta Paediatr.*, 1999, 88, 5, p. 571–575.

Sanchéz-Alvares, R., Almeida, A., Medina, JM. Oxidative Stress in Preterm Rat Brain Is Due to Mitochondrial Dysfunction. *Pediatr. Res.*, 2002, 51, 1, p. 34–39.

Saugstad, OD. Mechanisms of tissue injury by oxygen radicals: implications for neonatal disease. *Acta Paediatr.*, 1996, 85, p. 1–4.

Saugstad, OD. Protection of the fetal/infant brain: free radicals and antioxidants. In Cockburn F. (Ed.) *Advances in perinatal medicine*. The Parthenon Publishing Group, 1997, p. 178–182.

Seidl, Z., Süssová, J., Obenberger, J. Komplexní hodnocení dětí s DMO. Možnosti prognózy dětí s dětskou mozkovou obrnou (DMO) na základě komplexního vyšetření. *Rehabilitace a fyzikální lékařství*, 1995, 2, 1, s. 16–18.

Seidl, Z., Süssová, J., Obenberger, J., Vaněčková, M., Viták, T., Rydland, J. Magnetic resonance imaging in diplegic form of cerebral palsy. *Brain & Development,* 2001, 23, p. 46–49.

Seidl, Z., Süssová, J. Postižení mozku v obraze počítačové tomografie s dětskou mozkovou obrnou (soubor 80 dětí). *Československá pediatrie,* 1990, 45, s. 168–169.

Shi, Y., Pan, F., Li, H., et al. Role of Carbon Monoxide and Nitric Oxide in Newborn Infants With Postasphyxial Hypoxic-Ischemic Encephalopathy. *Pediatrics*, 2000, 106, 6, p. 1447–1451.

Sherington, CS. *The intergrative function of the nervous system.* Cambridge : University Press, 1909, p. 433.

Sherrington, CS. Decerebrate rigidity and reflex coordination of movements. *J. Physiol.* (Lond.), 1898, 22, p. 319–337.

Shumway-Cook, A., Woollacott, M.H. *Motor Control, Theory and Practical Applications*, Maryland, USA, 1995, p. 7–13.

Schejbalová, A., Frýdl, J., Mrzena, V., Popelka, p. Operační léčení pacientů s DMO na Ortopedické klinice FDL UK Praha – Motol. *Acta Chir. Orthop. Traum. Čech.*, 1990, 57, 1, s. 27–31.

Schejbalová, A., Smetana,V. Musculus iliopsoas – jedna z příčin neurogenní luxace: vlastní technika tenotomie a prolongace. *Acta Chir. Orthop. Traum. Čech.*, 2000, 67, 4, s. 234–238.

Schejbalová, A., Smetana,V. Transpozice a hemitranspozice musculus tibialis anterior při řešení pes equinovarus. *Acta Chir. Orthop. Traum. Čech.*, 2001, 68, 4, p. 239–243.

Schejbalová, A.: Moderní trendy v ortopedické terapii dětské mozkové obrny. *Čes.-slov. Pediat.*, 2003, 58, 9, s. 578–585.

Smetana, V. *Dětská mozková obrna z pohledu ortopéda - operační léčení projevů DMO na dolních končetinách.* (Habilitační práce) Praha, 1993, s. 215. Univerzita Karlova, 2. Lékařská fakulta.

Smetana, V., Schejbalová, A. Ortopedické operační léčení dětí postižených dětskou mozkovou obrnou. *Čes.-slov. Pediat.*, 1993, 48, 11, s. 670–674.

Smetana, V., Schejbalová, A. Operace podle Strayera jako základní operace k ovlivnění pes equinus u dětské mozkové obrny. *Acta. Chir. Orthop. Traum. Čech.*, 1993, 60, 4, s. 218–224.

Smetana, V., Schejbalová, A. Genu flectum u dětské mozkové obrny (DMO), prolongace flexorů kolenního kloubu s uvedením vlastní modifikace. *Acta Chir. Orthop. Traum. Čech.*, 1993, 60, 4, s. 225–231.

Smetana, V., Schejbalová, A. Význam tenotomie adduktorů při léčení projevů dětské mozkové obrny na dolních končetinách. *Acta Chir. Orthop. Traum. Čech.*, 1993, 60, 5, s. 301–305.

Smetana, V., Schejbalová, A. Naše zkušenosti s operací podle Griceho u pacientů postižených dětskou mozkovou obrnou. *Acta Chir. Orthop. Traum. Čech.*, 1994, 61, 1, s. 34–38.

Smetana, V., Schejbalová, A. Naše zkušenosti s operací dle Stephensona a Donovana při řešení addukční kontraktury u dětské mozkové obrny. *Acta Chir. Orthop. Traum. Čech.*, 1994, 61, 3, s. 167–170.

Smetana, V., Schejbalová, A. Kombinace operací podle Younga a Griceho při řešení pes planovalgus u dětí s dětskou mozkovou obrnou. *Acta Chir. Orthop. Traum. Čech.*, 1997, 64, 2, s. 86–89.

Smetana, V., Schejbalová A. Možnosti operačního řešení hallux valgus u pacientů s dětskou mozkovou obrnou. *Acta Chir. Orthop. Traum. Čech.*, 1994, 61, 6, s. 341–343.

Song, HR., Carroll, NC. Femoral varus derotation osteotomy with or without acetabuloplasty for unstable hips in cerebral palsy. *J. Pediatr. Orthop.*, 1998, 18,1, p. 62–68.

Speidl, B., Fleming, P., Henderson, J., et al. *Assessment of outcome and care after discharge. A Neonatal Vade Mecum*, 3rd ed., Arnold, 1998, p. 194–215.

Spira, R. Management of spasticity in cerebral palsied children by peripheral nerve block with phenol. *Dev. Med. Child Neurol.*, 1971, 13, p. 164–173.

Snyder, EY., Cloherty, JP. Perinatal Asphyxia. In Cloherty, JP. and Stark, AR. (Eds.) *Manual of Neonatal Care*, 4th ed. Eds. Lippincott-Raven, 1998, p. 515–533.

Stehlík, A. a kol. *Dítě s mozkovou obrnou v rodině*. Praha : Avicenum, 1977.

Steinbok, P., Schrag. Ch. Complications after selective posterior rhizotomy for spasticity in children with cerebral palsy, *Pediatr. Neurosurg.*, 1998, 28, p. 300–313

Steinbok, P. Outcomes after selective dorsal rhizotomy for spastic cerebral palsy. *Childs Nerv. Syst.*, 2001, 17, p. 1–18.

Steinwender, G., Saraph, V., Zwick, EB., Uitz, C., Linhart, W. Fixed and dynamic equinus in cerebral palsy evaluation of ankle function after multilevel surgery. *J. Pediatr. Orthop.*, 2001, 21, 1, p. 102–107.

Stewart, A. Outcome. In Levitt, G., Harvey, D., Cooke R. (Eds.) *Practical Perinatal Care: The Baby under1000 g.* Butterwirth Heinemann, 1999, p. 312–323.

Süssová, J., Seidl, Z., Faber, J. Strana postižení u hemiparetické formy DMO ve vztahu k výskytu epilepsie. *Rehablitace*, 1984, 17, s. 201–208.

Süssová, J., Seidl, Z., Obenberger, J. Relation of squinting and dilated occipital horns of lateral ventricles in hemiparetic form of cerebral palsy. *Sborn. lék.*, 1996, 97, 2, p. 203–207.

Süssová, J., Seidl, Z., Faber, J. Hemiparetic form of cerebral palsy in relation to epilepsy and mental retardation. *Dev. Med. Child Neurol.*, 1990, 32, p. 792–795.

Škapík, M., et al. *Využití balneoterapie ve vnitřním lékařství*. Praha : Grada Publishing, 1994, s. 138.

Škodová, E., Jedlička, I. *Klinická logopedie*. Praha : Portál, 2003.

Tachdjian MO. *Pediatric Orthopedics*. 2nd ed. Philadelphia, London, Toronto : Saunders, 1990, p. 3366.

Tenuta, J., Shelton, Y.A., Miller, F. Long- term follow – up of triple arthrodesis in patients with cerebral palsy. *J. Pediatr Orthop.*, 1993, 13, 6, p. 713–716.

Terjesen, T., Hellum, C. Femoral shortening osteotomy for chronic hip dislocation in patients with cerebral palsy. *Tidsskr. Nor. Laegeforen*, 1998, 118, 18, p. 2773–2776.

The United Kingdom Tizanidine Study Group. A double-blind placebo-controlled trial of tizanidine in the treatment of spasticity caused by multiple sclerosis. *Neurology*, 1994, 44 (suppl. 9), p. 70–79.

Thom, H. *Die infantilen Zerebralparesen.* 2 Auflage. Stuttgart, New York : Georg Thieme Verlag, 1982, p. 546.

Thomson, JD. Banta, JV. Scoliosis in cerebral palsy: an overview and recent results. *J. Pediatr. Orthop. B.*, 2001, 10, 1, p. 6–9.

Ticháčková, J. *Plavání tělesně postižených – míšní léze, DMO, amputace končetin.* (Absolventská práce) Brno, SZŠ a VOŠ Merhautova, 2002, s. 22–31.

Tichý, M., Kraus, J., Hořínek, D., Vaculík, M. Selective posterior rhizotomy in treatment of cerebral palsy, first experience in Czech Republic. *Bratisl. Lek. Listy*, 2003, 104, 2, p. 54–58.

Tichý, M., Kraus, J., Hořínek, D., Vaculík, M. Selektivní zadní rhizotomie v léčbě těžké formy spasticity u dětské mozkové obrny. *Bolest*, 2004, 7,1, s. 23–26.

Tommiska, V., Heinonen, K., Ikonen, p., et al. A National Short-Term Follow-Up Study of Extremely Low Birth Weight Infants Born in Finland in 1996–1997. *Pediatrics*, 2001, 107, 1, p. E2.

Tonkin, MA., Hatrick, NC., Eckersley, JR., Couzens, G. Surgery for cerebral palsy part 3: classification and operative procedures for tumb deformity. *J. Hand. Surg. Br.*, 2001, 26, 5, p. 465–470.

Topp, M., Uldall, P., Greisen, G. Cerebral plasy birth in eastrn Denmark, 1987–90: implications for neonatal care. *Paediatr. Perinat. Epidemiol.*, 2001, 15, 3, p. 271–277.

Trojan, p., et al. *Fyziologie a léčebná rehabilitace motoriky člověka.* Praha : Grada Publishing, 1996, s. 141–143.

Ubhi, T., Bhakta, BB., Ives, HL., Allgar, V., Roussonis, SH. Randomised double blind placebo controlled trial of the effect of botulinum toxin on walking in cerebral palsy. *Arch. Dis. Child.*, 2000, 83, p. 481–487.

Usanova, O. N. Efektivní diagnostické postupy a techniky nápravy používané ve školní, poradenské a klinické psychologii při určování a nápravě neuropsychologických potíží u dětí. Texty pro účastníky mezinárodního semináře. Praha, 2001.

ÚZIS ČR. *Rodička a novorozenec 2000. Zpráva o novorozenci.* Praha : ÚZIS ČR, 2001.

Van Hest, AE., House, JH., Cariello, C. Upper extremity surgical treatment of cerebral palsy. *J. Hand. Surg. Am.*, 1999, 24, 2, p. 323–330.

Vannucci, RC., Towfighi, J., Heitjan, DF., Brucklacher, RM. Carbon Dioxide Protects the Perinatal Brain From Hypoxic-Ischemic Damage: An Experimental Study in the Immature Rat. *Pediatrics*, 1995, 95, 6, p. 868–874.

Vater,W., Boudzio, M. *Od prvního hlasu k prvním slovům.* Tech-market, Praha 1996

Véle, F. *Kineziologie pro klinickou praxi.* Praha : Grada Publishing, 1997, p. 85, ISBN 80-7169-256-5.

Véle, F. Význam centrální nervové soustavy pro reedukaci pohybu (pohyb a jeho terapeutické ovlivnění z pohledu klinické neurofyziologie). Sborník přednášek, VIII. Novoměstské dny, Okresní nemocnice v Novém Městě na Moravě, květen 2000, s. 6–12.

Veličkovic, T. Basic Principles of the Neurodevelopmental Treatment (NDT) – Bobath, *Pediatr. Croat.*, 2002, 46 (Supl 1), p. 163–168 .

Vlach, O. *Léčení deformit páteře.* 1. vyd., Praha : Avicenum, 1986, s. 214.

Vodičková, R. Synergetická reflexní terapie. *Bulletin, Unie fyzioterapeutů ČR*, 1999, 10, s. 25–30.

Vohr, BR., Wright, LL., Dusick, AM., et al. Neurodevelopmental and functional outcomes of extremely low birth weight infants in the National Institute of Child Health and Human Development Neonatal Research Network, 1993–1994. *Pediatrics*, 2000, 105, p. 1216–1226.

Vojta,V. *Mozkové hybné poruchy v kojeneckém věku*. Praha : Nakladatelství Grada Avicenum, 1993, ISBN 80-85424-98-3.

Vojta, V., Peters, A. *Vojtův princip*. Praha: Grada Publishing, 1995, ISBN 80-7169-004-X.

Vojta,V., Peters, A. *Das Vojta-Princip*. Berlin Heidelberg New York : Springer Verlag, 2001, ISBN 3-540-60576-2.

Vojtova metoda, prof. MUDr. Václav Vojta, stránky stažené z internetové adresy: http://www.rl-corpus.cz

Volkert, R. Individuelle ortopädisch-technische Hilfsmittel zur Funktionsverbesserung der behinderten Hand. *Orthop. Tech.*, 1978, 2, 1, S. 13.

Volkert, R. Individuelle ortopädisch-technische Versorgungen bei Kindern mit Infantilen Zerebralparesen. *Z. Krankengymn.*, 1980, 32, 4, S. 389–394.

Volpe, JJ. Hypoxic-Ischemic Encephalopathy: Biochemical and Physiological Aspects. In VOLPE, JJ. *Neurology of the Newborn*, 4th ed., W.B. Saunders Comp., 2001, p. 217––276.

Volpe, JJ. Hypoxic-Ischemic Encephalopathy: Clinical Aspects. In VOLPE, JJ. *Neurology of the Newborn*, 4th ed. W.B.Saunders Comp., 2001, p. 331–394.

Volpe, JJ. Neurobiology of Periventricular Leukomalacia in the Premature Infant. Review Articles. *Pediatr. Res.*, 2001, 50, 5, p. 553–562.

von Koch, CS., Park, TS., Steinbok, P., Smyth, M., Peacock, WJ. Selective posterior rhizotomy and intrathecal baclofen for treatment of spasticity. *Pediatr. Neurosurg.*, 2001, 35, 2, p. 57–65.

Wade, DT. *Measurement in neurological rehabilitation*, New York : Oxford Univ. Press, First publ., 1992, repr. 93–96, 98, 2000.

Wall, SA., Chait, LA., Temlett, JA., Perkins, B., Hillen, G., Becker, P. Botulinum A chemodenervation: a new modality in cerebral palsy hands. *British Journal of Plastic Surgery*, 1993, 46, p. 703–706.

Weindling, AM., et al. A randomized controlled trial of eatly physiotherapy for high-risk infants. *Acta Paediatr.*, 1996, 85, p. 1107–11.

World Health Organization: Save the Childern, 2001.

Worster-Drought, C. Suprabulbar paresis. Suppl. 30. *Dev. Med. Child Neurol.*, 1974, p. 16.

Yoon, BH., Romero, R., Park, JS., et al. Fetal exposure to an intraamniotic inflammation and the development of cerebral palsy at the age of three years. *Am. J. Obstet. Gynecol.*, 2000, 182, p. 675–681.

Zoban, P. Antepartální a intrapartální rizikové faktory novorozenecké encefalopatie. Komentář. *Gynekologie po promoci*, 2001, 1, 6, p. 17–21.

Zoban, P., Černý, M. Hypoxické postižení mozku. In *Vybrané kapitoly z perinatologie*. Fuchs, V., Zoban, P., Tomášová, H., Černý, M. Praha : Karolinum, 2001, s. 288–295.

Zoban, P. *Pozdní morbidita novorozenců, vývoj v ČR*. XIX. konference Sekce perinatální medicíny České gynekologicko-porodnické společnosti ČLS JEP, Olomouc, 2002.

Zounková, I. Fyzioterapeutem získané klinické zkušenosti při aplikaci metody Vojtovy a K. a B. Bobathových. Zdravotnické noviny, *Lékařské listy*, 2000, XLIX, 29, s. 6–7.