

References

- Abel, O. (1918). Das Entwicklungstempo der Wirbeltierstämme. *Vereines zur Verbreitung Naturwissenschaftlicher Kenntnisse, Schriften*, Wien, 58, 91–120.
- Ager, D. V. (1976). The nature of the fossil record. *Proceedings of the Geologists' Association, London*, 87, 131–159.
- Aguirre, W. E. and Bell, M. A. (2012). Twenty years of body shape evolution in a threespine stickleback population adapting to a lake environment. *Biological Journal of the Linnean Society*, 105, 817–831.
- Aguirre, W. E., Doherty, P. K., and Bell, M. A. (2004). Genetics of lateral plate and gillraker phenotypes in a rapidly evolving population of threespine stickleback. *Behaviour*, 141, 1465–1483.
- Ashton, E. H. and Zuckerman, S. (1950). The influence of geographic isolation on the skull of the green monkey (*Cercopithecus aethiops sabacus*). I. A comparison between the teeth of the St. Kitts and the African Green monkey. *Proceedings of the Royal Society of London, Series B*, 137, 212–238.
- Ayala, F. J., Serra, L. and Prevosti, A. (1989). A grand experiment in evolution: the *Drosophila subobscura* colonization of the Americas. *Genome*, 31, 246–255.
- Babin-Fenske, J., Anand, M., and Alarie, Y. (2008). Rapid morphological change in stream beetle museum specimens correlates with climate change. *Ecological Entomology*, 33, 646–651.
- Bader, R. S. (1955). Variability and evolutionary rate in the oreodonts. *Evolution*, 9, 119–140.
- Baer, K. E. v. (1876). *Reden Gehalten in Wissenschaftlichen Versammlungen Und Kleinere Aufsätze Vermischten Inhalts. Zweiter Theil: Studien Aus Dem Gebiete Der Naturwissenschaften*. St. Petersburg, Schmitzdorff.
- Bailey, D. W. (1959). Rates of subline divergence in highly inbred strains of mice. *Journal of Heredity*, 50, 26–30.
- Baker, A. J. and Marshall, H. D. (1999). Population divergence in chaffinches *Fringilla coelebs* assessed with control-region sequences. In *Proceedings of the 22nd International Ornithological Congress, Durban*, eds. N. J. Adams and R. H. Slotow, Johannesburg, BirdLife South Africa, pp. 1899–1913.
- Baker, A. J., Peck, M. K., and Goldsmith, M. A. (1990). Genetic and morphometric differentiation in introduced populations of common chaffinches (*Fringilla coelebs*) in New Zealand. *Condor*, 92, 76–88.
- Baker, J. A., Heins, D. C., Foster, S. A., and King, R. W. (2008). An overview of life-history variation in female threespine stickleback. *Behaviour*, 145, 579–602.

- Banerjee, M. K. and Basu, A. (1991). Uttar Pradesh: basic anthropometric data. *All India Anthropometric Survey, North Zone (Anthropological Survey of India, Calcutta)*, 10, 1–626.
- Barrett, P. H., Gautrey, P. J., Herbert, S., Kohn, D., and Smith, S. (1987). *Charles Darwin's Notebooks, 1836–1844: Geology, Transmutation of Species, Metaphysical Enquiries*. Cambridge, British Museum (Natural History) and Cambridge University Press.
- Basu, A., Ganguly, P., Ghosh, G. C., and Basu, S. K. (1989). Maharashtra: basic anthropometric data. *All India Anthropometric Survey, North Zone (Anthropological Survey of India, Calcutta)*, 4, 1–550.
- Bather, F. A. (1920). Fossils and life. *Reports of the British Association for the Advancement of Science, London*, 1920, 61–86.
- Bell, M. A. and Aguirre, W. E. (2013). Contemporary evolution, allelic recycling, and adaptive radiation of the threespine stickleback. *Evolutionary Ecology Research*, 15, 377–411.
- Bell, M. A., Aguirre, W. E., and Buck, N. J. (2004). Twelve years of contemporary armor evolution in a threespine stickleback population. *Evolution*, 58, 814–824.
- Bell, M. A., Baumgartner, J. V., and Olson, E. C. (1985). Patterns of temporal change in single morphological characters of a Miocene stickleback fish. *Paleobiology*, 11, 258–271.
- Bell, M. A. and Haglund, T. R. (1982). Fine-scale temporal variation of the Miocene stickleback *Gasterosteus doryssus*. *Paleobiology*, 8, 282–292.
- Bell, M. A., Travis, M. P. and Blouw, D. M. (2006). Inferring natural selection in a fossil threespine stickleback. *Paleobiology*, 32, 562–577.
- Benson, R. B. J., Campione, N. E., Carrano, M. T., Mannion, P. D., Sullivan, C., Upchurch, P., and Evans, D. C. (2014). Rates of dinosaur body mass evolution indicate 170 Million years of sustained ecological innovation on the avian stem lineage. *PLoS Biology*, 12, e1001853.
- Berg, H. C. (1983). *Random Walks in Biology*. Princeton, Princeton University Press.
- Berg, L. S. (1926). *Nomogenesis, or Evolution Determined by Law*. London, Constable and Company.
- Berger, J. (1986). *Wild Horses of the Great Basin: Social Competition and Population Size*. Chicago, University of Chicago Press.
- Berry, R. J. (1964). The evolution of an island population of the house mouse. *Evolution*, 18, 468–483.
- Beurlen, K. (1932). Funktion und Form in der organischen Entwicklung. *Naturwissenschaften, Berlin*, 20, 73–80.
- Blanckenhorn, W. U. (2015a). Investigating yellow dung fly body size evolution in the field: Response to climate change? *Evolution*, 69, 2227–2234.
- Blanckenhorn, W. U. (2015b). Data from: Investigating yellow dung fly body size evolution in the field: response to climate change? *Dryad Digital Repository*, <https://doi.org/10.5061/dryad.3v4hn>.
- Boag, P. T. and Grant, P. R. (1981). Intense natural selection in a population of Darwin's finches (Geospizinae) in the Galapagos. *Science*, 214, 82–85.
- Bonner, J. T. (1965). *Size and Cycle: An Essay on the Structure of Biology*. Princeton, Princeton University Press.
- Bookstein, F. L. (1987). Random walk and the existence of evolutionary rates. *Paleobiology*, 13, 446–464.
- Bookstein, F. L., Gingerich, P. D., and Kluge, A. G. (1978). Hierarchical linear modeling of the tempo and mode of evolution. *Paleobiology*, 4, 120–134.
- Bowles, G. T. (1932). *New Types of Old Americans at Harvard and at Eastern Women's Colleges*. Cambridge, Harvard University Press.

- Brett, C. E. and Baird, G. C. (1995). Coordinated stasis and evolutionary ecology of Silurian to Middle Devonian faunas in the Appalachian Basin. In *New Approaches to Speciation in the Fossil Record*, eds. D. H. Erwin and R. L. Anstey, New York, Columbia University Press, pp. 285–315.
- Brinkmann, R. (1929). Statistisch-biostratigraphische Untersuchungen an mittelmajurassischen Ammoniten über Artbegriff und Stammesentwicklung. *Abhandlungen der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse, Neue Folge*, 13 (3), 1–249.
- Brown, C. R. and Brown, M. B. (1998a). Intense natural selection on body size and wing and tail asymmetry in cliff swallows during severe weather. *Evolution*, 52, 1461–1475.
- Brown, C. R. and Brown, M. B. (1998b). Fitness components associated with alternative reproductive tactics in cliff swallows. *Behavioral Ecology*, 9, 158–171.
- Brown, M. B. and Brown, C. R. (2011). Intense natural selection on morphology of cliff swallows (*Petrochelidon pyrrhonota*) a decade later: did the population move between adaptive peaks? *Auk*, 128, 69–77.
- Bruce, R. C. (1988). An ecological life table for the salamander *Eurycea wilderae*. *Copeia*, 1988, 15–26.
- Buckman, S. S. (1909). *Yorkshire Type Ammonites, Volume 1, Part 1*. London, Wheldon and Wesley.
- Bulmer, M. G. (1977). Periodical insects. *American Naturalist*, 111, 1099–1117.
- Bumpus, H. C. (1899). The elimination of the unfit as illustrated by the introduced sparrow, *Passer domesticus*. *Biological Lectures Delivered at the Marine Biology Laboratory, Woods Hole*, 1898, 209–226.
- Butler, M. A. and King, A. A. (2004). Phylogenetic comparative analysis: a modeling approach for adaptive evolution. *American Naturalist*, 164, 683–695.
- Calder, W. A. (1984). *Size, Function, and Life History*. Cambridge, Massachusetts, Harvard University Press.
- Calhoun, J. B. (1947). The role of temperature and natural selection in relation to the variations in the size of the English sparrow in the United States. *American Naturalist*, 81, 203–228.
- Carroll, S. P. and Boyd, C. (1992). Host race radiation in the soapberry bug: natural history with the history. *Evolution*, 46, 1052–1069.
- Carroll, S. P., Dingle, H., and Klassen, S. (1997). Genetic differentiation of fitness-associated traits among rapidly evolving populations of the soapberry bug. *Evolution*, 51, 1182–1188.
- Carroll, S. P., Hendry, A. P., Reznick, D. N., and Fox, C. W. (2007). Evolution on ecological time-scales. *Functional Ecology*, 21, 387–393.
- Carson, H. L. (1975). The genetics of speciation at the diploid level. *American Naturalist*, 109, 83–92.
- Carson, H. L. (1987). Population genetics, evolutionary rates and Neo-Darwinism. In *Rates of Evolution*, eds. K. S. W. Campbell and M. F. Day, London, Allen and Unwin, pp. 209–217.
- Caruso, N. M., Sears, M. W., Adams, D. C., and Lips, K. R. (2014). Widespread rapid reductions in body size of adult salamanders in response to climate change. *Global Change Biology*, 20, 1751–1759.
- Cavalli-Sforza, L. L. and Bodmer, W. F. (1971). *The Genetics of Human Populations*. San Francisco, W. H. Freeman and Company.
- Chaline, J. (1984). Le concept d'évolution polyphasée et ses implications. *Geobios, Lyon*, 17, 783–795.

- Charlesworth, B. (1980). *Evolution in Age-Structured Populations*. Cambridge, Cambridge University Press.
- Charlesworth, B. (1984). Some quantitative methods for studying evolutionary patterns in single characters. *Paleobiology*, 10, 308–318.
- Charmantier, A., Kruuk, L. E. B., Blondel, J., and Lambrechts, M. M. (2004). Testing for microevolution in body size in three blue tit populations. *Journal of Evolutionary Biology*, 17, 732–743.
- Cheetham, A. H. (1968). Morphology and systematics of the bryozoan genus *Metrarabdotos*. *Smithsonian Miscellaneous Collections*, 153, 1–121.
- Cheetham, A. H. (1986). Tempo of evolution in a Neogene bryozoan: rates of morphologic change within and across species boundaries. *Paleobiology*, 12, 190–202.
- Cheetham, A. H. and Jackson, J. B. C. (1995). Process from pattern: tests for selection versus random change in punctuated bryozoan speciation. In *New Approaches to Speciation in the Fossil Record*, eds. D. H. Erwin and R. L. Anstey, New York, Columbia University Press, pp. 184–207.
- Cheetham, A. H., Sanner, J. and Jackson, J. B. C. (2007). *Metrarabdotos* and related genera (Bryozoa: Cheilostomata) in the late Paleogene and Neogene of tropical America. *Journal of Paleontology* 81(1), Supplement, *Paleontological Society Memoir*, 67, 1–96.
- Chiba, S. (1996). A 40,000-year record of discontinuous evolution of island snails. *Paleobiology*, 22, 177–188.
- Chiba, S. (2007). Morphological and ecological shifts in a land snail caused by the impact of an introduced predator. *Ecological Research*, 22, 884–891.
- Clark, D. B. (1980). Population ecology of *Rattus rattus* across a desert-montane forest gradient in the Galápagos Islands. *Ecology*, 61, 1422–1433.
- Clutton-Brock, T. H., Guinness, F. E., and Albon, S. D. (1982). *Red Deer: Behaviour and Ecology of Two Sexes*. Chicago, University of Chicago Press.
- Clutton-Brock, T. H. and Pemberton, J. M. eds. (2004). *Soay Sheep: Dynamics and Selection in an Island Population*, Cambridge, Cambridge University Press.
- Clyde, W. C. and Gingerich, P. D. (1994). Rates of evolution in the dentition of early Eocene *Cantius*: comparison of size and shape. *Paleobiology*, 20, 506–522.
- Clyde, W. C., Hamzi, W., Finarelli, J. A., Wing, S. L., Schankler, D. M., and Chew, A. (2007). Basin-wide magnetostratigraphic framework for the Bighorn Basin, Wyoming. *Geological Society of America Bulletin*, 119, 848–859.
- Colbert, E. H. (1948). Evolution of the horned dinosaurs. *Evolution*, 2, 145–163.
- Conant, S. (1988). Geographic variation in the Laysan finch (*Telespyza cantans*). *Evolutionary Ecology*, 2, 270–282.
- Conover, D. O. and Munch, S. B. (2002). Sustaining fisheries yields over evolutionary time scales. *Science*, 297, 94–96.
- Cooch, E. G., Lank, D. B., Rockwell, R. F., and Cooke, F. (1991). Long-term decline in body size in a snow goose population: evidence of environmental degradation? *Journal of Animal Ecology*, 60, 483–496.
- Cooke, F., Rockwell, R. F., and Lank, D. B. (1995). *The Snow Geese of La Pérouse Bay: Natural Selection in the Wild*. Oxford, Oxford University Press.
- Cooper, G. A. and Williams, A. (1952). Significance of the stratigraphic distribution of brachiopods. *Journal of Paleontology*, 26, 326–337.
- Cope, E. D. (1887). *The Origin of the Fittest, Essays on Evolution*. New York, D. Appleton and Co. 467.
- Cope, E. D. (1896). *The Primary Factors of Organic Evolution*. Chicago, Open Court.
- COSEWIC. (2012). COSEWIC assessment and status report on the grizzly bear *Ursus arctos*. *Committee on the Status of Endangered Wildlife in Canada, Ottawa*, 1–84.

- COSEWIC. (2013a). COSEWIC assessment and status report on the Plains Bison *Bison bison bison* and the Wood Bison *Bison bison athabasca* in Canada. *Committee on the Status of Endangered Wildlife in Canada*, Ottawa, 1–109.
- COSEWIC. (2013b). COSEWIC assessment and status report on the giant threespine stickleback *Gasterosteus aculeatus* and the unarmoured threespine stickleback *Gasterosteus aculeatus* in Canada. *Committee on the Status of Endangered Wildlife in Canada*, Ottawa, 1–62.
- Coulson, T. and Crawley, M. J. (2004). Appendix 3: How average life tables can mislead. In *Soay Sheep: Dynamics and Selection in an Island Population*, eds. T. H. Clutton-Brock and J. M. Pemberton, Cambridge, Cambridge University Press, pp. 328–331.
- Crow, J. F. and Kimura, M. (1979). Efficiency of truncation selection. *Proceedings of the National Academy of Sciences USA*, 76, 396–399.
- Curry, G. B. (1982). Ecology and population structure of the recent brachiopod *Terebratulina* from Scotland. *Palaeontology*, 25, 227–246.
- Cuvier, G. (1812). *Recherches sur les ossements fossiles de quadrupèdes. Tomes I-IV*. Paris,
- Cuvier, G. (1817). *Le règne animal, distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée, Tome I*. Paris, Deterville.
- Cuvier, G. (1825). *Discours sur les révolutions de la surface du globe, et sur les changements qu'elles ont produits dans le règne animal*. Dufour et d'Ocagne, Paris (reprinted 1969 by Culture et Civilisation, 115 Avenue Gabriel Lebon, Brussels) 400.
- D'Agostino, R. B. (1986). Tests for the normal distribution. In *Goodness-of-Fit Techniques*, eds. R. B. D'Agostino and M. A. Stephens, New York, Marcel Dekker, pp. 367–419.
- Dall, W. H. (1877). On a provisional hypothesis of saltatory evolution. *American Naturalist*, 11, 135–137.
- Darwin, C. R. (1839). *Journal of Researches into the Geology and Natural History of the Various Countries Visited by the H. M. S. Beagle*. London, Henry Colburn.
- Darwin, C. R. (1859). *On the Origin of Species by Means of Natural Selection*. London, John Murray.
- Darwin, C. R. (1875). *The Variation of Animals and Plants under Domestication, Second Edition, Volumes 1 and 2*. London, John Murray.
- Darwin, E. (1794–1796). *Zoonomia; or the Laws of Organic Life, Volumes I and II*. London, J. Johnson.
- Dawkins, R. (1976). *The Selfish Gene*. Oxford, Oxford University Press.
- de Beer, G. R. (1960). Darwin's notebooks on transmutation of species. Part I. First notebook (July 1837–February 1838). *Bulletin of the British Museum (Natural History), Historical Series*, 2, 25–73.
- Depéret, C. (1907). *Les transformations du monde animal*. Paris, Ernest Flammarion.
- de Vries, H. (1901). *Die Mutationstheorie. Versuch Und Beobachtungen Über Die Entstehung Der Arten Im Pflanzenreich. Erster Band. Die Entstehung Der Arten Durch Mutation*. Leipzig, Verlag von Veit.
- de Vries, H. (1905). The evidence of evolution. *Annual Report of the Smithsonian Institution*, 1904, 389–396.
- de Vries, H. (1909). *The Mutation Theory: Experiments and Observations on the Origin of Species in the Vegetable Kingdom*. Chicago, Open Court Publishing.
- Diamond, J. M., Pimm, S. L., Gilpin, M. E., and LeCroy, M. (1989). Rapid evolution of character displacement in myzomelid honeyeaters. *American Naturalist*, 134, 675–708.

- Dobzhansky, T. (1937). *Genetics and the Origin of Species*. New York, Columbia University Press.
- Dobzhansky, T. and Spassky, B. (1967). Effects of selection and migration on geotactic and phototactic behaviour of *Drosophila*. I. *Proceedings of the Royal Society of London, Series B*, 168, 27–47.
- Dudley, J. W. and Lambert, R. J. (2004). 100 generations of selection for oil and protein in corn. *Plant Breeding Reviews*, 24, 79–110.
- Eastman, L. M., Morelli, T. L., Rowe, K. C., Conroy, C. J. and Moritz, C. (2012). Size increase in high elevation ground squirrels over the last century. *Global Change Biology*, 18, 1499–1508.
- Edwards, A. W. F. (1972). *Likelihood*. Cambridge, Cambridge University Press.
- Edwards, A. W. F. (1992). *Likelihood, Expanded Edition*. Baltimore, Maryland, Johns Hopkins University Press.
- Efron, B. and Tibshirani, R. J. (1986). Bootstrap methods for standard errors, confidence intervals, and other measures of statistical accuracy. *Statistical Science*, 1, 54–77.
- Efron, B. and Tibshirani, R. J. (1993). *An Introduction to the Bootstrap*. New York, Chapman and Hall 436.
- Eisenberg, J. F. (1981). *The Mammalian Radiations: An Analysis of Trends in Evolution, Adaptation, and Behavior*. Chicago, University of Chicago Press.
- Eldredge, N. (1971). The allopatric model and phylogeny in Paleozoic invertebrates. *Evolution*, 25, 156–167.
- Eldredge, N. (1972). Systematics and evolution of *Phacops rana* (Green, 1832) and *Phacops iowensis* Delo, 1935 (Trilobita) from the middle Devonian of North America. *Bulletin of the American Museum of Natural History*, 147, 45–114.
- Eldredge, N. and Gould, S. J. (1972). Punctuated equilibria: an alternative to phyletic gradualism. In *Models in Paleobiology*, ed. T. J. M. Schopf, San Francisco, Freeman, Cooper and Company, pp. 82–115.
- Elliott, E. B. (1863). *On the Military Statistics of the United States of America*. Berlin, International Statistical Congress: R. v. Decker.
- Elliott, E. B. (1865). On the military statistics of the United States of America. In *Fünfte Internationalen Statistischen Congresses*, ed. E. Engel, Berlin, Königlichen Geheimen Ober-Hofbuchdruckerei, pp. 715–758.
- Eloy de Amorim, M., Schoener, T. W., Santoro, G. R. C. C., Lins, A. C. R., Piovia-Scott, J., and Brand+úo, R. A. (2017). Lizards on newly created islands independently and rapidly adapt in morphology and diet. *Proceedings of the National Academy of Sciences USA*, 114, 8812–8816.
- Endler, J. A. (1986). *Natural Selection in the Wild*. Princeton, Princeton University Press 336.
- Erickson, G. M., Currie, P. J., Inouye, B. D., and Winn, A. A. (2006). Tyrannosaur life tables: an example of nonavian dinosaur population biology. *Science*, 313, 213–217.
- Erickson, G. M., Makovicky P. J., Inouye B. D., Zhou C.-F., and Gao, K. -Q. (2009). A life table for *Psittacosaurus lujiatunensis*: initial insights into ornithischian dinosaur population biology. *Anatomical Record*, 292, 1514–1521.
- Erickson, W. A. and Halvorson, W. L. (1990). Ecology and control of the roof rat (*Rattus rattus*) in Channel Islands National Park. *Cooperative National Park Resources Studies Unit, Institute of Ecology, University of California, Davis*, 38, 1–90.
- Ernest, S. K. M. (2003). Life history characteristics of placental nonvolant mammals. *Ecology*, 84, 3402.
- Estes, S. and Arnold, S. J. (2007). Resolving the paradox of stasis: models with stabilizing selection explain evolutionary divergence on all timescales. *American Naturalist*, 169, 227–244.

- Falconer, D. S. (1973). Replicated selection for body weight in mice. *Genetical Research, Cambridge*, 22, 291–321.
- Falconer, D. S. (1977). Why are mice the size they are? In *International Conference on Quantitative Genetics*, eds. E. Pollack, O. Kempthorne, and T. B. Bailey, Ames, Iowa State University Press, pp. 19–21.
- Falconer, D. S. (1981). *Introduction to Quantitative Genetics, Second Edition*. London, Longman Group Limited.
- Falconer, H. (1863). On the American fossil elephant of the regions bordering the Gulf of Mexico (*E. columbi* Falc.); with general observations on the living and extinct species. *Natural History Review, Dublin*, 3, 43–114.
- Fechner, G. T. (1860). *Elemente Der Psychophysik*. Leipzig, Breitkopf and Härtel.
- Feller, W. (1968). *An Introduction to Probability Theory and Its Applications*, Third Edition, Volume I. New York, John Wiley and Sons 509.
- Felsenstein, J. (1973). Maximum-likelihood estimation of evolutionary trees from continuous characters. *American Journal of Human Genetics*, 25, 471–492.
- Felsenstein, J. (1985). Phylogenies and the comparative method. *American Naturalist*, 125, 1–15.
- Festing, M. F. W. (1973). A multivariate analysis of subline divergence in the shape of the mandible in C57BL/Gr mice. *Genetical Research, Cambridge*, 21, 121–132.
- Fisher, R. A. and Ford, E. B. (1950). The “Sewall Wright effect.” *Heredity*, 4, 117–119.
- Foote, M. (1997). The evolution of morphological diversity. *Annual Review of Ecology and Systematics*, 28, 129–152.
- Forstén, A.-M. (1990). Dental size trends in an equid sample from the Sandalja II cave of northwestern Yugoslavia. *Paläontologische Zeitschrift*, 64, 153–160.
- Franks, S. J., Sim, S., and Weis, A. E. (2007). Rapid evolution of flowering time by an annual plant in response to a climate fluctuation. *Proceedings of the National Academy of Sciences USA*, 104, 1278.
- Freudenthal, M. and Martín-Suárez, E. (2013). Estimating body mass of fossil rodents. *Scripta Geologica*, 145, 1–130.
- Froese, D., Stiller, M., Heintzman, P. D., Reyes, A. V., Zazula, G. D., Soares, A. E. R., Meyer, M., Hall, E., Jensen, B. J. L., Arnold, L. J., MacPhee, R. D. E., and Shapiro, B. (2017). Fossil and genomic evidence constrains the timing of bison arrival in North America. *Proceedings of the National Academy of Sciences USA*, 114, 3457.
- Gaillard, J.-M., Pontier, D., Allaine, D., Lebreton, J.-D., Trouvilliez, J. and Clobert, J. (1989). An analysis of demographic tactics in birds and mammals. *Oikos*, 56, 59–76.
- Galton, F. (1869). *Hereditary Genius: An Inquiry into Its Laws and Consequences*. London, Macmillan.
- Galton, F. (1879). The geometric mean in vital and social statistics. *Proceedings of the Royal Society of London*, 29, 365–367.
- Galton, F. (1889). *Natural Inheritance*. London, MacMillan.
- Galton, F. (1894). Discontinuity in evolution. *Mind, a Quarterly Review of Psychology and Philosophy, London*, 3, 362–372.
- Garland, T. (1992). Rate tests for phenotypic evolution using phylogenetically independent contrasts. *American Naturalist*, 140, 509–519.
- Garland, T. and Janis, C. M. (1993). Does metatarsal/femur ratio predict maximal running speed in cursorial mammals? *Journal of Zoology, London*, 229, 133–151.
- Gates, R. R. (1911). The mutation theory [book review]. *American Naturalist*, 45, 254–256.
- Gaudry, A. (1896). *Essai De Paléontologie Philosophique*. Paris, Masson.
- Gauss, C. F. (1809). *Theoria Motus Corporum Celestium*. Hamburg, Perthes et Besser.
- Gavrilets, S. and Vose, A. (2005). Dynamic patterns of adaptive radiation. *Proceedings of the National Academy of Sciences USA*, 102, 18040–18045.

- Geary, R. C. (1935). The ratio of the mean deviation to the standard deviation as a test of normality. *Biometrika*, 27, 310–332.
- Geiger, M., Sánchez-Villagra, M., and Lindholm, A. K. (2018). A longitudinal study of phenotypic changes in early domestication of house mice. *Royal Society Open Science*, 5, 172099.
- Geoffroy Saint-Hilaire, É. (1828). Mémoire où l'on se propose de rechercher dans quels rapports de structure organique et de parenté sont entre eux les animalux des âges historiques et vivant actuellement et les espèces antédiluviennes et perdues. *Mémoires du Muséum National d'Histoire Naturelle, Paris*, 17, 209–229.
- Geoffroy Saint-Hilaire, I. (1836). *Histoire générale et particulière des anomalies de l'organisation chez l'homme et les animaux, ou traité de tératologie*, tome troisième. Paris, J.-B. Baillière.
- George, T. N. (1958). Rates of change in evolution. *Science Progress, a Quarterly Review of Current Scientific Investigations, London*, 46, 409–428.
- Gilchrist, G. W., Huey, R. B., Balanyà, J., Pascual, M., and Serra, L. (2004). A time series of evolution in action: a latitudinal cline in wing size in South American *Drosophila subobscura*. *Evolution*, 58, 768–780.
- Gilchrist, G. W., Huey, R. B., and Serra, L. (2001). Rapid evolution of wing size clines in *Drosophila subobscura*. *Genetica*, 112, 273–286.
- Gillespie, J. H. (1984). The molecular clock may be an episodic clock. *Proceedings of the National Academy of Sciences USA*, 81, 8009–8013.
- Gingerich, P. D. (1974). Stratigraphic record of early Eocene *Hyopsodus* and the geometry of mammalian phylogeny. *Nature*, 248, 107–109.
- Gingerich, P. D. (1976). Paleontology and phylogeny: patterns of evolution at the species level in early Tertiary mammals. *American Journal of Science*, 276, 1–28.
- Gingerich, P. D. (1983). Rates of evolution: effects of time and temporal scaling. *Science*, 222, 159–161.
- Gingerich, P. D. (1984). Punctuated equilibria – where is the evidence? *Systematic Zoology*, 33, 335–338.
- Gingerich, P. D. (1989). New earliest Wasatchian mammalian fauna from the Eocene of northwestern Wyoming: composition and diversity in a rarely sampled high-floodplain assemblage. *University of Michigan Papers on Paleontology*, 28, 1–97.
- Gingerich, P. D. (1993). Quantification and comparison of evolutionary rates. *American Journal of Science*, 293, 453–478.
- Gingerich, P. D. (1994). New species of *Apheliscus*, *Haplomylus*, and *Hyopsodus* (Mammalia, Condylarthra) from the late Paleocene of southern Montana and early Eocene of northwestern Wyoming. *Contributions from the Museum of Paleontology, University of Michigan*, 29, 119–134.
- Gingerich, P. D. (1995). Statistical power of EDF tests of normality and the sample size required to distinguish geometric-normal (lognormal) from arithmetic-normal distributions of low variability. *Journal of Theoretical Biology*, 173, 125–136.
- Gingerich, P. D. (2000). Arithmetic or geometric normality of biological variation: an empirical test of theory. *Journal of Theoretical Biology*, 204, 201–221.
- Gingerich, P. D. (2001). Rates of evolution on the time scale of the evolutionary process. *Genetica*, 112/113, 127–144.
- Gingerich, P. D. (2003). Mammalian responses to climate change at the Paleocene-Eocene boundary: Polecat Bench record in the northern Bighorn Basin, Wyoming. In *Causes and consequences of globally warm climates in the early Paleogene*, eds. S. L. Wing, P. D. Gingerich, B. Schmitz, and E. Thomas, Geological Society of America, Special Papers, 369: 463–478.

- Gingerich, P. D. (2010). Mammalian faunal succession through the Paleocene-Eocene thermal maximum (PETM) in western North America. *Vertebrata Palasiatica, Beijing*, 48, 308–327.
- Gingerich, P. D. (2014). Species in the primate fossil record. *Evolutionary Anthropology*, 23, 33–35.
- Gingerich, P. D. (2015). Body weight and relative brain size (encephalization) in Eocene Archaeoceti (Cetacea). *Journal of Mammalian Evolution*, 23, 17–31.
- Gingerich, P. D. (2019). Data from: Rates of Evolution: A Quantitative Synthesis (Cambridge University Press). *Dryad Digital Repository*, <https://doi.org/10.5061/dryad.1tn7123>.
- Gingerich, P. D., Smith, B. H. and Rosenberg, K. R. (1982). Allometric scaling in the dentition of primates and prediction of body weight from tooth size in fossils. *American Journal of Physical Anthropology*, 58, 81–100.
- Gingerich, P. D. and Smith, T. (2006). Paleocene-Eocene land mammals from three new latest Clarkforkian and earliest Wasatchian wash sites at Polecat Bench in the northern Bighorn Basin, Wyoming. *Contributions from the Museum of Paleontology, University of Michigan*, 31, 245–303.
- Goldschmidt, R. (1933). Some aspects of evolution. *Science*, 78, 539–547.
- Goldschmidt, R. (1940). *The Material Basis of Evolution*. New Haven, Yale University Press 436.
- Goodwin, N. B., Grant, A., Perry, A. L., Dulvy, N. K., and Reynolds, J. D. (2006). Life history correlates of density-dependent recruitment in marine fishes. *Canadian Journal of Fisheries and Aquatic Sciences*, 63, 494–509.
- Gould, S. J. (1969). An evolutionary microcosm: Pleistocene and Recent history of the land snail *P. (Poecilozonites)* in Bermuda. *Bulletin of the Museum of Comparative Zoology, Harvard University*, 138, 407–532.
- Gould, S. J. (1984). Smooth curve of evolutionary rate: a psychological and mathematical artifact. *Science*, 226, 994–995.
- Gould, S. J. (1985). The paradox of the first tier: an agenda for paleobiology. *Paleobiology*, 11, 2–12.
- Gould, S. J. (1988). Trends as changes in variance: a new slant on progress and directionality in evolution. *Journal of Paleontology*, 62, 319–329.
- Gould, S. J. and Eldredge, N. (1977). Punctuated equilibria: the tempo and mode of evolution reconsidered. *Paleobiology*, 3, 115–151.
- Gowe, R. S. and Fairfull, R. W. (1985). The direct response to long-term selection for multiple traits in egg stocks and changes in genetic parameters with selection. In *Poultry Genetics and Breeding*, eds. W. G. Hill, J. M. Manson, and D. Hewitt, Harlow, British Poultry Science Ltd., pp. 125–146.
- Gowe, R. S., Johnson, A. S., Downs, J. H., Gibson, R., Mountain, W. F., Strain, J. H., and Tinney, B. F. (1959). Environment and poultry breeding problems: 4. The value of a random-bred control strain in a selection; study. *Poultry Science*, 38, 443–462.
- Gradstein, F. M., Ogg, J. G., Schmitz, M. D., and Ogg, G. M. eds. (2012). *The Geological Time Scale 2012*, Amsterdam, Elsevier.
- Grant, B. R. and Grant, P. R. (2010). Songs of Darwin's finches diverge when a new species enters the community. *Proceedings of the National Academy of Sciences*, 107, 20156–20163.
- Grant, P. R. (1981). Speciation and the adaptive radiation of Darwin's finches. *American Scientist*, 69, 653–663.
- Grant, P. R. (1986). *Ecology and Evolution of Darwin's Finches*. Princeton, Princeton University Press.

- Grant, P. R. (1994). Population variation and hybridization: comparison of finches from two archipelagos. *Evolutionary Ecology*, 8, 598–617.
- Grant, P. R. and Grant, B. R. (1993). Evolution of Darwin's finches caused by a rare climatic event. *Proceedings of the Royal Society B: Biological Sciences*, 251, 111–117.
- Grant, P. R. and Grant, B. R. (2002). Unpredictable evolution in a 30-year study of Darwin's finches. *Science*, 296, 707–711.
- Grant, P. R. and Grant, B. R. (2014). *40 Years of Evolution: Darwin's Finches on Daphne Major Island*. Princeton, Princeton University Press.
- Grant, V. (1963). *The Origin of Adaptations*. New York, Columbia University Press.
- Grave, B. H. (1930). The natural history of *Bugula flabellata* at Woods Hole, Massachusetts, including the behavior and attachment of the larva. *Journal of Morphology*, 49, 355–383.
- Haeckel, E. H. P. A. (1866). *Generelle Morphologie Der Organismen. Zweiter Band. Allgemeine Entwicklungsgeschichte Der Organismen*. Berlin, Georg Reimer.
- Hairston, N. G. (1987). *Community Ecology and Salamander Guilds*. Cambridge, Cambridge University Press.
- Haldane, J. B. S. (1949). Suggestions as to quantitative measurement of rates of evolution. *Evolution*, 3, 51–56.
- Haldane, J. B. S. (1958). The theory of evolution, before and after Bateson. *Journal of Genetics*, 56, 11–27.
- Haller, B. C. and Hendry, A. P. (2014). Solving the paradox of stasis: squashed stabilizing selection and the limits of detection. *Evolution*, 68, 483–500.
- Hansen, T. F. (1997). Stabilizing selection and the comparative analysis of adaptation. *Evolution*, 51, 1341–1351.
- Hansen, T. F. and Martins, E. P. (1996). Translating between microevolutionary process and macroevolutionary patterns: the correlation structure of interspecific data. *Evolution*, 50, 1404–1417.
- Hargenvilliers, A.-A. (1817). *Recherches et considérations sur la formation et le recrutement de l'armée en France*. Paris, F. Didot.
- Harmon, L. J., Losos, J. B., Jonathan Davies, T., Gillespie, R. G., Gittleman, J. L., Bryan Jennings, W., Kozak, K. H., McPeck, M. A., Moreno-Roark, F., Near, T. J., Purvis, A., Ricklefs, R. E., Schluter, D., Schulte II, J. A., Seehausen, O., Sidlauskas, B. L., Torres-Carvajal, O., Weir, J. T., and Mooers, A. Ø. (2010). Early bursts of size and shape evolution are rare in comparative data. *Evolution*, 64, 2385–2396.
- Haugen, T. O. and Vøllestad, L. A. (2000). Population differences in early life-history traits in grayling. *Journal of Evolutionary Biology*, 13, 897–905.
- Haugen, T. O. and Vøllestad, L. A. (2001). A century of life-history evolution in grayling. *Genetica*, 112, 475–491.
- Hayami, I. (1984). Natural history and evolution of *Cryptopecten* (a Cenozoic-Recent pectinid genus). *Bulletin of the University Museum, University of Tokyo*, 24, 1–149.
- Hendry, A. P. and Kinnison, M. T. (1999). The pace of modern life: measuring rates of contemporary microevolution. *Evolution*, 53, 1637–1653.
- Hendry, A. P. and Quinn, T. P. (1997). Variation in adult life history and morphology among Lake Washington sockeye salmon (*Oncorhynchus nerka*) populations in relation to habitat features and ancestral affinities. *Canadian Journal of Fisheries and Aquatic Sciences*, 54, 75–84.
- Heppell, S. S., Caswell, H., and Crowder, L. B. (2000). Life histories and elasticity patterns: perturbation analysis for species with minimal demographic data. *Ecology*, 81, 654–665.
- Herschel, J. (1850). Quetelet on probabilities. *Edinburgh Review*, 92, 1–57.
- Hilgen, F. J., Lourens, L. J., Van Dam, J. A., Beu, A. G., Boyes, A. F., Cooper, R. A., Krijgsman, W., Ogg, J. G., Piller, W. E., and Wilson, D. S. (2012). The Neogene

- period. In *The Geological Time Scale 2012*, eds. F. M. Gradstein, J. G. Ogg, M. D. Schmitz, and G. M. Ogg, Amsterdam, Elsevier, pp. 923–978.
- Hoekstra, H. E., Hoekstra, J. M., Vignieri, S. N., Hoang, A., Beerli, P., and Kingsolver, J. G. (2002). Strength and tempo of directional selection in the wild. *Proceedings of the National Academy of Sciences USA*, 98, 9157–9160.
- Holmes, M. W., Boykins, G. K. R., Bowie, R. C. K., and Lacey, E. A. (2016). Cranial morphological variation in *Peromyscus maniculatus* over nearly a century of environmental change in three areas of California. *Journal of Morphology*, 277, 96–106.
- Hooks, A. P. (2013). *Prey plasticity responses to a native and nonnative predator*. M.Sc. thesis, Stony Brook, Stony Brook University, 45 pp.
- Hopkins, C. G. (1899). Improvement in the chemical composition of the corn kernel. *Illinois Agriculture Experimental Station Bulletin*, 55, 205–240.
- House, M. R. (1963). Bursts in evolution. *Report of the British Association for the Advancement of Science, London*, 19, 499–507.
- Huber, P. J. (1981). *Robust Statistics*. New York, John Wiley and Sons.
- Hudson, J. D. and Martill, D. M. (1994). The Peterborough Member (Callovian, Middle Jurassic) of the Oxford Clay Formation at Peterborough, UK. *Journal of the Geological Society*, 151, 113.
- Hughes, M., Gerber, S., and Wills, M. A. (2013). Clades reach highest morphological disparity early in their evolution. *Proceedings of the National Academy of Sciences USA*, 110, 13875–13879.
- Hunt, G. (2004). Phenotypic variation in fossil samples: modeling the consequences of time-averaging. *Paleobiology*, 30, 426–443.
- Hunt, G. (2006). Fitting and comparing models of phyletic evolution: random walks and beyond. *Paleobiology*, 32, 578–601.
- Hunt, G. (2007). Evolutionary divergence in directions of high phenotypic variance in the ostracode genus *Poseidonamicus*. *Evolution*, 61, 1560–1576.
- Hunt, G. and Roy, K. (2006). Climate change, body size evolution, and Cope's Rule in deep-sea ostracodes. *Proceedings of the National Academy of Sciences USA*, 103, 1347–1352.
- Hurst, H. E. (1951). Long-term storage capacity of reservoirs. *Transactions of the American Society of Civil Engineers*, 116, 770–808.
- Hutchinson, G. E. (1959). Homage to Santa Rosalia or why are there so many kinds of animals? *American Naturalist*, 93, 145–159.
- Huxley, J. S. (1924). Constant differential growth-ratios and their significance. *Nature*, 114, 895.
- Huxley, J. S. (1932). *Problems of Relative Growth*. London, Methuen.
- Huxley, J. S. (1942). *Evolution, the Modern Synthesis*. New York, Harper and Brothers.
- Huxley, J. S. and Teissier, G. (1936). Terminology of relative growth. *Nature*, 137, 780–781.
- Huxley, L. (1900). *Life and Letters of Thomas Henry Huxley, Volume 1*. London, Macmillan.
- Huxley, T. H. (1859). Letter to Charles Lyell dated 25 June 1859. In *Life and Letters of Thomas Henry Huxley, Volume 1 (1900)*, ed. L. Huxley, London, Macmillan, pp. 185–187.
- Ingram, T., Harmon, L. J., and Shurin, J. B. (2012). When should we expect early bursts of trait evolution in comparative data? Predictions from an evolutionary food web model. *Journal of Evolutionary Biology*, 25, 1902–1910.
- Jablonski, D. (1997). Body-size evolution in Cretaceous molluscs and the status of Cope's rule. *Nature*, 385, 250–252.
- Jaekel, O. (1902). *Über Verschiedene Wege Phylogenetischer Entwicklung*. Jena, Gustav Fischer.

- Jancović, I., Ahern, J. C. M., Karavanić, I., Stockton, T., and Smith, F. H. (2012). Epigravettian human remains and artifacts from Sandalja II, Istria, Croatia. *PaleoAnthropology*, 2012, 87–122.
- Janis, C. M. (1990). Correlation of cranial and dental variables with body size in ungulates and macropodids. In *Body Size in Mammalian Paleobiology: Estimation and Biological Implications*, eds. J. D. Damuth and B. J. MacFadden, Cambridge, Cambridge University Press, pp. 255–299.
- Jensen, H., Steinsland, I., Ringsby, T. H., and Saether, B.-E. (2008). Evolutionary dynamics of a sexual ornament in the house sparrow (*Passer domesticus*): the role of indirect selection within and between sexes. *Evolution*, 62, 1275–1293.
- Jepsen, G. L., Mayr, E., and Simpson, G. G. eds. (1949). *Genetics, Paleontology, and Evolution*, Princeton, Princeton University Press.
- Johannsen, W. L. (1909). *Elemente Der Exakten Erblichkeitslehre*. Jena, Gustav Fischer.
- Johnston, R. F. and Selander, R. K. (1964). House sparrows: rapid evolution of races in North America. *Science*, 144, 548–550.
- Johnston, R. F. and Selander, R. K. (1971). Evolution in the house sparrow. II. Adaptive differentiation in North American populations. *Evolution*, 25, 1–28.
- Kaufmann, R. (1933). Variationsstatistische Untersuchungen über die “Artabwandlung” und “Artumbildung” an der oberkambrischen Trilobitengattung *Olenus* Dalm. *Abhandlungen aus dem Geologisch-Palaeontologisches Institut der Universität Greifswald*, 10, 1–55.
- Kellogg, D. E. (1975). The role of phyletic change in the evolution of *Pseudocubus vema* (Radiolaria). *Paleobiology*, 1, 359–370.
- Kennett, J. P. (1966). The *Globorotalia crassiformis* bioseries in north Westland and Marlborough, New Zealand. *Micropaleontology*, 12, 235–245.
- Kerk, M. v. d., de Kroon, H., Conde, D. A., and Jongejans, E. (2013). Carnivora population dynamics are as slow and as fast as those of other mammals: implications for their conservation. *PLoS One*, 8, e70354.
- Kimura, M. (1964). Diffusion models in population genetics. *Journal of Applied Probability*, 1, 177–232.
- Kimura, Y., Flynn, L. J., and Jacobs, L. L. (2015). A palaeontological case study for species delimitation in diverging fossil lineages. *Historical Biology*, 28, 189–198.
- Klepaker, T. (1993). Morphological changes in a marine population of threespined stickleback, *Gasterosteus aculeatus*, recently isolated in fresh water. *Canadian Journal of Zoology*, 71, 1251–1258.
- Koontz, T. L., Shepherd, U. L., and Marshall, D. (2001). The effect of climate change on Merriam’s kangaroo rat, *Dipodomys merriami*. *Journal of Arid Environments*, 49, 581–591.
- Kristjánsson, B. K. (2005). Rapid morphological changes in threespine stickleback, *Gasterosteus aculeatus*, in freshwater. *Environmental Biology of Fishes*, 74, 357–363.
- Kristjánsson, B. K., Skúlason, S., and Noakes, D. L. G. (2002). Rapid divergence in a recently isolated population of threespine stickleback (*Gasterosteus aculeatus* L.). *Evolutionary Ecology Research*, 4, 659–672.
- Kruuk, L. E. B., Merilä, J., and Sheldon, B. C. (2001). Phenotypic selection on a heritable size trait revisited. *American Naturalist*, 158, 557–571.
- Kruuk, L. E. B., Slate, J., Pemberton, J. M., Brotherstone, S., Guinness, F., and Clutton-Brock, T. H. (2002). Antler size in red deer: heritability and selection but no evolution. *Evolution*, 56, 1683–1695.
- Kubitschek, H. E. (1974). Operation of selection pressure on microbial populations. *Symposia of the Society for General Microbiology*, 24, 105–130.

- Kurtén, B. (1955). Sex dimorphism and size trends in the cave bear, *Ursus spelaeus* Rosenmüller and Heinroth. *Acta Zoologica Fennica*, 90, 1–48.
- Kurtén, B. (1959). Rates of evolution in fossil mammals. *Cold Spring Harbor Symposia on Quantitative Biology*, 24, 205–215.
- Kurtén, B. (1963). The rate of evolution. In *Science in Archaeology: A Comprehensive Survey of Progress and Research*, eds. D. R. Brothwell and E. Higgs, Bristol, Thames and Hudson, pp. 217–223.
- Kurtén, B. (1965). Evolution in geological time. In *Ideas in Modern Biology*, ed. J. A. Moore, Garden City, New York, Natural History Press, pp. 329–354.
- Lack, D. (1940). Variation in the introduced English sparrow. *Condor*, 42, 239–241.
- Lack, D. (1947). *Darwin's Finches, an Essay on the General Biological Theory of Evolution*. Cambridge, Cambridge University Press.
- Lamarck, J.-B. (1809). *Philosophie Zoologique, Tome Premier, Tome Second*. Paris, Librairie Dentu.
- Lande, R. (1976). Natural selection and random genetic drift in phenotypic evolution. *Evolution*, 30, 314–334.
- Lande, R. (1977). On comparing coefficients of variation. *Systematic Zoology*, 26, 214–217.
- Langergraber, K. E., Prüfer, K., Rowney, C., Boesch, C., Crockford, C., Fawcett, K., Inoue, E., Inoue-Muruyama, M., Mitani, J. C., Muller, M. N., Robbins, M. M., Schubert, G., Stoinski, T. S., Viola, B., Watts, D., Wittig, R. M., Wrangham, R. W., Zuberbühler, K., Pääbo, S., and Vigilant, L. (2012). Generation times in wild chimpanzees and gorillas suggest earlier divergence times in great ape and human evolution. *Proceedings of the National Academy of Sciences*, 109, 15716–15721.
- Larsson, K., Jeugd, H. P. v. d., and Veen, I. T. v. d. F. P. (1998). Body size declines despite positive directional selection on heritable size traits in a barnacle goose population. *Evolution*, 52, 1169–1184.
- Lawler, R. R. (2011). Historical demography of a wild lemur population (*Propithecus verreauxi*) in southwest Madagascar. *Population Ecology*, 53, 229–240.
- Lee, A. H. and Werning, S. (2008). Sexual maturity in growing dinosaurs does not fit reptilian growth models. *Proceedings of the National Academy of Sciences USA*, 105, 582–587.
- Legendre, S. (1989). Les communautés de mammifères du Paléogène (Eocène supérieur et Oligocène) d'Europe occidentale: structures, milieux et évolution. *Münchener Geowissenschaftliche Abhandlungen, Reihe A, Geologie und Paläontologie*, 16, 1–110.
- Lerner, I. M. (1954). *Genetic Homeostasis*. Edinburgh, Oliver and Boyd.
- Lerner, I. M. and Dempster, E. R. (1951). Attenuation of genetic progress under continued selection in poultry. *Heredity*, 5, 75–94.
- Levin, S. A. (1992). The problem of pattern and scale in ecology. *Ecology*, 73, 1943–1967.
- Levinton, J. S. (1979). A theory of diversity equilibrium and morphological evolution. *Science*, 204, 335–336.
- Lewontin, R. C. (1966). On the measurement of relative variability. *Systematic Zoology*, 15, 141–142.
- Lexis, W. (1877). *Zur Theorie Der Massenerscheinungen in Der Menschlichen Gesellschaft*. Freiburg im Briesgau, Friedrich Wagner.
- Lich, D. K. (1990). *Cosomys primus*: a case for stasis. *Paleobiology*, 16, 384–395.
- Linnaeus, C. (1735). *Systema Naturae, Sive Regna Tria Naturae Systematice Proposita Per Classes, Ordines, Genera, Species*. Leiden, Lugduni Batavorum.
- Lister, A. M. (1989). Rapid dwarfing of red deer on Jersey in the last interglacial. *Nature*, 342, 539–542.

- Lovtrup (Løvtrup), S. (1974). *Epigenetics, a Treatise on Theoretical Biology*. New York, John Wiley and Sons.
- Lyell, C. (1832). *Principles of Geology, Volume II*. London, John Murray.
- Lynch, M. (1988). The rate of polygenic mutation. *Genetical Research, Cambridge*, 51, 137–148.
- Lynch, M. (1990). The rate of morphological evolution in mammals from the standpoint of the neutral expectation. *American Naturalist*, 136, 727–741.
- Mac Gillavry, H. J. (1968). Modes of evolution mainly among marine invertebrates. *Bijdragen tot de Dierkunde*, 38, 69–74.
- MacFadden, B. J. (1985). Patterns of phylogeny and rates of evolution in fossil horses: hipparions from the Miocene and Pliocene of North America. *Paleobiology*, 11, 245–257.
- MacFadden, B. J. (1986). Fossil horses from “Eohippus” (*Hyracotherium*) to *Equus*: scaling, Cope’s law, and the evolution of body size. *Paleobiology*, 12, 355–369.
- MacFadden, B. J. (1988). Fossil horses from “Eohippus” (*Hyracotherium*) to *Equus*, 2: rates of dental evolution revisited. *Biological Journal of the Linnean Society, London*, 35, 37–48.
- MacFadden, B. J. (1992). *Fossil Horses: Systematics, Paleobiology, and Evolution of the Family Equidae*. Cambridge, Cambridge University Press.
- MacFadden, B. J. and Hulbert, R. C. (1990). Body size estimates and size distribution of ungulate mammals from the Late Miocene Love Bone Bed of Florida. In *Body Size in Mammalian Paleobiology: Estimation and Biological Implications*, eds. J. D. Damuth and B. J. MacFadden, Cambridge, Cambridge University Press, pp. 337–363.
- MacLeod, N. (1991). Punctuated anagenesis and the importance of stratigraphy to paleobiology. *Paleobiology*, 17, 167–188.
- Maglio, V. J. (1973). Origin and evolution of the Elephantidae. *Transactions of the American Philosophical Society*, 63, 1–149.
- Malmgren, B. A., Berggren, W. A. and Lohmann, G. P. (1983). Evidence for punctuated gradualism in the late Neogene *Globorotalia tumida* lineage of planktonic foraminifera. *Paleobiology*, 9, 377–389.
- Malmgren, B. A. and Kennett, J. P. (1981). Phyletic gradualism in a late Cenozoic planktonic foraminiferal lineage; DSDP site 284, southwest Pacific. *Paleobiology*, 7, 230–240.
- Mandelbrot, B. B. (1967). How long is the coast of Britain? Statistical self-similarity and fractional dimension. *Science*, 156, 636–638.
- Mandelbrot, B. B. (1983). *Fractal Geometry of Nature*. San Francisco, W. H. Freeman.
- Manser, A., Lindholm, A. K., König, B., and Bagheri, H. C. (2011). Polyandry and the decrease of a selfish genetic element in a wild house mouse population. *Evolution*, 65, 2435–2447.
- Mayr, E. (1942). *Systematics and the Origin of Species*. New York, Columbia University Press.
- Mayr, E. (1954). Change of genetic environment and evolution. In *Evolution As a Process*, ed. J. S. Huxley, A. C. Hardy and E. B. Ford, London, George Allen and Unwin, pp. 157–180.
- Mayr, E. (1963). *Animal Species and Evolution*. Cambridge, Massachusetts, Harvard University Press.
- McAlister, D. (1879). The law of the geometric mean. *Proceedings of the Royal Society of London*, 29, 367–376.
- McCluskey, J., Olivier, T. J., Freedman, L., and Hunt, E. (1974). Evolutionary divergences between populations of Australian wild rabbits. *Nature*, 249, 278–279.

- McDonald, J. N. (1981). *North American Bison: Their Classification and Evolution*. Berkeley, University of California Press.
- McKellar, A. E. and Hendry, A. P. (2009). How humans differ from other animals in their levels of morphological variation. *PLoS One*, 4, e6876.
- McPeck, M.-A. (2008). The ecological dynamics of clade diversification and community assembly. *American Naturalist*, 172, E270–E284.
- McPhail, J. D. (1984). Ecology and evolution of sympatric sticklebacks (*Gasterosteus*): morphological and genetic evidence for a species pair in Enos Lake, British Columbia. *Canadian Journal of Zoology*, 62, 1402–1408.
- Meehan, T. (1875). *Change by Gradual Modification Not the Universal Law*. Salem, Massachusetts, Salem Press.
- Mendel, G. (1866). Versuche über Pflanzen-Hybriden. *Verhandlungen des naturforschenden Vereines in Brünn, Abhandlungen*, 4, 3–47.
- Michaux, B. (1988). Organotaxism: an alternative “way of seeing” the fossil record. *Journal of Theoretical Biology*, 133, 397–408.
- Millar, J. S. and Zammuto, R. M. (1983). Life histories of mammals: an analysis of life tables. *Ecology*, 64, 631–635.
- Miller, G. S. (1912). *Catalogue of the Mammals of Western Europe*. London, British Museum (Natural History).
- Millien, V., Ledevin, R., Boué, C., and Gonzalez, A. (2017). Rapid morphological divergence in two closely related and co-occurring species over the last 50 years. *Evolutionary Ecology*, 31, 847–864.
- Milner, J. M., Albon, S. D., Illius, A. W., Pemberton, J. M., and Clutton-Brock, T. H. (1999). Repeated selection of morphometric traits in the Soay sheep on St Kilda. *Journal of Animal Ecology*, 68, 472–488.
- Miracle, P. T. (1995). *Broad-Spectrum Adaptations Re-Examined: Hunter-Gatherer Responses to Late Glacial Environmental Changes in the Eastern Adriatic*. Ph.D., Ann Arbor, University of Michigan, 577 pp.
- Mivart, St. G. J. (1871). *On the Genesis of Species*. London, Macmillan 342.
- Møller, A. P. and Szép, T. (2005). Rapid evolutionary change in a secondary sexual character linked to climatic change. *Journal of Evolutionary Biology*, 18, 481–495.
- Moorjani, P., Amorim, C. E. G., Arndt, P. F., and Przeworski, M. (2016). Variation in the molecular clock of primates. *Proceedings of the National Academy of Sciences USA*, 113, 10607–10612.
- Moss, C. J. (2001). The demography of an African elephant (*Loxodonta africana*) population in Amboseli, Kenya. *Journal of Zoology*, 255, 145–156.
- Müller, F. (1886). Ein Zuchtungsversuch an Mais. *Kosmos, Stuttgart*, 19, 22–26.
- Nengovhela, A., Baxter, R. M., and Taylor, P. J. (2015). Temporal changes in cranial size in South African vlei rats (*Otomys*): evidence for the “third universal response to warming,” *African Zoology*, 50, 233–239.
- Neumayr, M. and Paul, C. M. (1875). Die Congerien- und Paludinenschichten Slavoniens und deren Faunen. Ein Beitrag zur Descendenz-Theorie. *Abhandlungen der Kaiserlich-Königlichen Geologischen Reichsanstalt, Wien*, 7(3), 1–113.
- Newell, N. D. (1949). Phyletic size increase – an important trend illustrated by fossil invertebrates. *Evolution*, 3, 103–124.
- Niel, C. and Lebreton, J.-D. (2005). Using demographic invariants to detect overharvested bird populations from incomplete data. *Conservation Biology*, 19, 826–835.
- Ogg, J. G., Hinnov, L. A., and Huang, C. (2012). Jurassic. In *The Geological Time Scale 2012*, eds. F. M. Gradstein, J. G. Ogg, M. D. Schmitz, and G. M. Ogg, Amsterdam, Elsevier, pp. 731–791.

- Orbigny, A. D. d. (1851). *Cours Élémentaire De Paléontologie Et De Géologie Stratigraphiques, Tome Second*. Paris, Victor Masson.
- Osborn, H. F. (1905). Present problems in paleontology. *Popular Science Monthly*, 66, 226–242.
- Osborn, H. F. (1925). The origin of species as revealed by vertebrate palaeontology. *Nature*, 115, 925–926, 961–963.
- Ovcharenko, V. N. (1969). Transitional forms and species differentiation of brachiopods. *Paleontological Journal*, 3, 57–63.
- Owen, R. (1868). *Anatomy of Vertebrates. Volume III. Mammals*. London, Longmans, Green, and Co. 915.
- Palmer, A. R. (1965). Biome – a new kind of biostratigraphic unit. *Journal of Paleontology*, 39, 149–153.
- Paradis, E. (2012). *Analysis of Phylogenetics and Evolution With R*, Second Edition. New York, Springer.
- Patton, J. L., Yang, S. Y., and Myers, P. (1975). Genetic and morphologic divergence among introduced rat populations (*Rattus rattus*) of the Galapagos archipelago, Ecuador. *Systematic Zoology*, 24, 296–310.
- Payne, J. L., Boyer, A. G., Brown, J. H., Finnegan, S., Kowalewski, M., Krause, R. A., Lyons, S. K., McClain, C. R., McShea, D. W., Novack-Gottshall, P. M., Smith, F. A., Stempien, J. A., and Wang, S. C. (2009). Two-phase increase in the maximum size of life over 3.5 billion years reflects biological innovation and environmental opportunity. *Proceedings of the National Academy of Sciences*, 106, 24–27.
- Pearson, E. S. and Hartley, H. O. (1966). *Biometrika Tables for Statisticians, Volume I*, Third Edition. Cambridge, Cambridge University Press.
- Pearson, K. (1894). Contributions to the mathematical theory of evolution. I. On the dissection of asymmetrical frequency curves. *Philosophical Transactions of the Royal Society of London, Series A*, 185, 71–110.
- Pearson, K. (1895). Contributions to the mathematical theory of evolution. II. Skew variation in homogeneous material. *Philosophical Transactions of the Royal Society of London, Series A*, 186, 343–414.
- Pearson, K. (1905). The problem of the random walk. *Nature*, 72, 294, 342.
- Pearson, K. (1924). *The Life, Letters and Labours of Francis Galton. Volume II—Researches of Middle Life*. London, Cambridge University Press.
- Pegueroles, G., Papacit, M., Quintana, A., Guillén, A., Prevosti, A., and Serra, L. (1995). An experimental study of evolution in progress: clines for quantitative traits in colonizing and Palearctic populations of *Drosophila*. *Evolutionary Ecology*, 9, 453–465.
- Peirce, C. S. (1873). On the theory of errors of observations. *U. S. Coast Survey Report for 1870, Appendix 21*, 1870, 200–224.
- Pergams, O. R. W. and Ashley, M. V. (1999). Rapid morphological change in island deer mice. *Evolution*, 53, 1573–1581.
- Pergams, O. R. W., Byrn, D., Lee, K. L. Y., and Jackson, R. (2015). Rapid morphological change in black rats (*Rattus rattus*) after an island introduction. *Peer J*, 3, e812.
- Peters, R. H. (1983). *Ecological Implications of Body Size*. Cambridge, Cambridge University Press.
- Petren, K., Grant, B. R., and Grant, P. R. (1999). A phylogeny of Darwin's finches based on microsatellite DNA length variation. *Proceedings of the Royal Society of London, Series B*, 266, 321–329.
- Pfeifer, S. P. (2017). The demographic and adaptive history of the African green monkey. *Molecular Biology and Evolution*, 34, 1055–1065.
- Philipitschenko, J. (1927). *Variabilität Und Variation*. Berlin, Gebrüder Borntraeger.

- Phillimore, A. B. and Price, T. D. (2008). Density-dependent cladogenesis in birds. *PLoS Biology*, 6, e71.
- Playfair, J. (1802). *Illustrations of the Huttonian Theory of the Earth*. Edinburgh, William Creech.
- Porter, W. T. (1894). The growth of St. Louis children. *Transactions of the Academy of Science of St. Louis*, 6, 263–380.
- Press, W. H., Flannery, B. P., Teukolsky, S. A., and Vetterling, W. T. (1989). *Numerical Recipes: The Art of Scientific Computing (Fortran Version)*. Cambridge, Cambridge University Press.
- Przybylo, R., Sheldon, B. C., and Merilä, J. (2000). Climatic effects on breeding and morphology: evidence for phenotypic plasticity. *Journal of Animal Ecology*, 69, 395–403.
- Quetelet, A. (1835). *Sur l'homme et le développement de ses facultés, ou essai de physique sociale*, Tome Premier. Paris, Bachelier.
- Quetelet, A. (1846). *Lettres sur la théorie des probabilités, appliquée aux sciences morales et politiques*. Bruxelles, M. Hayez.
- Quetelet, A. (1849). *Letters Addressed to H.R.H. the Grand Duke of Saxe Coburg and Gotha, on the Theory of Probabilities As Applied to the Moral and Political Sciences Translated From the French by Olinthus Gregory Downes*. London, C. and E. Layton.
- Quetelet, A. (1869). *Physique sociale, ou essay sur le développement des facultés*, Tome I. Brussels, C. Muquardt.
- Quetelet, A. (1870). *Anthropométrie ou mesure des différentes facultés de l'homme*. Brussels, C. Muquardt.
- R Core Team. (2017). *R: A Language and Environment for Statistical Computing*. Vienna, Austria, R Foundation for Statistical Computing.
- Rabosky, D. L. and Lovette, I. J. (2008). Explosive evolutionary radiations: decreasing speciation or increasing extinction through time? *Evolution*, 62, 1866–1875.
- Raup, D. M. and Crick, R. E. (1981). Evolution of single characters in the Jurassic ammonite *Kosmoceras*. *Paleobiology*, 7, 200–215.
- Reiss, M. J. (1989). *The Allometry of Growth and Reproduction*. Cambridge, Cambridge University Press.
- Rensch, B. (1943). Die paläontologischen Evolutionsregeln in zoologischer Betrachtung. *Biologia Generalis*, Vienna, 17, 1–55.
- Rensch, B. (1947). *Neuere Probleme Der Abstammungslehre: Die Transspezifische Evolution*. Stuttgart, Ferdinand Enke Verlag.
- Reznick, D. N., Shaw, F. H., Rodd, F. H., and Shaw, R. G. (1997). Evaluation of the rate of evolution in natural populations of guppies (*Poecilia reticulata*). *Science*, 275, 1934–1937.
- Richard, A. F., Dewar, R. E., Schwartz, M., and Ratsirarson, J. (2000). Mass change, environmental variability and female fertility in wild *Propithecus verreauxi*. *Journal of Human Evolution*, 39, 381–391.
- Richard-Hansen, C., Vié, J.-C., Vidal, N., and Kéramec, J. (1999). Body measurements on 40 species of mammals from French Guiana. *Journal of Zoology*, 247, 419–428.
- Richardson, L. F. (1961). The problem of contiguity: an appendix to Statistics of Deadly Quarrels. *General Systems Yearbook, Society for the Advancement of General Systems Theory*, 6, 139–187.
- Rickwood, A. E. (1977). Age, growth and shape of the intertidal brachiopod *Waltonia inconspicua* Sowerby, from New Zealand. *American Zoologist*, 17, 63–73.
- Roden, M. K., Parrish, R. R., and Miller, D. S. (1990). The absolute age of the Eifelian Tioga Ash bed, Pennsylvania. *Journal of Geology*, 98, 282–285.

- Sander, P. M., Christian, A., Clauss, M., Fechner, R., Gee, C. T., Griebeler, E.-M., Gunga, H.-C., Hummel, J., Mallison, H., Perry, S. F., Preuschoft, H., Rauhut, O. W. M., Remes, K., Tütken, T., Wings, O., and Witzel, U. (2011). Biology of the sauropod dinosaurs: the evolution of gigantism. *Biological Reviews*, 86, 117–155.
- Sato, A., Tichy, H., O'hUigin, C., Grant, P. R., Grant, B. R., and Klein, J. (2001). On the origin of Darwin's finches. *Molecular Biology and Evolution*, 18, 299–311.
- Saunders, W. B. (1984). *Nautilus* growth and longevity: evidence from marked and recaptured animals. *Science*, 224, 992–990.
- Schindewolf, O. H. (1936). *Paläontologie, Entwicklungslehre Und Genetik: Kritik Und Synthese*. Berlin, Gebrüder Borntraeger.
- Schindewolf, O. H. (1950). *Grundfragen Der Paläontologie: Geologische Zeitmessung, Organische Stammesentwicklung, Biologische Systematik*. Stuttgart, Erwin Nägele.
- Schluter, D. (1984). Morphological and phylogenetic relations among the Darwin's finches. *Evolution*, 38, 921–930.
- Schluter, D. (2000). *The Ecology of Adaptive Radiation*. Oxford, U.K., Oxford University Press.
- Schmalhausen, I. I. (1935). Determination of basic concepts and methods of investigations of growth [in Russian]. In *The Growth of Animals*, ed. S. Kaplansky, Moscow, pp. 8–60.
- Schmalhausen, I. I. (1943). [Rate of evolution and factors which determine it]. *Zhurnal Obshchei Biologii, Moskva*, 4, 253–285 [Russian with English summary].
- Schmidt-Nielsen, K. (1984). *Scaling: Why Is Animal Size So Important?* Cambridge, Cambridge University Press.
- Schneider, D. C. (1994). Scale-dependent patterns and species interactions in marine nekton. In *Aquatic Ecology: Scale, Pattern, and Process*, eds. P. Giller and D. H. A. Rafaelli, London, Blackwell, pp. 441–467.
- Secord, R., Bloch, J. I., Chester, S. G. B., Boyer, D. M., Wood, A. R., Wing, S. L., Kraus, M. J., McInerney, F. A., and Krigbaum, J. (2012). Evolution of the earliest horses driven by climate change in the Paleocene-Eocene thermal maximum. *Science*, 335, 959–962.
- Seeley, R. H. (1986). Intense natural selection caused by a rapid morphological transition in a living marine snail. *Proceedings of the National Academy of Sciences USA*, 83, 6897–6901.
- Selander, R. K. and Johnston, R. F. (1967). Evolution in the house sparrow. *I. Intra-population variation in North America*. *Condor*, 69, 217–258.
- Sewertzoff, A. N. (1931). *Morphologische Gesetzmässigkeiten Der Evolution*. Jena, Gustav Fischer Verlag.
- Sheets, H. D. and Mitchell, C. E. (2001). Uncorrelated change produces the apparent dependence of evolutionary rate on interval. *Paleobiology*, 27, 429–445.
- Simberloff, D. S., Dayan, T., Jones, C., and Ogura, G. (2000). Character displacement and release in the small Indian mongoose, *Herpestes javanicus*. *Ecology*, 81, 2086–2099.
- Simpson, G. G. (1943). Criteria for genera, species, and subspecies in zoology and paleontology. *Annals of the New York Academy of Sciences*, 44, 145–178.
- Simpson, G. G. (1944). *Tempo and Mode in Evolution*. New York, Columbia University Press.
- Simpson, G. G. (1953). *The Major Features of Evolution*. New York, Columbia University Press 434.
- Slobodkin, L. B. (1961). *Growth and Regulation of Animal Populations*. New York, Holt, Reinhart, and Winston.
- Smith, F. A., Boyer, A. G., Brown, J. H., Costa, D. P., Dayan, T., Ernest, S. K. M., Evans, A. R., Fortelius, M., Gittleman, J. L., Hamilton, M. J., Harding, L. E., Lintulaakso, K.,

- Lyons, S. K., McCain, C., Okie, J. G., Saarinen, J. J., Sibly, R. M., Stephens, P. R., Theodor, J., and Uhen, M. D. (2010). The evolution of maximum body size of terrestrial mammals. *Science*, 330, 1216–1219.
- Smith, F. A., Browning, H., and Shepherd, U. L. (1998). The influence of climate change on the body mass of woodrats *Neotoma* in an arid region of New Mexico, USA. *Ecography*, 21, 140–148.
- Smith, T. B., Freed, L. A., Lepson, J. K., and Carothers, J. H. (1995). Evolutionary consequences of extinctions in populations of a Hawaiian honeycreeper. *Conservation Biology*, 9, 107–113.
- Sokal, R. R. and Rohlf, F. J. (1981). *Biometry*, Second Edition. San Francisco, W. H. Freeman.
- Spanbauer, T. L., Fritz, S. C., and Baker, P. A. (2018). Punctuated changes in the morphology of an endemic diatom from Lake Titicaca. *Paleobiology*, 44, 89–100.
- Spitzer, F. L. (1964). *Principles of Random Walk*. Princeton, Van Nostrand.
- St. Louis, V. L. and Barlow, J. C. (1991). Morphometric analysis of introduced and ancestral populations of the Eurasian tree sparrow. *VWilson Bulletin*, 103, 1:12.
- Stanley, S. M. (1973). An explanation for Cope's rule. *Evolution*, 27, 1–26.
- Stanley, S. M. (1975). A theory of evolution above the species level. *Proceedings of the National Academy of Sciences USA*, 72, 646–650.
- Stanley, S. M. (1989). The empirical case for the punctuational model of evolution. *Journal of Social and Biological Structures*,
- Stanley, S. M. and Yang, X. (1987). Approximate evolutionary stasis for bivalve morphology over millions of years: a multivariate, multilineage study. *Paleobiology*, 13, 113–139.
- Stearns, S. C. (1983). The genetic basis of differences in life history traits among six populations of mosquitofish (*Gambusia affinis*) that shared ancestors in 1905. *Evolution*, 37, 618–627.
- Stigler, S. M. (1986). *The History of Statistics: Measurement of Uncertainty Before 1900*. Cambridge, MA, Harvard University Press.
- Stuart, Y. E., Campbell, T. S., Hohenlohe, P. A., Reynolds, R. G., Revell, L. J., and Losos, J. B. (2014). Rapid evolution of a native species following invasion by a congener. *Science*, 346, 463–466.
- Sylvester-Bradley, P. C. (1959). Iterative evolution in fossil oysters. In *Proceedings of the 15th International Congress of Zoology*, pp. 193–197.
- Sylvester-Bradley, P. C. (1977). Biostratigraphical tests of evolutionary theory. In *Concepts and Methods of Biostratigraphy*, eds. E. G. Kauffman and J. E. Hazel, Stroudsburg, Pa., Dowden, Hutchinson and Ross, pp. 41–63.
- Szuma, E. (2003). Microevolutionary trends in the dentition of the Red fox (*Vulpes vulpes*). *Journal of Zoological Systematics and Evolutionary Research*, 41, 47–56.
- Taylor, B. L., Chivers, S. J., Larese, J., and Perrin, W. F. (2007). *Generation Length and Percent Mature Estimates for IUCN Assessments of Cetaceans*. Administrative Report LJ-07-01, La Jolla, California, Southwest Fisheries Science Center.
- Taylor, J. M. (1974). Morphological skull variation in the Australian wild rabbit: a multivariate analysis. Bachelor of Medical Science thesis, Perth, University of Western Australia, 182 pp.
- Taylor, J. M., Freedman, L., Olivier, T. J., and McCluskey, J. (1977). Morphometric distances between Australian wild rabbit populations. *Australian Journal of Zoology*, 25, 721–732.
- Theriot, E. C., Fritz, S. C., Whitlock, C., and Conley, D. J. (2006). Late Quaternary rapid morphological evolution of an endemic diatom in Yellowstone Lake, Wyoming. *Paleobiology*, 32, 38–54.

- Todor, K. (2016). The growth of bacterial populations. In *Online Textbook of Bacteriology*, Madison, University of Wisconsin, http://textbookofbacteriology.net/growth_3.html.
- Tseng, M., Kaur, K. M., Soleimani Pari, S., Sarai, K., Chan, D., Yao, C. H., Porto, P., Toor, A., Toor, H. S., and Fograscher, K. (2018a). Decreases in beetle body size linked to climate change and warming temperatures. *Journal of Animal Ecology*, 87, <http://dx.doi.org/10.1111/1365-2656.12789>.
- Tseng, M., Kaur, K. M., Soleimani Pari, S., Sarai, K., Chan, D., Yao, C. H., Porto, P., Toor, A., Toor, H. S., and Fograscher, K. (2018b). Decreases in beetle body size linked to climate change and warming temperatures. *Dryad Digital Repository*, <https://doi.org/10.5061/dryad.5164v>.
- Turner, J. R. G. (1986). The genetics of adaptive radiation: a neo-Darwinian theory of punctuational evolution. In *Patterns and Processes in the History of Life*, eds. D. M. Raup and D. Jablonski, Berlin, Springer, pp. 183–207.
- Uyeda, J. C., Hansen, T. F., Arnold, S. J., and Jason Pienaar. (2011). The million-year wait for macroevolutionary bursts. *Proceedings of the National Academy of Sciences USA*, 108, 15908–15913.
- Van Valen, L. M. (1974). Two modes of evolution. *Nature*, 252, 298–300.
- Van Valkenburgh, B. (1990). Skeletal and dental predictors of body mass in carnivores. In *Body Size in Mammalian Paleobiology: Estimation and Biological Implications*, eds. J. D. Damuth and B. J. MacFadden, Cambridge, Cambridge University Press, pp. 181–205.
- Vandenbergh, N. Hilgen, F. J. Speijer, R. P. Ogg, J. G. Gradstein, F. M. Hammer, O. Hollis, C. J., and Hooker, J. J. (2012). The Paleogene period. In *The Geological Time Scale 2012*, eds. F. M. Gradstein, J. G. Ogg, M. D. Schmitz, and G. M. Ogg, Amsterdam, Elsevier, pp. 855–921.
- Vermeij, G. J. (1977). The Mesozoic marine revolution: evidence from snails, predators, and grazers. *Paleobiology*, 3, 245–258.
- Vermeij, G. J. (1982). Environmental change and the evolutionary history of the periwinkle *Littorina littorea* in North America. *Evolution*, 36, 561–580.
- Villermé, L. R. (1829). Mémoire sur la taille de l'homme en France. *Annales d'Hygiene Publique et de Medecine Legale*, 1, 351–399.
- Vrba, E. S. (1985). Environment and evolution: alternative causes of the temporal distribution of evolutionary events. *South African Journal of Science*, 81, 229–236.
- Waagen, W. H. (1869). Die Formenreihe des *Ammonites subradiatus*: Versuch einer paläontologischen Monographie. *Geognostisch-Paläontologische Beiträge, München*, 2, 179–256.
- Waddington, C. H. (1939). *An Introduction to Modern Genetics*. London, George Allen and Unwin.
- Waddington, C. H. (1942). Canalization of development and the inheritance of acquired characters. *Nature*, 150, 563–565.
- Wedekind, R. (1920). Über Virenzperioden (Blüteperioden). *Sitzungsberichte der Gesellschaft zur Beförderung der gesamten Naturwissenschaften zu Marburg*, 1920 (2), 18–31.
- Weidenreich, F. (1943). The skull of *Sinanthropus pekinensis*: a comparative study on a primitive hominid skull. *Palaeontologica Sinica, New Series D*, 10, 1–485.
- Weismann, A. (1872). *Ueber Den Einfluss Der Isolirung Auf Die Artbildung*. Leipzig, Wilhelm Engelmann 108.
- Weismann, A. (1883). *Über Die Vererbung*. Jena, Gustav Fischer.
- Weldon, W. F. R. (1893). On certain correlated variations in *Carcinus maenas*. *Proceedings of the Royal Society of London*, 54, 318–329.

- Weldon, W. F. R. (1895). Attempt to measure the death-rate due to the selective destruction of *Carcinus moenas* with respect to a particular dimension. *Proceedings of the Royal Society of London*, 57, 360–379.
- Wiedmann, J. (1973). Evolution or revolution of ammonoids at Mesozoic system boundaries? *Biological Reviews, Cambridge*, 48, 159–194.
- Wiens, J. A. (1989). Spatial scaling in ecology. *Functional Ecology*, 3, 385–397.
- Williams, H. S. (1910). Persistence of fluctuating variations as illustrated by the fossil genus *Rhipidomella*. *Geological Society of America Bulletin*, 21, 295–312.
- Williamson, P. G. (1981). Palaeontological documentation of speciation in Cenozoic molluscs from Turkana Basin. *Nature*, 293, 437–443.
- Williamson, S. H. (2017). *Daily Closing Value for the Dow Jones Average, 1885 to Present*. <https://measuringworth.com/DJA/>.
- Wood, A. R., Zelditch, M. L., Rountrey, A. N., Eiting, T. P., Sheets, H. D., and Gingerich, P. D. (2007). Multivariate stasis in the dental morphology of the Paleocene-Eocene condylarth *Ectocion*. *Paleobiology*, 33, 248–260.
- Wright, S. (1931). Evolution in Mendelian populations. *Genetics*, 16, 97–159.
- Wright, S. (1968). *Evolution and the Genetics of Populations. Volume 1: Genetic and Biometric Foundations*. Chicago, University of Chicago Press.
- Wright, S. (1969). *Evolution and the Genetics of Populations. Volume 2: The Theory of Gene Frequencies*. Chicago, University of Chicago Press.
- Yablokov, A. V. (1974). *Variability of Mammals*. New Delhi, Amerind Publishing.
- Yarwood, C. E. (1956). Generation time and the biological nature of viruses. *American Naturalist*, 40, 97–102.
- Yule, G. U. (1924). A mathematical theory of evolution, based on the conclusions of Dr. J. C. Willis, F.R.S. *Philosophical Transactions of the Royal Society of London, Series B*, 213, 21–87.
- Zakrzewski, R. J. (1969). The rodents from the Hagerman local fauna, upper Pliocene of Idaho. *Contributions from the Museum of Paleontology, University of Michigan*, 23, 1–36.
- Zeleny, C. (1922). The effect of selection for eye facet number in the white bar-eye race of *Drosophila melanogaster*. *Genetics*, 7, 1–115.
- Zeng, Z. and Brown, J. H. (1987). Population ecology of a desert rodent: *Dipodomys merriami* in the Chihuahuan Desert. *Ecology*, 68, 1328–1340.
- Zeuner, F. E. (1946). *Dating the Past: An Introduction to Geochronology*. London, Methuen and Company.
- Ziegler, A. M. (1966). The Silurian brachiopod *Eocoelia hemisphaerica* (J. de C. Sowerby) and related species. *Palaeontology*, 9, 523–543.
- Zimmermann, W. (1953). *Evolution: Die Geschichte Ihrer Probleme Und Erkenntnisse*. Freiburg, Karl Alber.