

Literatura

- AO, M., MAE, O., NAMBA, Y., et al.: Perforator based flap for coverage of lumbosacral defects. *Plast. Reconstr. Surg.*, 101, 1998, č. 4, s. 987–991.
- ARRONDO, D. I., HUIZI, E. X., GALA DE ANDREAS, M., et al.: Decubitus ulcers in intensive care units. Analysis and care. *Enferm. Intensiva*, 6, 1995, č. 4, s. 159–164.
- BAILEY, B. N.: *Bedsores*. London, Edward Arnold 1967, s. 97
- BARASH, P. G., CULLEN, B. F., STOELTING, R. K.: *Clinical anesthesia*. 31, 1992, s. 271–273.
- BARKER, D. E., ELKINS, C. W., POER, D. H.: Methods of closure of decubitus ulcers in the paralyzed patient. *Ann. Surg.*, 123, 1946, s. 523–525.
- BARKER, D. E.: Late results in surgery for decubitus ulcers. *Amer. J. Surg.*, 74, 1947, s. 180–182.
- BENEŠ, V.: *Poranění míchy*. Praha, Avicenum 1961, s. 189.
- BERKAS, E. M., MERRILL, D., CHESLER, M. D., et al.: Multiple decubitus ulcers treatment by hip disarticulation and soft tissue flaps from the lower limbs. *Plast. Reconstr. Surg.*, 27, 1961, s. 618–626.
- BLOCKSMA, R., KOSTRUBALA, J. G., GREELEY, P. W.: The surgical repair of decubitus ulcers in paraplegics: Further observations. *Plast. Reconst. Surg.*, 4, 1949, s. 123–130.
- BOSSE, A., WULF, M., WIETHEGE, T., et al.: Collagens and growth factors in heterotopic ossification. *Pathologe*, 15, 1994, č. 4, s. 216–225.
- BRAVO, C., ONDROVČÍK, P., VÁLKA, J.: Sledování účinku nejčastěji používaných roztoků a mastí v běžné chirurgické praxi v chirurgii. *Rozhl. Chir.*, 74, 1995, s. 273–276.
- BRODANOVÁ, M.: *Úvod do infúzní terapie a výživy*. Praha, Avicenum 1984, s. 94–99.
- CLARK, M., WATTS, S.: The incidence of pressure sores within a National Health Service Trust hospital during 1991. *J. Adv. Nurs.*, 20, 1994, s. 309–402.
- CROCE, E. J., SCHULLINGER, R. N., SHEARER, T. P.: Operative treatment of decubitus ulcer. *Ann. Surg.*, 123, 1946, s. 53–57.
- CULEY, F.: Nursing aspects of pressure sore prevention and therapy. *Br. J. Nurs.*, 7, 1998, č. 15, s. 879–882.
- CUZZELL, J.: Wound care forum for pressure sores. *Amer. J. Nurs.*, 8, 1986, s. 923–924.
- DANIEL, R. K., TERZIS, J. K., CUNINGHAM, D. M.: Sensory skin flaps for coverage of pressure sores in paraplegic patients. *Plast. Reconstr. Surg.*, 58, 1976, s. 317–328.
- DANSEREAU, J. G., CONWAY, H.: Closure of decubiti in paraplegics. Report on 2000 cases. *Plast. Reconstr. Surg.*, 33, 1964, s. 474–480.
- DAVIS, J. S.: The operative treatment of scars following bedsores. *Surg.*, 3, 1938, s. 21–25.
- DIBBELL, D. G.: Use of a long island flap to bring sensation to the sacral area in young paraplegics. (Case report). *Plast. Reconstr. Surg.*, 54, 1974, s. 220–223.

- DISA, J. J., CARLTON, J. M., GOLDBERG, N. H.: Efficacy of operative cure in pressure sore patient. *Plast. Reconstr. Surg.*, 89, 1992, s. 272–278.
- DUMURGIER, C., PUJOL, C., CHEVALLEY, J., et al.: Pressure sore carcinoma. *Paraplegia*, 6, 1991, s. 390–395.
- EDWARDS, M.: The rationale for the use of risk calculators in pressure sore prevention and the evidence of the reliability and validity of published scales. *J. Adv. Nurs. (Anglie)*, 20, 1994, s. 288–296.
- ESPOSITO, G., CAPRIO, G. di, ZICCARDI, P., et al.: Tissue expansion in the treatment of pressure ulcers. *Plast. Reconstr. Surg.*, 88, 1991, s. 501–507.
- GEORGIADE, N., PICKRELL, K., MAGUIRE, C.: Total tigh flaps for extensive decubitus ulcers. *Plast. Reconstr. Surg.*, 17, 1956, s. 220.
- GOLDBERG, J. A., BARWICK, W. J., LEVIN, L. S.: Restoration of sensation in paraplegia by a sensory innervated plantar fillet free flap. (Case Report). *Paraplegia*, 33, 1995, s. 116–119.
- GRABB, W. C., SMITH, J.: *Plastic Surgery*, Lippincott-Raven Pulol 1997.
- GRIFFITH, B. H., SCHULTZ, R. C.: The prevention and surgical treatment of recurrent decubitus. *Plast. Reconstr. Surg.*, 27, 1961, s. 235–241.
- HAMMERSHOY, E. M., SPERLING, L. S., HENRIKSEN, T. B., et al.: Quality assurance regarding pressure sores in a Danish hospital population. *Ugeskr. Laeger*, 156, 1994, s. 4229–4233.
- HAUGE, E. N.: The anatomical basis of a new method for re-innervation of the gluteal region in paraplegics. *A. Physiol. Scand. (Suppl.)*, 1991, č. 603, s. 19–21.
- HAYASHI, A., MARUYAMA, Y., SAZE, M., et al.: *Br. J. Plast. Surg.*, 51, 1998, č. 2, s. 113 až 117.
- HUNG-CHIC CHEN,, CHAN-JIN, W., NOORDHOFF, M. S.: Coverage of multiple extensive pressure sores with a single filled lower leg myocutaneus free flap. *Plast. Reconstr. Surg.*, 7, 1986, s. 396–399.
- HURWITZ, D. J., SWATZ, W. M., MATTHES, S. J.: The gluteal thigh flap. A riable and sensate flap for the closure of buttock and perineal wounds. *Plast. Reconstr. Surg.*, 68, 1981, s. 521–523.
- JARKOVSKÝ, J.: Rozbor proleženin gerontopsychiatrických nemocných. *Vnitř. Lék.*, 26, 1980, s. 279–281.
- KÁBRTOVÁ, A.: Rehabilitační soustředění pacientů po poranění míchy. *Přednáška*. Kováčová, 1994.
- KIERNEY, P. C., ENGRAV, L. H., ISIK, F. F., et al.: Results of 268 pressure sores in 158 patients managed jointly by plastic surgery and rehabilitation medicine. *Plast. Reconstr. Surg.*, 102, 1998, č. 3, s. 765–772.
- KOSTRUBALA, J. C., GREELEY, P. W.: The problem of decubitus ulcers in paraplegics. *Plast. Reconstr. Surg.*, 2, 1947, s. 403–409.
- KUIPERS, T., STARK, G. B., SPILKER, G.: Costs and long-term results of plastic surgery treatment of decubitus ulcers in paraplegic patients. *Handchir. Microchir. Plast. Chir.*, 27, 1995, č. 3, s. 161–165.
- LAMON, J. D. Jr., ALEXANDER, E. Jr.: Secondary closure of decubitus ulcers with the aid fo penicillin. *JAMA*, 127, 1945, s. 396–393.
- LUSCHER, M. J., ROCHE, R. de, KRUPP, S., et al.: The sensory tensor fasciae latae flap: a 9-year follow up. *Ann. Plast. Surg.*, 26, 1991, s. 306–310.
- MACKINNON, S. E., DELLON, A. L., PATTERSON, G. A.: Medial antebrachial cutaneous-lateral femoral cutaneous neurotisation to provide sensation to pressure-bearing areas in paraplegic patient. *Ann. Plast. Surg.*, 14, 1985, s. 541–544.
- MARUYAMA, Y., NAKAJIMA, H., WADA, M., et al.: A gluteus maximus myocutaneus island flap for the repair of a sacral decubitus ulcer. *Brit. J. Plast. Surg.*, 33, 1980, s. 150.

- MASSE, P.: La place du lambeau de rotation dans le traitement des escarres sacrées des paraplegiques. Indications, technique, résultats. Ann. Chir. Plast., 5, 1960, s. 39–48.
- MEEHAN, M.: National pressure ulcer prevalence survey. Adv. Wound Care, 7, 1994, č. 3, s. 27–30.
- MILLER, I.: Ústní sdělení.
- MINAMI, R. T., MILLS, R., PARDOVE, R.: Gluteus maximus myocutaneus flaps for repair of pressure sores. Plast. Reconstr. Surg., 60, 1977, s. 242–245.
- MORRIS, A. G., WATSON, A. H., STEVENSON, J. H.: Complications of plastic surgery. London, Bailliere Tindall 1989, s. 413–417.
- MÜLLEROVÁ, N.: Nový přístup k prevenci a ošetřování dekubitů – další aspekt kvality péče ve FN v Plzni. Ošetřovatelská péče, Listy asociací nemocnic, 2, 1996, s. 25–27.
- NAHAI, F., SIVERTON, J. S., HILL, H. L., et al.: The tensor fascia lata musculocutaneous. Ann. Plast. Surg., 1, 1979, s. 372–374.
- NEJEDLÝ, B., et al.: Vnitřní prostředí, klinická biochemie a praxe. Praha, Avicenum 1980, s. 496–499.
- NIETZEL, W.: Den richtigen Greif gefunden? Rotations-Therapie ohne wirksame Druckentlastung undenkbar! Krankenpf., 35, 1997, č. 3, s. 82–83.
- NOLA, G. T., VISTNERS, L. M.: Differential response of skin and muscle in the experimental production of pressure sores. Plast. Reconstr. Surg., 66, 1980, s. 728–733.
- NUNN, J. F.: Anaesthesia for patients with neuromuscular disorders. In: NUNN, J. F.: General anaesthesia. London, Butterworths 1989, s. 741–742.
- PARISH, L. C., WITKOVSKI, J. A., CREISSEY, J. C.: The decubitus Ulcer in Clinical Practice. Berlin-Heidelberg, Springer Verlag 1997.
- PARRY, S. W., MATTHES, S. J.: Bilateral gluteus maximus myocutaneus advancement flaps: Sacral coverage for ambulatory patients. Amer. Plast. Surg., 8, 1982, s. 443–445.
- REED, M. J.: Wound repair in older patients: preventing problems and managing the healing. Geriatrics, 53, 1998, č. 5, s. 88–94.
- RIEBELOVÁ, V.: Proleženiny – tlakové vředy, poznatky o jejich etiologii, patologické anatomii a patofyziologii. Rozh. Chir., 54, 1975, s. 693–697.
- RIEBELOVÁ, V.: Poznámky k prevenci a konzervativnímu léčení dekubitálních vředů. Rozhl. Chir., 56, 1977, s. 514–518.
- RIEBELOVÁ, V.: The tensor fasciae latae muschulocutaneous flap on operations for lateralis and nervus saphenus. A preparatory study. Acta Chir. Plast., 27, 1985, s. 104–112.
- ROYER, J., PICKRELL, K., GEORGIADE, N., et al.: Total thigh flaps for extensive decubitus ulcers. A 16 years review of 41 total thigh flaps. Plast. Reconstr. Surg., 43, 1969, s. 2.
- RODRIGUEZ, G. P., CLAUS-WALKER, J.: Biochemical changes in skin composition in spinal cord injury. A possible contribution to decubitus ulcers. Paraplegia, 26, 1985, s. 108–119.
- RUBIN, J. A., WHETZEL, T. P., STEVENSON, T. R.: The posterior thigh fasciocutaneous flap: Vascular anatomy and clinical application. Plast. Reconstr. Surg., 95, 1995, s. 1228–1239.
- SALZBERG, C. A., BYRNE, D. W., CAYTEN, C. G., et al.: A new pressure ulcer risk assessment scale for individuals with spinal cord injury. Amer. J. Phys. Med. Rehabil., 75, 1996, s. 96–104.
- STELLY, T., C., McNEIL, J., W., SNYPES, S., R., et al.: Hemicorporectomy. Clin. Anat., 8, 1995, č. 2, s. 116–123.
- STOCKAROVÁ, D., PILNÁČEK, J.: Our experience with use of musculocutaneous flaps in trochanter decubitus ulcers. Acta Chir. Plast., 26, 1984, s. 238–245.

- STOCKAROVÁ, D., PILNÁČEK, J., RUBÍN, J.: Možnosti léčby sakrálních dekubitů muskulokutánními laloky. *Rozhl. Chir.*, 65, 1986, s. 85–92.
- STOCKAROVÁ, D., PILNÁČEK, J., RUBÍN, J.: Přínos muskulokutánních laloků v léčení chronických osteomyelitid bérce. *Rozhl. Chir.*, 65, 1986, s. 93–102.
- SUMIYA, T., KAWAMURA, K., TOKUHIRO, A., et al.: A survey of wheel chair use by paraplegic individuals in Japan. Part 1: Characteristics of wheelchair cushions. *Spinal Cord*, 35, 1997, č. 9, s. 590–594.
- SUMIYA, T., KAWAMURA, K., TOKUHIRO, A., et al.: A survey of wheel chair use by paraplegic individuals in Japan. Part 2: Prevalence of pressure sores. *Spinal Cord*, 35, 1997, č. 9, s. 595–598.
- THOMPSON, C. E., RICE, M. L. Jr.: Secondary amyloidosis in spinal cord injury. *Ann. Int. Med.*, 31, 1949 s. 1057.
- TOBIN, G. R., SANDERS, B. P., WEINER, L. J.: The biceps femoris myocutaneous advancement flap: as useful modification for ischial pressure ulcer reconstruction. *Ann. Plast. Surg.*, 6, 1981, s. 39.
- WENDSCHE, P., et al.: Poranění páteře a míchy. Komplexně ošetřovatelská péče u paraplegiků. Brno, IDVPZ 1993, s. 189.
- WENDSCHE, P.: Současný trend léčení poranění krční páteře. *Zprav. Úraz. Chir.*, 3, 1995, s. 1–6.
- WESBECHER, M., BARBER, S., VIVIANS, R., et al.: Treatment of a young adult paraplegic with multiple stage III and IV pressure ulcers. *J. Wound. Ostomy. Continence Nurs.*, 25, 1998, č. 1, s. 55–58.
- WINGATE, G. B., FRIEDLAND, J. A.: Repair of ischial pressure ulcers with gracilis myocutaneous island flaps. *Plast. Reconstr. Surg.*, 62, 1978, s. 245–250.
- WHITE, J. C., HAMM, W. G.: Primary closure of bedsores by plastic surgery. *Ann. Surg.*, 124, 1946, s. 1136–1140.
- YAMAMOTO, Y., TSUTSUMIDA, A., MURAZUMI, M., et al.: Long-term outcome of pressure sores treated with flap coverage. *Plast. Reconstr. Surg.*, 100, 1997, č. 5, s. 1212 až 1217.
- YUAN, R. T. W.: The use of tissue expansion of lower extremity wounds in paraplegic patients. *Plast. Reconstr. Surg.*, 5, 1989, s. 892–895.
- ZADÁK, Z.: Parenterální výživa v klinické praxi. Hradec Králové, Karlova Univerzita 1984, s. 138–154.
- ZHOU, G., QIAO, Q., CHEN, G. Y., et al.: Clinical experience and surgical anatomy of 32 free anterolateral thigh flap transplantations. *Brit. J. Plast. Surg.*, 2, 1991, s. 91–96.