

---

## Contents

|  |     |
|--|-----|
| Preface .....  | VII |
| Bachmann, W. (Kronach); Lotz, N.; Sieger, C. (Munich): Insulin/Sulphonylurea Combinations – A New Beginning? .....   | 1   |
| <i>Clinico-Experimental Studies</i>  |     |
| Klujko, J.; Schmidt, F. H.; Kühnle, H. F; Reiter, J. (Mannheim): Investigation of the Mechanism of Action of Glibenclamide and Insulin Administered in Combination in Type-II Diabetic Patients with Secondary Failure to Oral Treatment .....   | 13  |
| Hasche, H. (Rotenburg/Wümme): Combination Therapy with Different Doses of Glibenclamide. A Double-Blind Study on 36 Patients with Secondary Failure .....  | 25  |
| Schwarz, O. (Mannheim); Rosak, C.; Althoff, P. H.; Schöffling, K. (Frankfurt/M.); Schmidt, F. H. (Mannheim): Combined Treatment of Type-II Diabetic Patients with Insulin and Glibenclamide after Secondary Failure to Tablets. A Double-Blind, Placebo-Controlled Crossover Study ..... | 33  |
| Stahl, T. (Freudenberg): Combination of Regular Insulin with Glibenclamide in Secondary Failure .....  | 44  |
| Henrichs, H. R. (Quakenbrück): Sulphonylurea/Insulin Combination in Diabetes Mellitus Following Secondary Failure to Tablets. Results of a Double-Blind Crossover Study of Glibenclamide .....   | 51  |
| Willms, B.; Boustani, A. (Bad Lauterberg): Comparison of Pre-Prandial Administration of Regular Insulin with Administration of NPH Insulin at Bed Time and of Conventional Insulin Therapy in Combination Therapy with Insulin/Sulphonylurea in Type-II Diabetes .....                   | 68  |
| Bottermann, P. (Munich): Insulin Treatment in Type-II Diabetes in Presence of Secondary Failure to Sulphonylurea Therapy. A Contribution to the Discussion .....   | 78  |
| Bergis, K. H. (Bad Mergentheim): Insulin by Night – Sulphonylurea by Day: A Practical Compromise for the Stabilization on Insulin in Type-II Diabetics with Secondary Sulphonylurea Failure .....  | 85  |

*Long-Term Clinical Trials*

|  |     |
|--|-----|
| Knick, B.; Röckel, A.; Panitz, N.; Groth, U. (Wiesbaden): Comparative Study of SU/Insulin (with Special Consideration of a Group of Kidney-Failure Patients) versus Combined Metformin/Insulin Therapy in Type-II Diabetes | 94  |
| Sachse, G. (Wiesbaden); Mäser, E.; Federlin, K. (Gießen): Long-Term Results of Combined Insulin/Sulphonylurea Therapy: Results, Critical Assessment, Clinical Consequences .....   | 103 |
| Bachmann, W. (Kronach); Lotz, N.; Mehnert, H. (Munich); Rosak, C.; Schöffling, K. (Frankfurt/M.): Controlled Multicenter Double-Blind Clinical Study of the Efficacy of Combined Treatment .....                           | 109 |
| Stürmer, W. (Würzburg); Schrezenmeir, J. (Mainz); Kasper, H. (Würzburg): Long-Term Results of Combined Sulphonylurea/Insulin Therapy in Patients with a Positive Glibenclamide Test under Clamp Conditions .....           | 121 |
| Lotz, N. (Munich); Bachmann, W. (Kronach); Mehnert, H. (Munich): Combination of Sulphonylurea and Insulin in the Long-Term Treatment of 'Secondary Failure' in Type-II Diabetes .....                                      | 131 |
| Petzold, R.; Klein, E. (Bad Oeynhausen): Insulin/Sulphonylurea Therapy in 'Secondary Failure' – Clinical Observations .....  | 146 |
| Klein, W. (Berlin): Combined Sulphonylurea/Insulin Treatment in Secondary Failure .....  | 151 |
| Bachmann, W. (Kronach); Lotz, N.; Mehnert, H. (Munich): Concluding Remarks   | 154 |