

CONTENTS

Studies on protozoan parasite of Australian fishes I. New species of the genera <i>Coccomyxa</i> Léger et Hesse, 1907, <i>Ortholinea</i> Shulman, 1962 and <i>Kudoa</i> Meglitsch, 1947 (Myxozoa, Myxosporida)	289–306
J. LOM, K. ROHDE, I. DYKOVÁ	289–306
Biometric variability of <i>Proteocephalus neglectus</i> (Cestoda: Proteocephalidae) in two different age groups of the rainbow trout from the Dobšiná water reservoir (East Slovakia)	307–316
V. HANZLOVÁ, M. ŠPAKULOVÁ	307–316
Redescription of <i>Proteocephalus neglectus</i> La Rue, 1911 (Cestoda: Proteocephalidae), a trout parasite, including designation of its lectotype	317–323
V. HANZLOVÁ, T. SCHOLZ	317–323
Nematode parasites of fishes of the Paraná River, Brazil. Part I. Trichuroidea, Oxyuroidea and Cosmocercoidae	327–353
F. MORAVEC, A. KOHN, B. M. FERNANDES	327–353
A scanning electron microscope study of <i>Toxocara genettae</i> Warren 1972 (Ascaridae), with data on morphometric variation	355–367
M. L. SANMARTÍN, F. ALVAREZ, H. GUÓN BOTELLA, R. IGLESIAS, J. ESTÉVEZ, R. LÓPEZ-ROMÁN	355–367
<i>Strongyloides ophiussensis</i> sp. n. (Nematoda: Strongylidae), parasite of an insular lizard, <i>Podarcis pityusensis</i> (Sauria: Lacertidae)	369–373
V. ROCA, M. J. HORNERO	369–373
Immunization of rabbits with antigens from <i>Psoroptes cuniculi</i> , the rabbit scab mite	375–382
J. UHLÍŘ	375–382

Continued on inside cover