

CONTENTS

ARTICLES

- 641 Monoclonal antibody against Rab8 from *Bombyx mori* (Lepidoptera: Bombycidae) — T. UNO, T. NAKADA, Y. UNO, K. KANAMARU, H. YAMAGATA, M. NAKAMURA & M. TAKAGI
- 647 Linkage analysis of the visible mutations *Sel* and *Xan* of *Bombyx mori* (Lepidoptera: Bombycidae) using SSR markers — X. MIAO, M. LI, F. DAI, C. LU, M.R. GOLDSMITH & Y. HUANG
- 653 Effects of polymorphic Robertsonian rearrangements on the frequency and distribution of chiasmata in the water-hyacinth grasshopper, *Cornops aquaticum* (Orthoptera: Acrididae) — P.C. COLOMBO
- 661 Direction of karyotype evolution in the bug family Nabidae (Heteroptera): New evidence from 18S rDNA analysis — C. NOKKALA, V. KUZNETSOVA, S. GROZEEVA & S. NOKKALA
- 667 Morphological and molecular evidence of a possible hybrid zone of *Leptidea sinapis* and *L. reali* (Lepidoptera: Pieridae) — R. VEROVNIK & P. GLOGOVČAN
- 675 The effects of Pleistocene glaciations on the phylogeography of *Melitaea cinxia* (Lepidoptera: Nymphalidae) — N. WAHLBERG & I. SACCHERI
- 685 Crustacean red pigment-concentrating hormone Panbo-RPCH affects lipid mobilization and walking activity in a flightless bug *Pyrrhocoris apterus* (Heteroptera) similarly to its own AKH-peptides — R. SOCHA, D. KODRÍK & R. ZEMEK
- 693 The role of temperature and photoperiod in diapause induction in the brassica leaf beetle, *Phaedon brassicae* (Coleoptera: Chrysomelidae) — X.-P. WANG, F.-S. XUE, Y.-Q. TAN & C.-L. LEI
- 699 Diapause, cold hardiness and flight ability of Cry1Ac-resistant and -susceptible strains of *Helicoverpa armigera* (Lepidoptera: Noctuidae) — G.-M. LIANG, K.-M. WU, B. RECTOR & Y.-Y. GUO
- 705 Effect of photoperiod on parasitization by *Trichogramma principium* (Hymenoptera: Trichogrammatidae) — S.Y.A. REZNIK & N.P. VAGHINA
- 715 Mating periodicity and post-mating refractory period in the zoophytophagous plant bug *Macrolophus caliginosus* (Heteroptera: Miridae) — C. GEMENO, O. ALOMAR, J. RIUDAVETS & C. CASTAÑÉ
- 721 Adult feeding increases fecundity in female *Helicoverpa armigera* (Lepidoptera: Noctuidae) — Z.-M. SONG, Z. LI, D.-M. LI, B.-Y. XIE & J.-Y. XIA
- 725 Ejaculate size varies with remating interval in the grasshopper *Chorthippus parallelus erythropus* (Caelifera: Acrididae) — K. REINHARDT
- 731 Microhabitat preferences of *Maculinea teleius* (Lepidoptera: Lycaenidae) in a mosaic landscape — P. BATÁRY, N. ÖRVÖSSY, Á. KÖRÖSI, M. VÁLYI NAGY & L. PEREGOVITS
- 737 The multicolored Asian ladybeetle *Harmonia axyridis* (Coleoptera: Coccinellidae) in Quebec agroecosystems ten years after its arrival — É. LUCAS, C. VINCENT, G. LABRIE, G. CHOUINARD, F. FOURNIER, F. PELLETIER, N.J. BOSTANIAN, D. CODERRE, M.-P. MIGNAULT & P. LAFONTAINE
- 745 Size-dependent colouration in larvae of *Orgyia antiqua* (Lepidoptera: Lymantriidae): A trade-off between warning effect and detectability? — S.-L. SANDRE, T. TAMMARU & T. MÄND
- 753 Probability distribution, sampling unit, data transformations and sequential sampling of European vine moth, *Lobesia botrana* (Lepidoptera: Tortricidae) larval counts from Northern Greece vineyards — A.A. IFOULIS & M. SAVOPOULOU-SOULTANI
- 763 Climate signals are reflected in an 89 year series of British Lepidoptera records — R.L.H. DENNIS & T.H. SPARKS
- 769 Body weight distributions of central European Coleoptera — W. ULRICH
- 777 Spermathecal morphology and sperm dynamics in the female *Schizocosa malitiosa* (Araneae: Lycosidae) — G. USETA, B.A. HUBER & F.G. COSTA
- 787 Flightless Hawaiian Hemerobiidae (Neuroptera): Comparative morphology and biology of a brachypterous species, its macropterous relative and intermediate forms — C.A. TAUBER, M.J. TAUBER & J.G. GIFFIN
- 801 Revision of *Afrocrania* (Coleoptera: Chrysomelidae: Galerucinae) Part II: Species in which the males lack head cavities or extended elytral extrusions — T. WAGNER
- 815 Phylogenetic revision of the New Zealand genus *Dasytricheta*, with a discussion of its systematic position within the subfamily Aleocharinae (Coleoptera: Staphylinidae) — G. PAŚNIK
- 827 *Pristinochterus* gen. n. (Hemiptera: Ochteridae) from the Upper Mesozoic of northeastern China — Y. YAO, W. CAI & D. REN
- 837 A new Middle Jurassic "grylloblattodean" family from China (Insecta: Juraperlidae fam. n.) — D.-Y. HUANG & A. NEL