

CONTENTS

- 598 Editorial — P. SASKA

ARTICLES

- 601 Elytral surface structure in *Poecilus lepidus* (Coleoptera: Carabidae): What about the nature of its inheritance? — D. MOSSAKOWSKI & W. PAARMANN
- 608 The effects of overwintering, sex, year, field identity and vegetation at the boundary of fields on the body condition of *Anchomenus dorsalis* (Coleoptera: Carabidae) — E. BARANOVSKÁ, M. KNAPP & P. SASKA
- 615 Weed seed choice by carabid beetles (Coleoptera: Carabidae): Linking field measurements with laboratory diet assessments — S. PETIT, A. BOURSault & D.A. BOHAN
- 621 Thermal constants of egg development in carabid beetles — variation resulting from using different estimation methods and among geographically distant European populations — P. SASKA, M. VLACH, J. SCHMIDTOVÁ & A.V. MATALIN
- 631 Prey detection in carabid beetles (Coleoptera: Carabidae) in woodland ecosystems by PCR analysis of gut contents — L. ŠERIĆ JELASKA, D. FRANJEVIĆ, S.D. JELASKA & W.O.C. SYMONDSOHN
- 639 Consequence of the transformation of a primeval forest into a managed forest for carabid beetles (Coleoptera: Carabidae) — a case study from Białowieża (Poland) — J. SKŁODOWSKI
- 649 Quantifying predation pressure along an urbanisation gradient in Denmark using artificial caterpillars — M. FERRANTE, A. LO CACCIATO & G.L. LÖVEI
- 655 The effect of the construction and renovation of a highway bypass in Central Poland on the carabid beetle fauna (Coleoptera: Carabidae) — I. DYMITYRSZYN
- 663 Dispersal of individuals of the flightless grassland ground beetle, *Carabus hungaricus* (Coleoptera: Carabidae), in three populations and what they tell us about mobility estimates based on mark-recapture — Z. ELEK, L. DRAG, P. POKLUDA, L. ČÍŽEK & S. BÉRCES
- 669 Effect of conventional and non-inversion tillage systems on the abundance and some life history traits of carabid beetles (Coleoptera: Carabidae) in winter triticale fields — A. KOSEWSKA, T. SKALSKI & M. NIETUPSKI
- 677 Changes in carabid beetle fauna (Coleoptera: Carabidae) along successional gradients in post-industrial areas in Central Poland — A. SCHWERK
- 686 Edge effects on ground-dwelling beetles (Carabidae and Staphylinidae) in oak forest-forest edge-grassland habitats in Hungary — B. TÓTHMÉRÉSZ, D.D. NAGY, S. MÍZSER, D. BOGYÓ & T. MAGURA
- 692 Form, function and evolutionary significance of stridulatory organs in ant nest beetles (Coleoptera: Carabidae: Paussini) — A. DI GIULIO, S. FATTORINI, W. MOORE, J. ROBERTSON & E. MAURIZI
- 703 Benign neglect enhances urban habitat heterogeneity: Responses of vegetation and carabid beetle (Coleoptera: Carabidae) to the cessation of mowing of park lawns — S. VENN & D.J. KOTZE
- 715 Old forest edges may promote the distribution of forest species in carabid assemblages (Coleoptera: Carabidae) in Croatian Forests — A. BRIGIĆ, M. STARČEVIĆ, B. HRAŠOVEC & Z. ELEK

NOTE

- 726 Isolation and characterization of eleven polymorphic microsatellite markers from the beneficial carabid beetle, *Poecilus cupreus* (Coleoptera: Carabidae), and genetic structuring among three populations from western France — R. MARREC, S. RUAUT, C. RIBOUT, M. PLANTEGENEST & B. GAUFFRE