

CONTENTS

ARTICLES

- 501 Genetic variation in East-Adriatic populations of the Asian tiger mosquito, *Aedes albopictus* (Diptera: Culicidae), inferred from *NADH5* and *COI* sequence variability — T. ŽITKO, A. KOVACIĆ, Y. DESDEVISES & J. PUZINA
- 509 Comparative analysis of the location of rDNA in the Palaearctic bushcricket genus *Isophya* (Orthoptera: Tettigoniidae: Phaneropterinae) — B. GRZYWACZ, A. MARYAŃSKA-NADACHOWSKA, D.P. CHOBANOV, T. KARAMYSHEVA & E. WARCHALOWSKA-ŚLIWA
- 519 The genetic structure of phenologically differentiated Large Blue (*Maculinea arion*) populations (Lepidoptera: Lycaenidae) in the Carpathian Basin — J. BERECZKI, J.P. TÓTH, A. TÓTH, E. BÁTORI, K. PECSENYE & Z. VARGA
- 529 One variable species or multiple cryptic? Mitochondrial phylogeny of Central and North American *Chlosyne lacinia* (Lepidoptera: Nymphalidae) — T.C. BONEBRAKE, W.B. WATT, A. PEREZ & C.L. BOGGS
- 537 Divergent patterns in the mitochondrial and nuclear diversity of the specialized butterfly *Plebejus argus* (Lepidoptera: Lycaenidae) — M. SIELEZNIEW, D. PONIKWICKA-TYSZKO, M. RATKIEWICZ, I. DZIEKAŃSKA, A. KOSTRO-AMBROZIAK & R. RUTKOWSKI
- 547 Effect of insect-resistant transgenic maize on growth and development, utilization of nutrients and in vivo activity of the detoxification enzymes of the Asian corn borer, *Ostrinia furnacalis* (Lepidoptera: Pyralidae) — M.J. SHI, P.L. LU, X.L. SHI & Y.Z. YANG
- 553 Molecular phylogenetic analysis of a scale insect (*Drosicha mangiferae*; Hemiptera: Monophlebidae) infesting mango orchards in Pakistan — M. ASHFAQ, J. ARA, A.R. NOOR, P.D.N. HEBERT & S. MANSOOR
- 561 Cooperative defence of brood in *Nicrophorus quadripunctatus* (Coleoptera: Silphidae) — S. SUZUKI
- 565 Two ways of finding a host: A specialist and a generalist parasitoid species (Hymenoptera: Chalcidoidea: Pteromalidae) — R.S. PETERS
- 575 *Praon volucre* (Hymenoptera: Braconidae: Aphidiinae), a natural enemy of *Macrosiphum euphorbiae* (Hemiptera: Aphidiidae): Life table and intrinsic rate of population increase — J.C. LINS JR, V.H.P. BUENO, D.B. SILVA, M.V. SAMPAIO & J.C. VAN LENTEREN
- 581 Seed preferences of *Pyrrhocoris apterus* (Heteroptera: Pyrrhocoridae): Are there specialized trophic populations? — M. KRISTENOVÁ, A. EXNEROVÁ & P. ŠTYS
- 587 Effect of the size of the pupae, adult diet, oviposition substrate and adult population density on egg production in *Musca domestica* (Diptera: Muscidae) — B. PASTOR, H. ČÍČKOVÁ, M. KOZÁNEK, A. MARTÍNEZ-SÁNCHEZ, P. TAKÁČ & S. ROJO
- 597 Influence of male density on guarding behaviour in the water strider *Aquarius paludum* (Heteroptera: Gerridae) — A. MATSUEDA & K. FUJISAKI
- 603 *Chilocorus renipustulatus* (Coleoptera: Coccinellidae) dominates predatory ladybird assemblages on *Sorbus aucuparia* (Rosales: Rosaceae) — E. KULA & O. NEDVĚD
- 609 Landscape structure affects activity density, body size and fecundity of *Pardosa* wolf spiders (Araneae: Lycosidae) in winter oilseed rape — T. DRAPELA, T. FRANK, X. HEER, D. MOSER & J.G. ZALLER
- 615 Abundance and species richness of overwintering ground beetles (Coleoptera: Carabidae) are higher in the edge than in the centre of a woodlot — A. ROUME, A. OUIN, L. RAISON & M. DECONCHAT
- 623 Butterfly diversity and biogeography on the Croatian karst mountain Biokovo: Vertical distribution and preference for altitude and aspect? — I. MIHOĆI, V. HRŠAK, M. KUĆINIĆ, V. MIĆETIĆ STANKOVIĆ, A. DELIĆ & N. TVRTKOVIĆ
- 635 The parasitoid species complex associated with sexual and parthenogenetic Naryciinae (Lepidoptera: Psychidae): Integrating ecological and molecular analyses — J.A. ELZINGA, K. ZWAKHALS, J. MAPPES & A. GRAPPUTO
- 651 Altitudinal trends in the phenology of butterflies in a mountainous area in central Spain — J.I. DE ARCE CRESPO & D. GUERRÍERREZ
- 659 Biogeography of tenebrionid beetles (Coleoptera: Tenebrionidae) in the circum-Sicilian islands (Italy, Sicily): Multiple biogeographical patterns require multiple explanations — S. FATTORINI
- 673 Morphometry and number of spermatozoa in drone honeybees (Hymenoptera: Apidae) reared under different conditions — A.M. MAZEED
- 677 New stem-Phasmatodea from the Middle Jurassic of China — L. SHANG, O. BÉTHOUX & D. REN
- 687 Discovery of Eutheini (Coleoptera: Staphylinidae: Scydmaeninae) in Australia, with implications for phylogeny and biogeography of *Paraneseuthia* — P. JAŁOSZYŃSKI

NOTES

- 697 Isolation of nine polymorphic microsatellite loci from the leaf beetle, *Lema (Lema) coronata* (Coleoptera: Chrysomelidae: Criocerinae) — Y. MATSUMURA & I. YAO
- 701 Deterrence of feeding in *Rhodnius prolixus* (Hemiptera: Reduviidae) after treatment of antennae with a nitric oxide donor — V. SFARA, E.N. ZERBA & R.A. ALZOGARAY
- 705 Differences in the sugar composition of the honeydew of polyphagous brown soft scale *Coccus hesperidum* (Hemiptera: Sternorrhyncha: Coccoidea) feeding on various host plants — K. GOLAN & A. NAJDA
- 711 Contribution to the biochemical characterization of the silk and structure characterization of the cocoons of the horse chestnut leaf miner *Cameraria ohridella* (Lepidoptera: Gracillariidae) — V. HNĚVŠOVÁ, D. KODRÍK & F. WEYDA