

CONTENTS

ŘÍHA M., HLADÍK M., MRKVÍČKA T., PRCHALOVÁ M., ČECH M., DRAŠTÍK V., FROUZOVÁ J., JŮZA T., KRATOCHVÍL M., PETERKA J., VAŠEK M., KUBEČKA J. Post-spawning dispersal of tributary spawning fish species to a reservoir system	1
CUI L., ABBAS K., YU Y., WANG W., ZHOU L., ZHOU X. First record of the natural occurrence of pentaploid loach, <i>Misgurnus anguillicaudatus</i> in Hubei Province, China	14
ARSLAN A., ZIMA J. The banded karyotype of the 2n = 58 chromosomal race of mole rats from Erzincan, Turkey.....	19
SHCHIPANOV N.A., PAVLOVA S.V. Contact zones and ranges of chromosomal races of the common shrew, <i>Sorex araneus</i> , in north-eastern European Russia	24
NISSEN H., KRÜGER F., FICHTNER A., SOMMER R.S. Local variability in the diet of Daubenton's bat (<i>Myotis daubentonii</i>) in a lake landscape of northern Germany	36
BAO F., WU S., SU CH., YANG L., ZHANG F., MA G. Air temperature changes in a burrow of Chinese pangolin, <i>Manis pentadactyla</i> , in winter.....	42
MALECHA A.W., ANTCZAK M. Diet of the European polecat <i>Mustela putorius</i> in an agricultural area in Poland....	48
BRZEZIŃSKI M., CHIBOWSKA P., ROMANOWSKI J. Dam reservoir affects diet of otters inhabiting mountain river in SE Poland	54
PALMEGIANI I., GAZZOLA A., APOLLONIO M. Wolf diet and its impact on the ungulates community in a new recolonized area of Western Alps: Gran Paradiso National Park.....	59
STACHE A., HELLER E., HOTHORN T., HEURICH M. Activity patterns of European roe deer (<i>Capreolus capreolus</i>) are strongly influenced by individual behaviour	67
XU F., MA M., YANG W., BLANK D., DING P., ZHANG T. Group size effect on vigilance and daytime activity budgets of the <i>Equus kiang</i> (Equidae, Perissodactyla) in Arjinshan National Nature Reserve, Xinjiang, China	76