

Proceedings of the 12th International Conference on Polysaccharides-Glycoscience

CONTENTS:

KEY-NOTE LECTURES

<u>T. Budtova*</u>	Bio-aerogels	6
<u>X. Tan, B. Deng, X. Jiang, P. Zhang, Z. Hu, E. Li, M. A. Sullivan, B. L. Schulz, R. G. Gilbert*</u>	Glycogen: higher-level molecular structure and diabetes	7
<u>W. Praznik, R. Loeppert, H. Viernstein, M. Mueller</u>	Characterization of heteropolysaccharides from seed mucilage of chia (<i>Salvia hispanica</i> L.), basil (<i>Ocimum basilicum</i> L.) and flax (<i>Linum usitatissimum</i> L.) and their healthy functionality	9
<u>M. A. Radeloff</u>	Modifying starches – trends and developments	12

ORAL PRESENTATIONS

<u>E. Achbergerová*, D. Šmejkalová, G. Huerta-Angeles, K. Nešporová, M. Čepa, J. Šógorková, V. Velebný</u>	Fluorescently labelled nanocarriers based on hyaluronic acid: synthesis and <i>in vitro</i> studies	15
<u>S. Balicki, R. Gancarz, I. Pawlaczyk-Graja*</u>	The influence of ultrasounds assisted extraction parameters on the acidic polysaccharides content in the alkaline extracts of <i>Erigeron canadensis</i> L. – a preliminary study	20
<u>R. Bleha*, B. Jindřichová, L. Burketová, A. Synytsya, O. Valentová</u>	Structural analysis of polysaccharides from <i>Leptosphaeria maculans</i> and <i>Alternaria brassicicola</i>	24
<u>D. Čožíková*, M. Hermannová, V. Moravcová, V. Velebný</u>	Preparation of hyaluronan fragments with low polydispersity obtained from acid-catalyzed hydrolysis	29
<u>N. V. Dolgopyatova, Yu. A. Kuchina, V. Yu. Novikov, I. N. Kononova</u>	Enzymes effect on the molecular weight and molecular weight distribution of chondroitin sulfate, derived from marine aquatic organisms	31
<u>S. Gíllarová*, S. Henke, Z. Bubník</u>	Acid and enzymatic hydrolysis of guar gum followed chromatographic separation	35
<u>A. R. Górecki, J. Lewandowicz, K. Penkacik</u>	Effect of high hydrostatic pressure treatment and pullulanase debranching of buckwheat starch on resistant starch formation, rheological properties and microstructural changes	42
<u>M. Chylińska*, B. Prats Mateu, N. Gierlinger, K. Deryło, M. Tchórzewski, M. Szymańska-Chargot, A. Zdunek</u>	Changing of biochemical parameters and cell wall polysaccharides distribution and composition during physiological development of tomato fruit	47
<u>J. Lewandowicz*, H. M. Baranowska, A. Szwengiel, J. Le Thanh-Blicharz</u>	Molecular structure vs. functional properties of waxy and normal corn starch	53

<i>D. Lukin, V. Kolpakova, V. Ananyev, N. Lukin, I. Usachev, A. Sardjveladze, D. Solomin</i>	Application of thermoplastic starch and starch containing waste of food industry in biodegradable polymer compositions	58
<i>A. Matelová*, G. Huerta-Angeles, Z. Brůnová, D. Šmejkalová, R. Vícha, V. Velebný</i>	Synthesis of novel amphiphilic hyaluronan containing $\pi \cdots \pi$ conjugated moieties as carriers for drug delivery	63
<i>N. Sokolan*, N. Voron'ko, S. Derkach, J. Kuchina</i>	Polyelectrolyte complexes of polysaccharides with gelatin	67
<i>A. Szewngiel*, G. Lewandowicz, J. Nowak</i>	Advantages and disadvantages of the use of aqueous and organic GPC columns for the separation of starch macromolecules	71
<i>L. Štěrbová*, J. Bradová, J. Rysová, R. Winterová</i>	Effects of fat type and content on starch digestibility in wheat-based biscuits	77
<i>M. Witek*, I. Maciejaszek, J. Banaś, J. Majcherczyk, K. Surówka</i>	State of water in protein-polysaccharide complexes studied by ^1H NMR relaxation	81

POSTERS

<i>M. Bączkiewicz, T. Fortuna, S. Pietrzyk, D. Galkowska, L. Juszcak, M. Witczak, A. Augustyńska</i>	Alpha-amylase degradation of double modified potato starch	87
<i>J. Rożnowski*, M. Bączkiewicz, I. Przetaczek-Rożnowska, K. Harańczyk</i>	Selected properties of starch phosphate obtained from starches isolated from lentil seeds	91
<i>E. Baeva*, A. Synytsya, J. Čopíková</i>	Microwave superheated water extraction of (1-6)- β -D-glucan from fruiting bodies of <i>Pleurotus ostreatus</i>	96
<i>R. Rezler*, H. M. Baranowska, K. Pers</i>	Rheological and water binding properties of starch gels	100
<i>K. Tremmel-Bede*, L. Láng, Z. Bedő, G. Vida, M. Rakszegi</i>	Identification and development of wheat lines with increased level of arabinoxylan	103
<i>J. Bradová*, L. Štěrbová, T. Sedláček, V. Dvořáček, J. Hermuth</i>	The effect of harvest year and cultivar on starch digestibility in wheat grain and its milling and baking products	108
<i>K. Buksa*, A. Nowotna, W. Praznik</i>	Molecular structure of water extractable and water unextractable arabinoxylan affected by alkali dissolving conditions	112
<i>K. Buksa*, D. Litwinek, M. Kowalczyk, A. Nowotna, J. Boreczek, H. Gambuś</i>	The influence of polysaccharides on sugar, alcohol and organic acid profile in wholemeal bread obtained by spontaneous sourdough fermentation process	115
<i>I. Burešová*, K. Vaculová, M. Zacharová, O. Jirsa, I. Sedláčková</i>	The comparison of different methods used for measuring rheological characteristics of barley suspension and dough during heating and cooling	118
<i>V. Dvořáček, E. Matějová, L. Štěrbová, J. Bradová</i>	Rye translocation – possible breeding tool for nutritional quality improvement of wheat polysaccharides	122
<i>S. Gillarová*, S. Henke, Z. Bubník</i>	Continuous chromatography separation of guar gum hydrolysates	125
<i>D. Gumul*, M. Krystyjan, J. Korus, R. Ziobro, A. Areczuk, J. Sobolewska-Zielińska, H. Gambuś</i>	The changes in content of phenolic acids and flavonoids in starch breads with addition of non-wheat flours during its production	130
<i>J. Korus, D. Gumul*, M. Krystyjan, R. Ziobro, M. Kruczek</i>	Total polyphenol content and antioxidant activity during production stages of starch bread with buckwheat and amaranth flours	133
<i>M. Halaj, P. Capek</i>	The structural features of extracellular biopolymers produced by green microalgae	136
<i>Z. Hromádková*, Z. Košťálová, M. Ondřejovič, D. Chmelová</i>	Isolation, composition and structure of polysaccharides from <i>Larix decidua</i>	142

<u>Z. Hromádková*</u> , <u>Z. Košťálová</u> , <u>T. Yilmaz</u> , <u>S. Tavman</u>	Characterization of pectic polysaccharides extracted from hazelnut skin by hot water	148
<u>M. Hrušková</u> , <u>I. Švec</u>	Flax fiber – influence on bread, cookies and pasta characteristics	153
<u>D. Chena Aldao</u> , <u>E. Šárka</u> , <u>P. Smrčková</u> , <u>L. Gruberová</u>	Use of starch of modified digestibility for pharmaceutical applications	156
<u>D. Myśliwiec</u> , <u>M. Chylińska*</u> , <u>M. Szymańska-Chargot</u> , <u>A. Zdunek</u>	Investigation of adsorption kinetics, equilibrium and isotherm of xyloglucan on cellulose	161
<u>K. Królikowska*</u> , <u>S. Pietrzyk</u> , <u>I. Przetaczek-Rożnowska</u>	Stability and rheological properties of oil-in-water emulsions formed by the octenylsuccinate starches differing in amylose content	166
<u>I. Przetaczek-Rożnowska*</u> , <u>K. Królikowska</u> , <u>B. Lewandowska</u>	Effect of species addition on selected properties of corn starch	170
<u>M. Krystyan*</u> , <u>D. Gumul</u> , <u>R. Sabat</u> , <u>R. Ziobro</u> , <u>J. Korus</u> , <u>J. Sobolewska-Zielińska</u> , <u>A. Areczuk</u>	The content of dietary fiber in gluten-free dough and bread supplemented with amaranth and buckwheat flour and mechanical properties of the dough	175
<u>M. Krystyan*</u> , <u>W. Ciesielski</u> , <u>D. Gumul</u> , <u>K. Buksa</u> , <u>R. Ziobro</u> , <u>J. Korus</u> , <u>M. Sikora</u> , <u>M. Kruczek</u> , <u>J. Sobolewska-Zielińska</u>	The structure and properties of freeze-dried corn starch	178
<u>J. Le Thanh-Blicharz*</u> , <u>A. Szwengiel</u> , <u>J. Lewandowicz</u>	Molecular mass distribution vs. emulsifying activity of maltodextrins	181
<u>Z. Małysek</u> , <u>J. Le Thanh-Blicharz</u> , <u>A. Drożdżyńska</u> , <u>G. Lewandowicz</u>	The effect of degree of substitution on rheological properties of acetylated distarch adipate	185
<u>J. Le Thanh-Blicharz</u> , <u>A. Szwengiel</u> , <u>G. Lewandowicz*</u>	Molecular structure of starch modified by high pressure homogenisation of pastes	189
<u>Ł. Masewicz</u> , <u>J. Lewandowicz</u> , <u>J. Le Thanh-Blicharz</u> , <u>M. Kempka</u> , <u>H. M. Baranowska*</u>	Diffusion of water in potato starch pastes	193
<u>D. Longmire</u> , <u>H. Hudelja</u> , <u>R. Bleha</u> , <u>J. Čopíková</u>	Polysaccharides in cocoa beans	196
<u>M. Marounek</u> , <u>Z. Volek</u> , <u>T. Taubner</u> , <u>D. Dušková</u>	Effect of amidated alginate on cholesterol in serum, liver and faeces of rats fed fat and cholesterol-containing diet	201
<u>O. Mikala</u> , <u>M. Milichovský</u> , <u>J. Gojný</u> , <u>M. Hájek</u>	Influence of vapours of essential oils on the porous lignocellulose materials during paper aging	204
<u>D. Murdzheva</u> , <u>I. Vasileva</u> , <u>N. Petkova</u> , <u>I. Ivanov</u> , <u>M. Todorova</u> , <u>P. Denev*</u>	New approach for synthesis of amidated alginic acids	208
<u>D. Murdzheva</u> , <u>N. Petkova</u> , <u>I. Vasileva</u> , <u>M. Todorova</u> , <u>M. Ognyanov</u> , <u>I. Ivanov</u> , <u>P. Denev*</u>	Accelerated modification of low-methoxylated celery pectin	213
<u>B. Pančíková*</u> , <u>M. Sluková</u> , <u>P. Skřivan</u>	The comparison of different milling processes on structure and properties of flour	218
<u>S. Pepeliaev*</u> , <u>J. Jílková</u> , <u>Z. Černý</u> , <u>D. Smirnou</u> , <u>L. Franke</u> , <u>V. Velebný</u>	New colorimetric assay for hyaluronidase activity determination	220
<u>S. Pietrzyk*</u> , <u>K. Królikowska</u> , <u>J. Šek</u> , <u>U. Lech</u> , <u>A. Kosiorowska</u>	The effect of oxidation process parameters on functional properties of modified starch	223
<u>J. Pinkrová*</u> , <u>I. Boháčenko</u> , <u>Z. Šmídová</u>	Modified HPLC method with RI detection for determination of cereal β -(1-3),(1-4)-D-glucan	227
<u>J. Pravda</u> , <u>J. Čopíková</u> , <u>P. Matricardi</u>	Preparation of nanohydrogels for <i>Varroa</i> control	230
<u>M. Pravda*</u> , <u>R. Buffa</u> , <u>J. Bystroňová</u> , <u>D. Čožíková</u> , <u>M. Hermannová</u>	Hydrogels based on hyaluronan-tyramine conjugate – influence of conjugate structure on hydrogel properties	234
<u>P. Paják*</u> , <u>S. Czyjt-Kurylo</u> , <u>I. Przetaczek-Rożnowska</u> , <u>T. Fortuna</u>	The influence of guar and cassia gums on the physical properties of starch-based films	239
<u>I. Przetaczek-Rożnowska*</u>	Some physicochemical properties of starches isolated from <i>Cucurbita</i> varieties	243

<u>P. Smrčková</u> , R. Filepová, E. Šárka, A. D. Antonio Chena	Characterisation of starch in food materials by GPC	246
<u>J. Sobolewska-Zielińska*</u> , D. Gumul, M. Krystyjan, D. Galkowska	Physical properties of low-fat mayonnaise with maltodextrin and modified starch	250
D. Galkowska*, J. Cieśla, <u>J. Sobolewska-Zielińska</u>	Non-enzymatic browning of model sponge cakes	253
<u>A. Szewngiel*</u> , M. Bednarek	Optimisation of dissolution conditions of potato starch for aqueous SEC	259
E. Šárka, V. Jankovec, P. Smrčková, D. A. Chena Aldao, A. Synytsya	Chemical reactions of wheat starch characterized by low volume of the reaction liquid	265
<u>I. Švec</u> , M. Hrušková	Flax fiber – influence on composite flour behaviour	270
<u>Y. Troshchynska</u> , A. Synytsya, J. Kyselka, J. Štetina	Flaxseed meal mucilage polysaccharides: chemical characterization and purification	273
<u>T. Uhlířová*</u> , E. Gregorová, W. Pabst	Bread cutter tomography – from stereology to topology	278
<u>K. Vaculová*</u> , I. Sedláčková, O. Jirsa, E. Bajerová	The influence of controlled germination on the content of selected nutrients in grain and milling fractions of barley for prospective human consumption	283
<u>I. Wiede*</u> , B. Wiede, M. Sluková, L. Jurkaninová, K. Vaculová	Characterization of β -glucan extracted from native and extruded barley flour	288
M. Witek*, I. Maciejaszek, J. Majcherczyk, J. Banaś, <u>K. Surówka</u>	The impact of hydration on water mobility in protein- polysaccharide electrocomplexes	292