

EXPERIMENTAL CELL RESEARCH

Volume 315, Number 3, February 1, 2009

CONTENTS

| | |
|--|-----|
| Highlights | v |
| <i>Research Articles</i> | |
| Elske J. Schabort, Mathilde van der Merwe, Benjamin Loos, Frances P. Moore, and Carola U. Niesler. TGF- β 's delay skeletal muscle progenitor cell differentiation in an isoform-independent manner | 373 |
| Raffaella Solito, Federico Corti, Silvia Fossati, Emiliya Mezhericher, Sandra Donnini, Jorge Ghiso, Antonio Giachetti, Agueda Rostagno, and Marina Ziche. Dutch and arctic mutant peptides of β amyloid ₁₋₄₀ differentially affect the FGF-2 pathway in brain endothelium | 385 |
| Po-Feng Lee, Alvin T. Yeh, and Kayla J. Bayless. Nonlinear optical microscopy reveals invading endothelial cells anisotropically alter three-dimensional collagen matrices | 396 |
| Michelle N. Yarmo, Anne Landry, André S.D. Molgat, AnneMarie Gagnon, and Alexander Sorisky. Macrophage-conditioned medium inhibits differentiation-induced Rb phosphorylation in 3T3-L1 preadipocytes | 411 |
| Nicolás Pregi, Shirley Wenker, Daniela Vittori, Claudia Pérez Leirós, and Alcira Nesse. TNF-alpha-induced apoptosis is prevented by erythropoietin treatment on SH-SY5Y cells | 419 |
| Jasleen Shant, Kunrong Cheng, Bernard S. Marasa, Jian-Ying Wang, and Jean-Pierre Raufman. Akt-dependent NF- κ B activation is required for bile acids to rescue colon cancer cells from stress-induced apoptosis | 432 |
| Elisabeth Hedström, Natalia Issaeva, Martin Enge, and Galina Selivanova. Tumor-specific induction of apoptosis by a p53-reactivating compound | 451 |
| Piotr Rieske, Brian J. Augelli, Robert Stawski, John Gaughan, S. Ausim Azizi, and Barbara Krynska. A population of human brain cells expressing phenotypic markers of more than one lineage can be induced <i>in vitro</i> to differentiate into mesenchymal cells | 462 |
| Hisashi Ideno, Rieko Takanabe, Akemi Shimada, Kazuhiko Imaizumi, Ryoko Araki, Masumi Abe, and Akira Nifuji. Protein related to DAN and cerberus (PRDC) inhibits osteoblastic differentiation and its suppression promotes osteogenesis <i>in vitro</i> | 474 |

Available online at



www.sciencedirect.com



0014-4827 (20090201)315:3;1-#

| | |
|--|-----|
| Philipp Baumann, Sonja Mandl-Weber, Fuat Oduncu, and Ralf Schmidmaier. The novel orally bioavailable inhibitor of phosphoinositol-3-kinase and mammalian target of rapamycin, NVP-BEZ235, inhibits growth and proliferation in multiple myeloma | 485 |
| Anja Apel, Ariane Groth, Sabine Schlesinger, Helge Bruns, Peter Schemmer, Markus W. Büchler, and Ingrid Herr. Suitability of human mesenchymal stem cells for gene therapy depends on the expansion medium..... | 498 |
| Claire Robinson and Andreas F. Kolb. Analysis of mammary specific gene locus regulation in differentiated cells derived by somatic cell fusion | 508 |
| Michelle L. Milstein, Timothy D. Houle, and Steven E. Cala. Calsequestrin isoforms localize to different ER subcompartments: Evidence for polymer and heteropolymer-dependent localization..... | 523 |
| M.L. Fanarraga, J.C. Villegas, G. Carranza, R. Castaño, and J.C. Zabala. Tubulin cofactor B regulates microtubule densities during microglia transition to the reactive states | 535 |
| Elena Kolobova, Andrey Efimov, Irina Kaverina, Arun K. Rishi, John W. Schrader, Amy-Joan Ham, M. Cecilia Larocca, and James R. Goldenring. Microtubule-dependent association of AKAP350A and CCAR1 with RNA stress granules | 542 |



Covered in the abstract and citation database BIOBASE, Biological & Agricultural Index, Biological Abstracts, Biosis Previews, Chemical Abstracts, Current Awareness in Biological Sciences, Current Contents, EMBASE, EMBiology, Genetics Abstracts, MEDLINE®, Science Citation Index, and SCOPUS®. Full text available on ScienceDirect®.