


Contents in Brief

List of Topics	ix
Acknowledgments	xxix
Prologue	xxxiii
A Note to the Reader	xxxix

Donated by the Universities of
UMEÅ and UPPSALA through
BAHA'I-STUDENT-SOCIETIES
in Sweden

Introduction to the Cell


PART



1. Evolution of the Cell	3
2. Small Molecules, Energy, and Biosynthesis	43
3. Macromolecules: Structure, Shape, and Information	91
4. How Cells are Studied	143

The Molecular Organization of Cells


PART



5. Basic Genetic Mechanisms	199
6. The Plasma Membrane	255
7. Internal Membranes and the Synthesis of Macromolecules	319
8. The Cell Nucleus	385
9. Energy Conversion: Mitochondria and Chloroplasts	483
10. The Cytoskeleton	549
11. Cell Growth and Division	611
12. Cell-Cell Adhesion and the Extracellular Matrix	673
13. Chemical Signaling Between Cells	717

From Cells to Multicellular Organisms

PART



14. Germ Cells and Fertilization	769
15. Cellular Mechanisms of Development	813
16. Differentiated Cells and the Maintenance of Tissues	891
17. The Immune System	951
18. The Nervous System	1013
19. Special Features of Plant Cells	1099

Index

List of Topics

PART

I

Introduction to the Cell

CHAPTER

1

The Evolution of the Cell

From Molecules to the First Cell

Simple Biological Molecules Can Form Under Prebiotic Conditions	4
Polynucleotides Are Capable of Directing Their Own Synthesis	4
Self-replicating Molecules Undergo Natural Selection	6
Information Flows from Polynucleotides to Polypeptides	7
Membranes Defined the First Cell	8
Mycoplasmas Are the Simplest Living Cells	10
Summary	11

From Procarvates to Eucaryotes

Procarvotic Cells Are Structurally Simple But Biochemically Diverse	11
Metabolic Reactions Evolve	12
Cyanobacteria Can Fix CO ₂ and N ₂	14
Bacteria Can Carry Out the Aerobic Oxidation of Food Molecules	15
Eucaryotic Cells Contain Several Distinctive Organelles	15
Eucaryotic Cells Depend on Mitochondria for Their Oxidative Metabolism	19
Chloroplasts May Be Descendants of Procarvotic Algae	19
Eucaryotic Cells Contain a Rich Array of Internal Membranes	21
Eucaryotic Cells Have a Cytoskeleton	21
Protozoa Include the Most Complex Cells Known	23
Genes Can Be Switched On and Off	23
Eucaryotic Cells Have Vastly More DNA Than They Need for the Specification of Proteins	26
In Eucaryotic Cells the Genetic Material Is Packaged in Complex Ways	26
Summary	27

From Single Cells

to Multicellular Organisms

Single Cells Can Associate to Form Colonies	28
The Cells of a Higher Organism Become Specialized and Cooperate	29
Multicellular Organization Depends on Cohesion Between Cells	29
Epithelial Sheets of Cells Enclose a Sheltered Internal Environment	30
Cell-Cell Communication Controls the Spatial Pattern of Multicellular Organisms	31
Cell Memory Permits the Development of Complex Patterns	31
Basic Developmental Programs Tend to Be Conserved in Evolution	32
Eucaryotic Organisms Possess a Complex Machinery for Reproduction	32
The Cells of the Vertebrate Body Exhibit More Than 200 Different Modes of Specialization	33
Cells of the Immune System Are Specialized for the Task of Chemical Recognition	34
Nerve Cells Allow a Rapid Adaptation to a Changing World	35
Developing Nerve Cells Must Assemble to Form a Nervous System	36
Nerve Cell Connections Determine Patterns of Behavior	37
Summary	39
References	41

CHAPTER

2

Small Molecules, Energy, and Biosynthesis

The Chemical Components of a Cell

Cell Chemistry Is Based on Carbon Compounds	44
Cells Use Four Basic Types of Small Molecules	45

X List of Topics

Sugars Are Food Molecules of the Cell
Fatty Acids Are Components of Cell Membranes
Amino Acids Are the Subunits of Proteins
Nucleotides Are the Subunits of DNA and RNA
Summary

Biological Order and Energy

Biological Order Is Made Possible by the Release of Heat Energy from Cells
Photosynthetic Organisms Use Sunlight to Synthesize Organic Compounds
Chemical Energy Passes from Plants to Animals
Cells Obtain Energy by the Oxidation of Biological Molecules
The Breakdown of Organic Molecules Takes Place in Sequences of Enzyme-catalyzed Reactions
Part of the Energy Released in Oxidation Reactions Is Coupled to the Formation of ATP
The Hydrolysis of ATP Generates Order in Cells
Summary

Food and the Derivation of Cellular Energy

Food Molecules Are Broken Down in Three Stages to Give ATP
Glycolysis Can Produce ATP Even in the Absence of Oxygen
Oxidative Catabolism Yields a Much Greater Amount of Usable Energy
Metabolism Is Dominated by the Citric Acid Cycle
The Transfer of Electrons to Oxygen Drives ATP Formation
Amino Acids and Nucleotides Are Part of the Nitrogen Cycle
Summary

Biosynthesis and the Creation of Order

The Energy for Biosynthesis Comes from the Hydrolysis of ATP
Biosynthetic Reactions Are Often Directly Coupled to ATP Hydrolysis
Coenzymes Are Involved in the Transfer of Specific Chemical Groups
Biosynthesis Requires Reducing Power
Biological Polymers Are Synthesized by Repetition of Elementary Dehydration Reactions
Summary

The Coordination of Catabolism and Biosynthesis

Metabolism Is Organized and Regulated
Metabolic Pathways Are Regulated by Changes in Enzyme Activity
Catabolic Reactions Can Be Reversed by an Input of Energy
Enzymes Can Be Switched On and Off by Covalent Modification

50	Reactions Are Compartmentalized Both Within Cells and Within Organisms	87
51		
56	<i>Summary</i>	89
56	References	89
62		

CHAPTER 3

Macromolecules: Structure, Shape, and Information

63	Molecular Recognition Processes	91
64	The Information Carried by a Macromolecule Is Expressed by Means of Weak Noncovalent Bonds	92
64	Diffusion Is the First Step to Molecular Recognition	93
65	Thermal Motions Not Only Bring Molecules Together But Also Pull Them Apart	96
66	Molecular Recognition Processes Can Never Be Perfect	97
67	<i>Summary</i>	98
67	Nucleic Acids	98
	Genes Are Made of DNA	98
67	DNA Molecules Consist of Two Long Complementary Chains Held Together by Base Pairs	99
67	The Structure of DNA Provides an Explanation for Heredity	103
69	Errors in DNA Replication Cause Mutations	103
71	The Nucleotide Sequence of a Gene Determines the Amino Acid Sequence of a Protein	106
72	Portions of the DNA Sequence Are Copied into RNA	107
73	Sequences of Nucleotides in mRNA Are "Read" in Sets of Three	107
74	tRNA Molecules Match Amino Acids to Groups of Nucleotides	108
75	The RNA Message Is Read from One End to the Other by a Ribosome	109
	<i>Summary</i>	111
	Protein Structure	111
75	The Shape of a Protein Molecule Is Determined by Its Amino Acid Sequence	111
77	Common Folding Patterns Recur in Different Protein Chains	113
79	Proteins Are Enormously Versatile in Structure	115
80	Proteins Show Different Levels of Structural Organization	118
81	Relatively Few of the Many Possible Polypeptide Chains Would Be Useful	118
82	New Proteins Often Evolve by Minor Alterations of Old Ones	118
82	New Proteins Often Evolve Through the Combination of Different Polypeptide Domains	120
84	Protein Subunits Can Self-assemble into Large Structures in the Cell	121
84	A Single Type of Protein Subunit Can Interact with Itself to Form Geometrically Regular Aggregates	121
87	Self-assembling Aggregates Can Include Different Protein Subunits and Nucleic Acids	124

There Are Limits to Self-assembly	126	Image Reconstruction Techniques Based on	
<i>Summary</i>	127	Diffraction Can Be Used to Extract Additional	
Protein Function	127	Information from Electron Micrographs	157
A Protein's Conformation Determines Its Chemistry	128	X-ray Diffraction Reveals the Three-dimensional	
Substrate Binding Is the First Step in Enzyme		Arrangement of the Atoms in a Molecule	158
Catalysis	129	<i>Summary</i>	159
Enzymes Accelerate Reaction Rates Without Shifting		Cell Culture	160
Equilibria	131	Cells Can Be Grown in a Culture Dish	161
Many Enzymes Make Reactions Proceed Preferentially		Chemically Defined Media Permit Identification of	
in One Direction by Coupling Them to ATP		Specific Growth Factors	162
Hydrolysis	132	Eucaryotic Cell Lines Are a Convenient Source of	
Multienzyme Complexes Help to Increase the Rate of		Homogeneous Cells	162
Cell Metabolism	132	Cells Can Be Fused Together to Form Hybrid Cells	163
Intracellular Membranes Increase the Rates of		<i>Summary</i>	165
Diffusion-limited Reactions	133	The Fractionation of Cells	
Protein Molecules Can Reversibly Change Their Shape	134	and Their Contents	165
Allosteric Proteins Are Involved in Metabolic		Cells Can Be Isolated from a Tissue and Separated	
Regulation	134	into Different Types	166
Allosteric Proteins Are Vital for Cell Signaling		Organelles and Macromolecules Can Be Separated by	
Proteins Can Be Pushed or Pulled into Different		Ultracentrifugation	166
Shapes	135	The Molecular Details of Complex Cellular Processes	
Energy-driven Changes in Protein Conformations Can		Can Be Determined Only by Studies of Fractionated	
Do Useful Work	137	Cell Components	169
ATP-driven Membrane-bound Allosteric Proteins Can		Proteins Can Be Separated by Chromatography	170
Act as Pumps	137	The Size and Subunit Composition of a Protein Can	
Protein Molecules Can Harness Ion Gradients to Do		Be Determined by SDS Polyacrylamide-Gel	
Useful Work	139	Electrophoresis	173
<i>Summary</i>	139	More Than 1000 Proteins Can Be Resolved on a Single	
References	139	Gel by Two-dimensional Polyacrylamide-Gel	
		Electrophoresis	175
		Selective Cleavage of a Protein Generates a Distinctive	
		Set of Peptide Fragments	177
		Short Amino Acid Sequences Can Be Analyzed by	
		Automated Machines	177
		<i>Summary</i>	178
		Tracing Cellular Molecules with	
		Radioactive Isotopes and Antibodies	179
		Radioactive Atoms Can Be Detected with Great	
		Sensitivity	179
		Radioisotopes Are Used to Trace Molecules in Cells	
		and Organisms	179
		Antibodies Can Be Used to Detect and Isolate Specific	
		Molecules	181
		Hybridoma Cell Lines Provide a Permanent Source of	
		Monoclonal Antibodies	182
		Antibodies and Other Macromolecules Can Be	
		Injected into Living Cells	184
		<i>Summary</i>	184
		Recombinant DNA Technology	185
		Restriction Nucleases Hydrolyze DNA Molecules at	
		Specific Nucleotide Sequences	185
		Selected DNA Sequences Are Produced in Large	
		Amounts by Cloning	187
		Copies of Specific mRNA Molecules Can Be Cloned	188

CHAPTER

4

How Cells Are Studied**Microscopy**

The Light Microscope Can Resolve Details 0.2 μm	143
Apart	144
Different Components of the Cell Can Be Selectively	
Stained	145
Tissues Are Usually Fixed and Stained for Microscopy	145
Living Cells Can Be Seen in a Phase-Contrast or	
Differential-Interference-Contrast Microscope	146
The Electron Microscope Resolves the Fine Structure	
of the Cell	148
Biological Specimens Require Special Preparation for	
the Electron Microscope	149
Three-dimensional Images Can Be Obtained by	
Electron Microscopy	151
Freeze-Fracture and Freeze-Etch Electron Microscopy	
Provide Novel Views of the Cell	152
Individual Macromolecules Can Be Resolved in the	
Electron Microscope	154
The Detailed Structure of Molecules in a Crystalline	
Array Can Be Calculated from the Diffraction Patterns	154
They Create	

Cloned DNA Fragments Can Be Rapidly Sequenced	189	Recombinant DNA Techniques Allow Even the Minor Proteins of a Cell to Be Studied	191
Nucleic Acid Hybridization Reactions Provide a Sensitive Way of Detecting Specific Nucleotide Sequences	189	Mutant Genes Can Now Be Made to Order	192
<i>In Situ</i> Hybridization Techniques Are Used to Localize Specific Nucleic Acid Sequences in Chromosomes and Cells	191	Summary	194
		References	194

The Molecular Organization of Cells

PART



Basic Genetic Mechanisms

Protein Synthesis

The Decoding of DNA into Protein	199
RNA Polymerase Copies DNA into RNA	200
Protein Synthesis Is Inherently Very Complex	202
Specific Enzymes Couple Each Amino Acid to Its Appropriate tRNA Molecule	205
Amino Acids Are Added to the Carboxyl-Terminal End of a Growing Polypeptide Chain	206
Each Amino Acid Added Is Selected by a Complementary Base-pairing Interaction Between Its Linked tRNA Molecule and an mRNA Chain	207
The Events in Protein Synthesis Are Catalyzed on the Ribosome	208
A Ribosome Moves Stepwise Along the mRNA Chain	210
A Protein Chain Is Released from the Ribosome Whenever One of Three Different Termination Codons Is Reached	211
The Initiation Process Sets the Reading Frame for Protein Synthesis	211
The Overall Rate of Protein Synthesis in Eucaryotes Is Controlled by Initiation Factors	212
Many Inhibitors of Procarvotic Protein Synthesis Are Useful as Antibiotics	212
Summary	213

DNA Repair Mechanisms

DNA Sequences Are Maintained with a Very High Fidelity	214
Directly Detected Spontaneous-Mutation Rates Are Consistent with Evolutionary Estimates	214
Most Mutations in Proteins Are Deleterious and Are Eliminated by Natural Selection	215

CHAPTER

5

Low Mutation Rates Mean That Related Organisms Must Be Made from Essentially the Same Proteins	216
If Left Uncorrected, Spontaneous DNA Damage Would Change DNA Sequences Rapidly	216
The Stability of Genes Is Due to DNA Repair	217
The Structure and Chemistry of the DNA Double Helix Make It Easy to Repair	220
Summary	221
DNA Replication Mechanisms	221
Base-pairing Underlies DNA Replication as well as DNA Repair	221
The DNA Replication Fork Is Asymmetrical	223
The High Fidelity of DNA Replication Requires a "Proofreading" Mechanism	224
DNA Replication in the 5'-to-3' Direction Is Intrinsically More Accurate	226
A Special Nucleotide Polymerizing Enzyme Is Needed to Synthesize Short Primer Molecules on the Lagging Strand	226
Special Proteins Help Open Up the DNA Double Helix in Front of the Replication Fork	227
Other Special Proteins Prevent DNA Tangling	228
The Replication of DNA in Eucaryotes and Procarvotes Is Basically Similar	230
Replication Forks Are Created at Replication Origins	231
Summary	232
Viruses	232
Viruses Are Mobile Genes	232
The Genetic Component of a Virus Is Either DNA or RNA	233
The Outer Coat of a Virus May Be a Protein Capsid or a Membrane Envelope	233
Viral Chromosomes Usually Code for One or More Enzymes Involved in the Replication of the Viral Nucleic Acid	235
Viral Genomes Come in a Variety of Forms	235

Viral Chromosomes Can Integrate Themselves into Host Chromosomes	236	Membrane Transport Proteins Can Be Visualized by Freeze-Fracture Electron Microscopy	272
RNA Viruses Also Replicate Through the Formation of Complementary Chains	236	Vectorial Labeling Reagents Can Be Used to Study Some Plasma Membrane Proteins of Nucleated Cells	274
Viral Genetic Elements Can Make Cells Cancerous	238	When Two Cells Are Fused Together, Their Plasma Membrane Proteins Rapidly Mix	275
How Did Viruses Evolve?	239	Membrane Proteins Cluster into Patches When They Are Cross-linked by Antibodies	276
Summary	240	Cross-linked Membrane Proteins Are Actively Swept to One Pole of the Cell in the Process of "Capping"	278
Genetic Recombination Mechanisms	240	Conflicting Views on How Cells Cap: Flow Versus Pull Hypotheses	279
General Recombination Processes Are Guided by Base-pairing Interactions Between Complementary Strands of Homologous DNA Helices	241	Antibody-induced Redistribution Can Be Used to Determine Whether Two Polypeptides Are Associated with Each Other in the Plasma Membrane	280
General Recombination Is Initiated at a Nick in One Strand of a DNA Double Helix	242	Lateral Diffusion Rates of Membrane Proteins Can Be Quantified	281
Special Proteins Enable DNA Single Strands to Pair with a Homologous Region of DNA Double Helix	243	Cells Have Ways of Restricting the Lateral Mobility of Certain of Their Membrane Proteins	282
Genetic Recombination Usually Involves a Cross-Strand Exchange	244	Summary	283
General Recombination Aids DNA Repair Processes	247	Membrane Carbohydrate	283
Site-specific Recombination Enzymes Move Special Sequences of DNA in and out of Genomes	247	The Carbohydrate in Biological Membranes Is Confined to the Noncytoplasmic Surface	284
Summary	250	Cell-Surface Carbohydrate Is Suspected to Be Important in Cell-Cell Interactions, But This Has Been Difficult to Demonstrate	285
References	250	Summary	286
<hr/>			
CHAPTER		6	
The Plasma Membrane			
The Lipid Bilayer	256	Membrane Transport of Small Molecules	286
Membrane Lipids Are Amphipathic Molecules That Spontaneously Form Bilayers	256	Protein-free Lipid Bilayers Are Impermeable to Ions But Freely Permeable to Water	287
The Lipid Bilayer Is a Two-dimensional Fluid	258	Membrane-bound Transport Proteins Transfer Specific Small Molecules Across Cell Membranes	287
The Fluidity of a Lipid Bilayer Depends on Its Composition	259	Transport Proteins Form a Continuous Protein Pathway Across the Lipid Bilayer	289
The Lipid Bilayer Serves as a Solvent for Membrane Proteins	260	Carrier Proteins Behave Like Membrane-bound Enzymes	289
The Lipid Bilayer of the Plasma Membrane Is Asymmetrical	261	The Membrane Potential That Exists Across the Plasma Membrane Is Maintained by a Na ⁺ -K ⁺ Pump	290
Glycolipids Are Found on the Surface of All Plasma Membranes, But Their Function Is Unknown	262	The Ubiquitous Plasma Membrane Na ⁺ -K ⁺ Pump Is an ATPase	291
Summary	263	The Na ⁺ -K ⁺ ATPase Helps Control Cell Volume by Controlling the Solute Concentration Inside Cells	294
Membrane Proteins	264	Some Ca ²⁺ Pumps Are Also Membrane-bound ATPases	295
Many Membrane Proteins Are Held in the Bilayer by Hydrophobic Interactions with Lipid Molecules	264	Membrane-bound Enzymes That Synthesize ATP Are Transport ATPases Working in Reverse	295
The Use of SDS Polyacrylamide-Gel Electrophoresis Has Revolutionized the Study of Membrane Proteins	266	Active Transport Can Be Driven by Ion Gradients	295
The Cytoplasmic Side of Membrane Proteins Can Be Studied in Red Blood Cell Ghosts	266	Active Transport in Bacteria Can Occur by "Group Translocation" and Can Involve Water-soluble Binding Proteins	297
Spectrin Is Loosely Associated with the Cytoplasmic Side of the Red Blood Cell Membrane	268	Some Transmembrane Protein Channels Are "Gated" and Open Only Transiently	298
Glycophorin Extends Through the Red Cell Lipid Bilayer as a Single α-Helix	268	Asymmetrically Distributed Ion Channels Can Generate Ion Currents That Polarize Cells	300
Band III of the Human Red Blood Cell Membrane Is a Transport Protein	269	Ionophores Increase the Ion Permeability of Synthetic and Biological Membranes	301
Bacteriorhodopsin Is a Proton Pump That Traverses the Bilayer as Seven α-Helices	271	Summary	301

Membrane Transport of Macromolecules and Particles: Exocytosis and Endocytosis

Exocytosis Occurs by the Fusion of Intracellular Vesicles with the Plasma Membrane	303
Triggered Exocytosis Is a Localized Response of the Plasma Membrane and Its Underlying Cytoplasm	304
Membrane Fusion Involves Bilayer Adherence Followed by Bilayer Joining	305
Endocytosis Occurs Continually in Most Cells	306
Most Endocytotic Vesicles Ultimately Fuse with Lysosomes	307
Many Endocytotic Vesicles Are Coated	307
Coated Pits and Vesicles Provide a Specialized Pathway for Receptor-mediated Endocytosis of Specific Macromolecules	309
Many Cell-Surface Receptors Associate with Coated Pits Only After Ligand Binding	311
Some Macromolecules Can Penetrate Cell Membranes Directly	311
Specialized Phagocytic Cells Ingest Particles That Bind to Specific Receptors on Their Surface	311
Phagocytosis Is a Localized Response That Proceeds by a "Membrane-zipping" Mechanism	312
Membrane Vesicular Traffic: How Is It Powered, Guided, and Regulated?	313
<i>Summary</i>	314

References

The Binding of Many Ribosomes to an Individual Messenger RNA Molecule Generates Polysomes	328
Protein Synthesis Is Blocked by Specific Inhibitors	330
Some Proteins Regulate the Rate of Their Own Synthesis by Binding to the Messenger RNA Molecules on Which They Are Made	330
In Eucaryotes Only One Species of Polypeptide Chain Can Be Synthesized on Each Messenger RNA Molecule	331
Polyproteins Are Often Made in Eucaryotes	332
Many Proteins Undergo Covalent Modification After Their Synthesis	333
Some Proteins Are Degraded Soon After They Have Been Synthesized	334
Not All Proteins Synthesized in the Cytosol Remain There	335
<i>Summary</i>	335

The Endoplasmic Reticulum 335

Attached Ribosomes Define "Rough" Regions of ER	336
Smooth ER Is Abundant in Certain Specialized Cells	338
Rough and Smooth Regions of ER Can Be Physically Separated	339
Rough Regions of ER Contain Specific Proteins Responsible for the Binding of Ribosomes	340
Membrane-bound Ribosomes Synthesize Proteins That Pass Through the Membrane During Their Translation	341
Direct Evidence Favoring Vectorial Discharge Has Come from Experiments in Bacteria	342

Membrane-bound Ribosomes Are Derived from Free Ribosomes That Are Directed to the ER Membrane by Special Signal Sequences	343
There Is Genetic Evidence for the Signal Hypothesis	344
Some Proteins Cross Membranes by a Posttranslational Import Mechanism Rather Than by Vectorial Discharge	344
Most Proteins Synthesized in the Rough ER Are Glycosylated	345

The Oligosaccharide Is Added to the Growing Polypeptide Chain on the Luminal Side of the ER	347
The Oligosaccharide Is Donated to the Polypeptide by an Activated Lipid and Then Almost Immediately Modified	347
Lipids Are Synthesized in the ER Membrane	350
Lipid Biosynthesis Is Asymmetric	351
Special Proteins Transfer Phospholipids from the ER to Mitochondria	351
The Luminal Side of an Internal Organelle Is Topologically Equivalent to the Outside of the Cell	352
Membrane Growth by Continuous Expansion of the ER Ensures the Propagation of Transmembrane Asymmetry	353
<i>Summary</i>	354

The Golgi Apparatus 355

The Golgi Apparatus Consists of Stacks of Disc-shaped Cisternae with Associated Small Vesicles	355
--	-----

CHAPTER

7

Internal Membranes and the Synthesis of Macromolecules

The Compartmentalization of Higher Cells

Large Eucaryotic Cells Need Internal Membranes	320
Internal Membranes Divide the Cell into Specialized Compartments	320
Even Complex Eucaryotic Cells Have Only a Few Major Intracellular Compartments	320
Intracellular Compartments Permit the Cell to Carry Out Many Incompatible Chemical Reactions Simultaneously	323
Viruses Reveal the Existence of Highly Organized Pathways Between Host Cell Compartments	323
Different Viruses Follow Different Pathways Through the Cell	325
<i>Summary</i>	326

The Cytosol

Most Intermediary Metabolism Takes Place in the Cytosol	326
Many Proteins Are Synthesized by Ribosomes in the Cytosol	328

The Golgi Apparatus Is Structurally and Biochemically Polarized

The Golgi Apparatus Is Not Yet Understood in Biochemical Terms

Carbohydrate Structures Are Modified in the Golgi Apparatus

Oligosaccharide-trimming Pathways Are Elaborate and Precisely Programmed

The Correct Program of Trimming Is Set by an Irreversible "Switch" Thrown Early in Oligosaccharide Processing

Carbohydrate Modification in the Golgi Apparatus Can Be Detected by Autoradiography

Proteins Destined for Secretion Are Packaged into Golgi-associated Secretory Vesicles That Then Fuse with the Plasma Membrane

Membrane Components Are Recycled

Membrane Fusion and Recycling Can Be Demonstrated by Electron Microscopy

Coated Vesicles Are Thought to Play a Major Part in the Intracellular Sorting of Proteins

Why Have a Golgi Apparatus?

Summary

Lysosomes and Peroxisomes

Lysosomes Are the Principle Sites of Intracellular Digestion

Histochemical Staining Demonstrates That Lysosomes Are Heterogeneous Organelles

Primary Lysosomes Are Formed by Budding from the Golgi Apparatus

Lysosomal Function in Cultured Cells from Patients Having a Lysosomal Enzyme Deficiency Can Be Corrected by Adding the Missing Enzyme to the Culture Medium

Only Lysosomal Hydrolases with a Mannose-Phosphate-containing Oligosaccharide Are Internalized

Intracellular Hydrolases Are Probably Directed to Primary Lysosomes by the Mannose-Phosphate Marker

The Membrane of the Peroxisome Is Formed by Budding from the Smooth ER

Peroxisomes Use Oxygen to Carry Out Catabolic Reactions

The Enzyme Content of Peroxisomes Varies with Cell Type

A Sorting Mechanism Similar to That Used in the Lysosomal Pathway May Operate for Other Intracellular Compartments

Summary

Organelles with Double Membranes: The Nucleus, Mitochondria, and Chloroplasts

Bacteria with Inner and Outer Cell Membranes May Be Viewed as Models of the Double-Membrane Organelles

Membrane Proteins May Move Between the Inner and Outer Bilayers at Adhesion Sites

Adhesion Sites May Also Play a Part in Transporting Water-soluble Molecules into the Lumen of Double-Membrane Organelles

Summary

References

CHAPTER

8

The Cell Nucleus

The Organization of DNA into Chromosomes

Histones Are Among the Most Highly Conserved Proteins Known 386

The Association of Histones with DNA Leads to the Formation of Nucleosomes, the Unit Particles of Chromatin 388

Nucleosomes Are Packed Together to Form Regular Higher-Order Structures 388

Histone H1 Proteins Help Pack Nucleosomes Together 389

Not All Nucleosomes Are Packed in Precisely the Same Way 390

367 The DNA Helix Is Punctuated by Proteins That Bind to Specific DNA Sequences 392

Proteins Can Recognize Specific DNA Sequences by Hydrogen Bonding to Base Pairs and by Sensing Helix Geometry 394

Histones Restrict the Accessibility of DNA to Other DNA-binding Proteins and Can Thereby Affect Gene Regulation 395

Nucleosome Beads Can Be Nonrandomly Positioned in Chromatin 395

Each Chromosome Probably Contains One Very Long DNA Molecule Organized in a Series of Looped Domains 397

Bands on Mitotic Chromosomes Reveal an Even Higher Level of Organization 397

Interphase Genes Can Be Seen in Polytene Chromosomes 401

Each Looped Domain of Chromatin May Correspond to a Separate Unit of Function 402

Different Domains of Chromatin Appear to Contain Differently Organized Nucleosomes 404

Most Chromosomal DNA Does Not Code for Essential Proteins 405

Summary 406

RNA Synthesis and RNA Processing 406

Three Different RNA Polymerases Make RNA in Eucaryotes 407

RNA Polymerase II Transcribes Some DNA Sequences Much More Often Than Others 408

Transcription Occurs on DNA Bound Up in Nucleosomes 411

New RNA Is Packaged in Ribonucleoprotein Particles 411

The Precursors of Messenger RNA Are Covalently Modified at Both Ends 412

RNA Processing Removes Long Nucleotide Sequences from the Middle of RNA Molecules	414	The Regulation of Gene Expression in Eucaryotic Cells Includes Types of Controls Not Found in Bacteria	447
Small Nuclear Ribonucleoprotein Particles May Help Guide RNA Processing	415	Eucaryotic Gene Activation Mechanisms Appear to Loosen Chromatin Structure All Along a Gene	448
Multiple Intron Sequences Can Be Removed from a Single RNA Transcript	417	Eucaryotic Gene Activation Probably Occurs in Two Stages	449
The Same RNA Transcript Can Be Processed in Different Ways to Produce mRNAs Coding for Several Proteins	418	Looped Chromatin Domains Unfold as They Are Transcribed in Polytene Chromosomes	450
Different Proteins May Be Made from a Single DNA Coding Sequence at Different Stages of Cell Development	419	Lampbrush Chromosomes Display Their Transcription Units in Extended Looped Domains	452
What Is a Gene?	421	Altering the Ends of a Primary RNA Transcript Can Change the Protein Made	454
The Export of RNA from the Nucleus Requires Molecular Signals	422	Are Important Controls Also Exerted by Alterations in the Specificity of RNA Processing and Export?	455
Ribosomal RNAs and Transfer RNAs Are Made on Tandemly Arranged Sets of Identical Genes	422	Summary	455
The Nucleolus Is a Ribosome-producing Machine	424	Gene Regulatory Mechanisms and Cell "Memory"	456
The Nucleolus Has a Highly Organized Structure	426	A Highly Condensed Fraction of Interphase Chromatin Contains Specially Inactivated Genes	456
The Nucleolus Is Reassembled on Specific Chromosomes After Each Mitosis	427	Two Subclasses of Heterochromatin Can Be Distinguished	457
Summary	427	An Inactive X Chromosome Is Inherited	458
The Nuclear Envelope	429	Eucaryotic Genes Can Be Turned Off by a Novel Heritable Mechanism	459
The Nucleus Is Enclosed by a Double Membrane	429	Cooperatively Bound Clusters of Gene Regulatory Proteins Can, in Principle, Be Directly Inherited	460
The Nuclear Lamina Helps to Determine Nuclear Shape	430	Directly Inherited Patterns of DNA Methylation Help to Control Some Mammalian Genes	460
Nuclear Transport Occurs Through Nuclear Pores	432	In Special Cases, Local DNA Sequences Are Reversibly Rearranged to Turn Genes On and Off	462
Nuclear Pores Probably Transport Large Particles Selectively	433	Selected Portions of Chromosomes Can Be Either Deleted or Amplified in Somatic Cells	464
The Inner Surface of the Nuclear Lamina Helps to Organize the Chromosomes	434	Major Rearrangements of Chromosomes Are Usually Deleterious	465
Summary	435	Summary	467
Basic Elements in the Control of Gene Expression	435	The Organization and Evolution of DNA Sequences	467
The Cells of a Multicellular Organism All Contain the Same DNA	436	Genetic Recombination Drives Evolutionary Changes	468
Different Cell Types Synthesize Different Sets of Proteins	436	A Large Fraction of the DNA of Most Eucaryotes Consists of Repeated Nucleotide Sequences	468
Different Cell Types Transcribe Different Sets of Genes	437	Frequent Genetic-Recombination Events Expand and Contract Serially Repeated Satellite DNA Sequences	468
Repressor Proteins Inhibit the Transcription of Specific Genes in Bacteria	438	The Evolution of Globins Shows How Random DNA Duplications Contribute to the Evolution of Organisms	470
Gene Activator Proteins Probably Predominate in Higher Eucaryotic Cells	440	Recombination Events Also Create New Types of Protein Chains by Joining Coding Domains	471
Two Interacting Gene Regulatory Proteins That Repress Each Other's Synthesis Produce a Stable "Memory" in a Bacterial Cell	442	Short, Interspersed Repeated Sequences Are Common in Eucaryotic Genomes	472
Gene Regulatory Proteins That Act on Many Genes Simultaneously Are Probably Used in Combinations to Generate Different Tissues in Eucaryotes	443	The Concept of "Selfish" DNA	472
In Principle, Many Different Cell Types Can Be Efficiently Specified by Combinations of a Few Gene Regulatory Proteins	444	Cataclysmic Changes in Genomes Can Increase Biological Diversity	473
Input from Several Different Gene Regulatory Proteins May Be Needed to Turn On a Single Gene	445	Summary	475
In Bacteria, DNA Supercoiling Facilitates the Transcription of Genes	446	References	476

Energy Conversion: Mitochondria and Chloroplasts

The Mitochondrion

484

The Mitochondrion Contains an Outer Membrane, an Inner Membrane, and Two Internal Compartments 485

The Inner Membrane Is Folded into Cristae 486

Mitochondrial Oxidation Begins When Large Amounts of Acetyl CoA Are Produced in the Matrix Space from Pyruvate and from Fatty Acids 488

The Citric Acid Cycle Oxidizes the Acetyl Group on Acetyl CoA to Generate NADH and FADH₂ for the Respiratory Chain 490

On the Inner Mitochondrial Membrane, a Chemiosmotic Process Converts Oxidation Energy into ATP 490

The Respiratory Chain Transfers Electrons from NADH to Oxygen 493

Energy Released by the Passage of Electrons Along the Respiratory Chain Is Stored as an Electrochemical Proton Gradient Across the Inner Membrane 495

The Energy Stored in the Electrochemical Proton Gradient Across the Inner Membrane Is Harnessed to Produce ATP and to Transport Metabolites into the Matrix Space 496

The Rapid Conversion of ADP to ATP in Mitochondria Keeps the ATP Pool in the Cytoplasm Highly Charged 498

The Difference Between ΔG° and ΔG : A Large Negative Value of ΔG Is Required for ATP Hydrolysis to Be Useful to the Cell 498

Cellular Respiration Is Remarkably Efficient 499

Summary 500

The Respiratory Chain

500

Functional Inside-Out Particles Can Be Isolated from Mitochondria 500

ATP Synthetase Can Be Purified and Added Back to Membranes in an Active Form 501

ATP Synthetase Can Function in Reverse to Hydrolyze ATP and Pump H⁺ 502

The Respiratory Chain Can Be Shown to Pump H⁺ Across the Inner Mitochondrial Membrane 503

The Respiratory Chain Contains Large Enzyme Complexes Embedded in the Inner Membrane 504

Electron Transfers Are Mediated by Random Collisions in the Bilayer Between Diffusing Donors and Acceptors 505

A Large Drop in Redox Potential Across Each of the Three Respiratory Enzyme Complexes Provides the Energy Needed to Pump Protons 506

The Mechanisms of Respiratory Proton Pumping Are Not Well Understood 507

H⁺ Ionophores Dissipate the H⁺ Gradient and Thereby Uncouple Electron Transport from ATP Synthesis 508

Respiratory Control Normally Restrains the Electron Flow Through the Chain 508

Natural Uncouplers Convert the Mitochondria in Brown Fat into Heat-generating Machines 509

Chemiosmotic Mechanisms Represent a Common Theme Among the Diversity of Bacteria 509

Summary 510

The Chloroplast

510

Chloroplasts Have an Extra Compartment but Still Resemble Mitochondria 511

Two Unique Reactions Occur in Chloroplasts: The Light-driven Production of ATP and NADPH and the Conversion of CO₂ to Carbohydrate 513

Carbon Fixation Is Catalyzed by Ribulose Biphosphate Carboxylase, the World's Most Abundant Enzyme 514

In the Carbon-Fixation Cycle, Three Molecules of ATP and Two Molecules of NADPH Are Consumed for Each CO₂ Molecule Fixed 515

In Tropical Plants, Carbon Fixation Is Compartmentalized to Facilitate Growth at Low CO₂ Concentrations 516

Light Energy Captured by Chlorophyll Is Used to Produce a Strong Electron Donor from a Weak One 517

Noncyclic Photophosphorylation Produces Both NADPH and ATP 517

Chloroplasts Can Make ATP by Cyclic Photophosphorylation Without Making NADPH 519

The Geometry of Proton Translocation Is Similar in Mitochondria and Chloroplasts 520

Like the Inner Mitochondrial Membrane, the Inner Chloroplast Membrane Contains Carrier Proteins to Facilitate Metabolite Exchange with the Cytosol 521

Chloroplasts Also Carry Out Other Biosynthesis 521

Summary 522

The Evolution of Electron-Transport Chains

522

The Earliest Cells Produced ATP by Fermentation Processes 523

The Evolution of Energy-conserving Electron-Transport Chains Enabled Anaerobic Bacteria to Use Nonfermentable Organic Compounds as a Source of Energy 523

By Providing an Inexhaustible Source of Reducing Power, Photosynthetic Bacteria Overcame a Major Crisis in the Evolution of Cells 525

Summary 527

The Biogenesis of Mitochondria and Chloroplasts

528

Mitochondria and Chloroplasts Contain Separate Genetic Systems That Are Required for Their Replication 528

As a Cell Grows, the Number of Mitochondria and Chloroplasts It Contains Is Increased by Organelle Division 529

Most Genomes of Mitochondria and Chloroplasts Are Relatively Small, Circular DNA Molecules 530

Mitochondria and Chloroplasts Contain Complete Genetic Systems 532

Yeasts Have Many Advantages for the Study of Mitochondrial Biogenesis, Including the Availability of Genetic Analysis 533

In Mammals, Mitochondrial Genes Are Maternally Inherited	533
Petite Mutants in Yeast Demonstrate the Overwhelming Importance of the Cell Nucleus in Mitochondrial Biogenesis	534
A Large Fraction of the DNA in Yeast Mitochondria Is Noncoding	535
The Complete Nucleotide Sequence of the Human Mitochondrial Genome Is Known, and It Has Some Surprising Features	536
Chloroplasts Have a More Complex Genome Than Yeast and Animal Mitochondria	537
The RNA Transcripts Made on Mitochondrial DNA Are Extensively Processed After Their Synthesis	537
Proteins Are Imported into Mitochondria and Chloroplasts by an Energy-requiring Process	538
Chloroplasts Make Most of Their Own Lipids, While Mitochondria Rely Mainly on Import	540
The Biosynthesis of Mitochondria and Chloroplasts Is Largely Controlled by the Nucleus	540
How Can Drugs That Inhibit Mitochondrial Protein Synthesis Be Used as Antibiotics Without Harming the Patient?	540
Mitochondria and Chloroplasts Have Probably Evolved from Endosymbiotic Bacteria	541
Why Do Mitochondria and Chloroplasts Have Their Own Genetic Systems?	542
Summary	543
References	544

CHAPTER

10

The Cytoskeleton

Muscle Contraction	550
A Myofibril Is the Contractile Element of a Skeletal Muscle Cell	550
Myofibrils Are Composed of Repeating Assemblies of Thick and Thin Filaments	550
Contraction Occurs as Filaments Slide Past Each Other	551
Thin Filaments Consist Principally of Actin	553
Thick Filaments Are Composed Principally of Myosin	553
ATP Hydrolysis Drives Muscle Contraction	554
Myosin is an Actin-activated ATPase	555
Myosin Heads Bind to Actin Filaments	555
Myosin Heads "Walk" Along Actin Filaments	556
Actin and Myosin Are Held in Position in the Myofibril by Other Proteins	558
Muscle Contraction Is Initiated by a Rise in Intracellular Ca^{2+}	558
Troponin and Tropomyosin Mediate the Ca^{2+} Regulation of Muscle Contraction	559
Smooth Muscle Myosin Is Activated by Ca^{2+} -dependent Phosphorylation	560
Summary	561
Ciliary Movement	561
Cilia and Flagella Propagate Bending Movements	561

The Core of a Cilium Contains a Bundle of Parallel Microtubules in a "9 + 2" Arrangement	562
Microtubules Are Hollow Tubes Formed from Tubulin	563
The Ciliary Axoneme Contains Links, Spokes, and Sidearms Made of Protein	563
The Axoneme Moves by a Sliding Microtubule Mechanism	566
Dynein Is Responsible for the Sliding	567
Cilia Can Be Dissected Genetically	567
The Conversion of Microtubule Sliding to Ciliary Bending Depends on the Inner Sheath	569
Summary	569
General Features of Microtubules and Actin Filaments as Dynamic Assemblies	569
Microtubules Are Highly Labile Structures That Are Sensitive to Specific Antimitotic Drugs	570
Actin Filaments Are Continually Formed and Broken Down in Cells	571
Specific Drugs Change the State of Actin Polymerization and Thereby Affect Cell Behavior	572
The Polymerization of Actin and Tubulin Can Be Studied <i>in Vitro</i>	573
Actin Filaments and Microtubules Are Polar Structures That Grow at Different Rates from Their Two Ends	574
Actin and Tubulin Polymers Can Undergo a "Treadmilling" of Subunits	576
Summary	577

Microtubule Organizing Centers and Microtubule-associated Proteins 578

Cytoplasmic Microtubules Are Associated with Other Proteins	578
Microtubules Grow from Discrete Organizing Centers	578
Centrioles and Basal Bodies Are Structurally Identical and Functionally Interconvertible	579
Centrioles Usually Arise by Duplication	580
What Is the Function of Centrioles?	581
Summary	581

Actin Filaments and Actin-binding Proteins in Nonmuscle Cells 582

Microvilli Contain Bundles of Actin Filaments	582
Rigid Arrays of Cross-linked Actin Filaments Play a Part in the Detection of Sound	583
Dynamic Actin-Filament Structures Occur on the Surface of Many Cells	584
Nonmuscle Cells Contain Small "Musclelike" Assemblies	586
The Assembly of Myosin in Nonmuscle Cells Is Ca^{2+} -dependent	587
Actin Filaments Are Often Anchored in Cell Membranes	588
Amoeboid and Fibroblast Locomotion Depend on Actin Filaments	588
Amoeboid Movement Is Accompanied by Gel-to-Sol Transitions in the Cytoplasm	589

Cross-linking Proteins Can Produce an Actin Gel	590	Tumor Viruses as Tools for Studying the Control of the Cell Cycle	621
Fragmenting Proteins Can Produce a Ca^{2+} -dependent Liquefaction of Actin Gels	590	Oncogenes Are Identified Through the Methods of Molecular Genetics	623
The Contraction of Cytoplasmic Gels Is Mediated by Myosin	591	The Oncogenes of RNA Tumor Viruses Are the Best Understood	625
<i>Summary</i>	591	Oncogenes Frequently Code for Protein Kinases	626
Intermediate Filaments	593	Increased Levels of a Normal Cellular Protein Can Destroy Normal Cell Growth Regulation	627
Intermediate Filaments Consist of Fibrous Polypeptides That Vary Greatly in Size	593	Active Oncogenes Have Been Isolated Directly from Human Tumors	628
Disassembly of Intermediate Filaments May Require Filament Destruction	594	<i>Summary</i>	628
Different Cell Types Contain Intermediate Filaments of Distinct Composition	594	Events in the S Phase	629
Keratin Filaments Strengthen Epithelial Cell Sheets	595	During the S Phase, Clusters of Replication Forks Become Simultaneously Active on Each Chromosome	630
Do Intermediate Filaments Have Nonstructural Functions?	596	New Histones Are Assembled into Chromatin as DNA Replicates	632
<i>Summary</i>	596	The Orientation of Replication Origins Relative to Genes May Have Important Biological Consequences	635
Organization of the Cytoskeleton	597	Different Genetic Regions on the Same Chromosome Replicate at Distinct Times During the S Phase	635
The Cytoskeleton Can Be Seen in Three Dimensions in the Electron Microscope	597	Replication Times During the S Phase Are Correlated with Interphase Chromatin Structure	637
Organelles and Soluble Proteins Can Be Associated with the Cytoskeleton	599	How Is the Timing of DNA Replication Controlled?	638
Extensive Changes in the Cytoskeleton Are Produced by a Single Gene of a Transforming Virus	600	Chromatin-bound Factors Ensure That Each Region of the DNA Is Replicated Only Once During Each S Phase	638
Microtubules May Be the Overall Organizers of the Cytoskeleton	600	<i>Summary</i>	639
The Cytoskeleton Enables Cells to Respond to the Physical Nature of a Solid Surface	602	The Logic of the Cycle	640
Cytoskeleton Organization Can Be Passed from a Parent Cell to Its Daughters	602	Most Proteins Are Synthesized Continuously Throughout Interphase	640
Cytoskeletal Organization Can Be Transmitted Across Cell Membranes	603	The Cell Cycle Proceeds by an Obligatory Series of Sequential Reactions	641
How Do Cells Move?	604	Controls Operating Early in the Cycle Largely Determine the Cell Mass at Mitosis	642
<i>Summary</i>	605	The G_2 Phase Serves to Prepare Cells for Mitosis	644
References	605	Cell-Cycle Times Can Be Drastically Shortened When Not Limited by the Rates of Biosynthesis	644
		<i>Summary</i>	645

CHAPTER

11**Cell Growth and Division**

The Control of Cell Division	611	Cell Division	646
The Cells in a Multicellular Organism Divide at Very Different Rates	612	Cell Division Is Traditionally Divided into Six Stages	646
Differences in Cell-Cycle Times Are Due Mainly to Variations in the Length of G_1	613	Some Events in Mitosis Depend on the Mitotic Spindle While Others Do Not	647
Determination of Cell-Cycle Times	613	The Spindle Contains Organized Arrays of Microtubules in Dynamic Equilibrium Between Assembly and Disassembly	647
A Special "Trigger Protein" May Control Cell Division	617	Spindle Formation at Prophase Is Largely Controlled by the Mitotic Centers That Define the Poles	652
Cell Division Is Regulated by a Variety of "Feedback Control" Mechanisms	618	Each Mitotic Chromosome Consists of Two Chromatids and a Pair of Kinetochores That Bind to the Ends of Microtubules	652
Normal Mammalian Cells Grown in Culture Will Stop Dividing When They Run Out of Space	618	The Chromosome Alignment at Metaphase Is Generated by Interactions Between the Kinetochores and Polar Spindle Fibers	653
Positional Signals Also Control Cell Division	619	Metaphase Chromosomes Are Held in a Deceptively Static State by Balanced Bipolar Forces	654
Cancer Cells Have Lost Their Normal Growth Control	620		
<i>Summary</i>	621		

The Sister Chromatids Separate Suddenly at Anaphase	655
At Anaphase the Chromosomes Are Pulled Toward the Poles and the Two Poles Are Pushed Apart	655
A Dyneinlike ATPase Is Thought to Generate the Force That Pushes the Poles Apart in Anaphase, But a Different Mechanism Seems to Move the Chromosomes to the Poles	658
At Telophase the Nuclear Envelope Initially Re-forms Around Individual Chromosomes	659
The Mitotic Spindle Determines the Site of Cytoplasmic Cleavage During Cytokinesis	660
Actin and Myosin Generate the Forces for Cleavage	662
Cytokinesis Occurs by a Completely Different Mechanism in Plant Cells with Cell Walls	663
Cytokinesis Must Allow the Cytoplasmic Organelles to Be Faithfully Inherited	666
The Elaborate Mitotic Process of Higher Organisms Evolved Gradually from the Prokaryotic Fission Mechanisms	666
Summary	668

References

CHAPTER 12 Cell-Cell Adhesion and the Extracellular Matrix

Intercellular Recognition and Cell Adhesion

Myxobacteria Exploit the Advantages of Social Behavior	674
The Assembly of Slime Mold Amoebae into a Multicellular Slug Involves Chemotaxis and Specific Cell Adhesion	674
Species-specific Cell Aggregation in Sponges Is Mediated by a Large Extracellular Aggregation Factor	676
Dissociated Embryonic Vertebrate Cells Preferentially Associate with Cells of the Same Tissue	679
Cells from Different Tissues Display a Hierarchy of Adhesiveness	680
Summary	681

Cell Junctions

Desmosomes Anchor Cells Together	682
Tight Junctions Form a Permeability Barrier Across Cell Sheets	683
Gap Junctions Allow Small Molecules to Pass Directly from Cell to Cell	684
Why Are So Many Cells Electrically and Metabolically Coupled via Gap Junctions?	686
Most Cells in Early Embryos Are Ionically Coupled	688
Cells May Control the Permeability of Their Gap Junctions	689
Gap Junctions Are Composed of Channels That Directly Connect the Interiors of the Interacting Cells	690
Summary	692

The Extracellular Matrix 692

The Extracellular Matrix Consists Primarily of Fibrous Proteins Embedded in a Hydrated Polysaccharide Gel	692
Collagen Is the Major Protein of the Extracellular Matrix	693
Collagen Chains Have an Unusual Amino Acid Composition	696
During Secretion, Procollagen Molecules Are Cleaved to Form Collagen Molecules, Which Self-assemble into Fibrils	697
Once Formed, Collagen Fibrils Are Greatly Strengthened by Covalent Cross-linking	699
Elastin Is a Cross-linked, Random-Coil Protein That Gives Tissues Their Elasticity	701
Proteoglycans and Hyaluronic Acid Are Major Constituents of the Extracellular Matrix	702
Glycosaminoglycan Chains Occupy Vast Amounts of Space and Form Hydrated Gels	705
Glycosaminoglycan Chains May Be Highly Organized in the Extracellular Matrix	705
Fibronectin Is an Extracellular-Matrix Glycoprotein That Promotes Cell Adhesion	707
The Basal Lamina Is a Specialized Extracellular Matrix That Contains a Unique Type of Collagen	709
Basal Laminae Perform Diverse and Complex Functions	711
Intracellular Actin Filament Bundles Direct the Organization of Matrix Macromolecules on Cell Surfaces and Thereby Organize the Extracellular Matrix	711
In Reciprocal Fashion, an Ordered Extracellular Matrix Influences the Organization and Behavior of the Cells It Contains	712
Summary	713

References

713

CHAPTER 13 Chemical Signaling Between Cells

Three Different Strategies of Chemical Signaling: Local Chemical Mediators, Hormones, and Neurotransmitters 718

The Hypothalamus Is the Main Regulator of the Endocrine System	719
Different Cells Respond in Different Ways to the Same Chemical Signal	719
Some Cellular Responses to Chemical Signals Are Rapid and Transient, While Others Are Slow and Long-lasting	723
Signaling Molecules Can Be Either Water-soluble or Lipid-soluble	724
Local Chemical Mediators Are Rapidly Destroyed After They Are Secreted	724
Some Signaling Molecules Released by Nerve Terminals Probably Act as Local Chemical Mediators Rather Than as Neurotransmitters	726
Some Hormones and Local Chemical Mediators Act as Specific Growth Factors	727
Summary	729

Signaling Mediated by Intracellular Receptors: Mechanisms of Steroid Hormone Action

729	
Steroid-Hormone-Receptor Complexes Bind to Chromatin and Regulate the Transcription of Specific Genes	730
Steroid Hormones Often Induce Both Primary and Secondary Responses	731
Steroid Hormones Regulate Different Genes in Different Target Cells	731
<i>Summary</i>	733

Signaling Mediated by Cell-Surface Receptors: Cyclic AMP and Calcium Ions as Second Messengers

733	
The Use of Labeled Ligands Revolutionized the Study of Cell-Surface Receptors	733
Protein Hormones and Growth Factors Are Ingested by Receptor-mediated Endocytosis	734
Cell-Surface Receptors Act as Transducers by Regulating Enzymes or Ion Channels in the Plasma Membrane	735
Cyclic AMP Is a Ubiquitous Intracellular Mediator	736
Receptor and Adenylate Cyclase Molecules Are Separate Proteins that Functionally Interact in the Plasma Membrane	737
Receptors Activate Adenylate Cyclase Molecules Indirectly Through a GTP-binding Protein	738
Ca ²⁺ Also Functions as a Ubiquitous Intracellular Mediator	739
The Activation of Some Cell-Surface Receptors Opens Membrane-bound Ca ²⁺ Channels	742
<i>Summary</i>	742

The Mode of Action of Cyclic AMP and Calcium Ions as Second Messengers

743	
Cyclic AMP Activates Intracellular Protein Kinases	743

Cyclic AMP Also Regulates the Dephosphorylation of Cellular Proteins	745
Phosphorylase Kinase Illustrates How an Enzyme Can Be Activated Only Transiently by an Extracellular Chemical Signal	746
The Reversible Covalent Modification of Proteins Regulates a Large Number of Cellular Processes	747
Ca ²⁺ Alters the Conformation of Intracellular Ca ²⁺ -binding Proteins	748
Does Cyclic GMP Act as an Intracellular Second Messenger?	749
Extracellular Signals Are Amplified Enormously by the Use of Second Messengers and Enzymatic Cascades	749
The Concentration of a Molecule Can Change Quickly If It Is Continuously Degraded or Removed at a Rapid Rate	750
Cells Can Respond Gradually or Suddenly to Signals	751
<i>Summary</i>	753

Target Cell Adaptation

754	
Some Cells Become Desensitized by Endocytosing Their Surface Receptors	754
Some Cell-Surface Receptors Are Reversibly Inactivated by Prolonged Ligand Binding	754
Morphine-induced Target Cell Desensitization Is Not the Result of Receptor Inactivation	755
Chemical Signaling in Unicellular Organisms Provides Useful Models for Study	756
Bacterial Chemotaxis Is a Simple Kind of Intelligent Behavior	757
Chemotaxis-deficient Mutants Have Revealed Four Classes of Proteins Involved in Bacterial Chemotaxis	759
Protein Methylation Is Responsible for Adaptation	760
A Model for Bacterial Chemotaxis	761
<i>Summary</i>	763

References

763	
------------	--

PART

III

From Cells to Multicellular Organisms

CHAPTER

14

Germ Cells and Fertilization

The Benefits of Sex

769	
In Multicellular Animals the Diploid Phase Is Complex and Long, the Haploid Simple and Fleeting	770

Sexual Reproduction Gives a Competitive Advantage to Organisms in an Unpredictably Variable Environment	771
Sexual Reproduction Helps Establish Favorable Alleles in a Large Population	771
New Genes Evolve by Duplication and Divergence	773
Sexual Reproduction Keeps a Diploid Species Diploid	773
A Diploid Species Has a Spare Copy of Each Gene Free to Mutate to Serve a New Function	773

A Diploid Species Can Rapidly Enrich Its Genome by the Addition of New Genes	775
<i>Summary</i>	776
Meiosis	776
Meiosis Involves Two Nuclear Divisions Rather Than One	778
Genetic Reassortment Is Enhanced by Crossing Over Between Homologous Nonsister Chromatids	778
A Synaptonemal Complex Mediates Chromosome Pairing	779
Recombination Nodules Are Thought to Mediate the Chromatid Exchanges	786
Chiasmata Play an Important Part in Chromosome Segregation in Meiosis	786
Pairing of the Sex Chromosomes Ensures That They Also Segregate	787
Meiotic Division II Resembles a Normal Mitosis	787
<i>Summary</i>	788
Gametes	788
An Egg Is the Only Cell in a Higher Animal Able to Develop into a New Individual	788
Eggs Are Highly Specialized Cells with Unique Features	789
Eggs Develop in Stages	790
Many Eggs Grow to Their Large Size Through Special Mechanisms	791
Hormones Induce Egg Maturation and Ovulation	793
Oogenesis Is Wasteful	795
Sperm Are Highly Adapted for Delivering Their DNA to an Egg	796
Sperm Are Produced Continuously in Many Mammals	797
Sperm Nuclei Are Haploid, But Sperm Cell Differentiation Is Directed by the Diploid Genome	797
<i>Summary</i>	800
Fertilization	801
A Sperm Must Be Activated Before It Can Fertilize an Egg	801
Sperm-Egg Adhesion Is Mediated by Species-specific Proteins	803
Egg Activation Is Mediated by Changes in Intracellular Ion Concentrations	804
The Rapid Depolarization of the Egg Membrane Prevents Further Sperm-Egg Fusions, Thereby Mediating the Fast Block to Polyspermy	805
The Cortical Reaction Is Responsible for the Late Block to Polyspermy	806
An Increase in Intracellular Free Ca^{2+} Initiates Egg Development	807
A Rise in the Intracellular pH in Some Organisms Induces the Late Synthetic Events of Egg Activation	808
Mammalian Eggs Can Be Fertilized <i>in Vitro</i>	808
<i>Summary</i>	809
References	811

Cellular Mechanisms of Development

Cleavage and Blastula Formation	814
Cleavage Produces Many Cells from One	814
The Polarity of the Embryo Depends on the Polarity of the Egg	815
The Blastula Consists of an Epithelium Surrounding a Cavity	815
<i>Summary</i>	816
Gastrulation, Neurulation, and Somite Formation	816
Gastrulation Transforms a Hollow Ball of Cells into a Three-layered Structure	816
The Ability of Cells to Extend, Adhere, and Contract Is the Universal Basis of Morphogenetic Movement	818
Gastrulation in Amphibians	819
Movements Are Organized About the Blastopore	820
The Endoderm Will Form the Gut and Associated Organs Such as the Lungs and Liver	821
The Mesoderm Will Form Connective Tissues, Muscles, and the Vascular and Urogenital Systems	821
The Ectoderm Will Form the Epidermis and the Nervous System	822
The Neural Tube Is Formed Through Coordinated Changes in Cell Shape	822
Blocks of Mesoderm Cells Uncouple to Form Somites on Either Side of the Body Axis	823
The Vertebrate Body Plan Is First Formed in Miniature and Then Maintained as the Embryo Grows	824
<i>Summary</i>	824
Early Steps in Pattern Formation: The Mouse	825
Mammalian Development Involves an Added Complication	826
The Steps Before Gastrulation	826
Organogenesis and Growth Before Birth	828
Studies of Chimeras Show That All of the Cells of the Very Early Mammalian Embryo Are Functionally Equivalent	828
Position in the Morula Determines the Fate of a Cell	830
A Group of Founder Cells, Rather Than a Single Founder Cell, Gives Rise to a Particular Tissue or Organ	830
Teratomas Can Arise From Embryos That Develop in the Wrong Environment	831
Cells from Teratocarcinomas Can Cooperate with Normal Cells in a Chimera to Make a Normal Mouse	832
<i>Summary</i>	832
Determination and Differentiation	832
In Higher Eucaryotes, the Behavior of a Cell Depends on Its History as well as on Its Environment and Its Genome	834

Cells Often Become Determined for a Future Specialized Role Long Before They Differentiate Overtly			
The Time of Cell Determination Can Be Discovered by Transplantation Experiments	834		
The Genetic Control of Development Is Best Studied in <i>Drosophila</i>			
The State of Determination of Imaginal Disc Cells Is Heritable	835		
Groups of Cells Occasionally Transdetermine	836		
Homoeotic Mutants Reveal Genes Whose Activities Control Cell Determination	838		
The Bithorax Complex Controls Differences Between Thoracic and Abdominal Segments	838		
The Larval Body Is Constructed by Modulation of a Fundamental Pattern of Repeating Segments	840		
Mitotic Recombination Can Be Exploited to Produce Marked Mutant Clones of Cells	841		
Sharp Demarcation Lines Separate Polyclonal Compartments	842		
Different Sets of Genes Are Active in the Cells of Different Compartments	842		
The State of Determination Is Built Up Combinatorially	844		
The Extent of Cell Proliferation in <i>Drosophila</i> Is Not Determined by Counting Cell Divisions: Fast-growing Clones May Nearly Fill Their Compartment But Do Not Make It Big	846		
Cell Determination in Vertebrates Resembles Cell Determination in <i>Drosophila</i>	846		
Summary	847		
Patterns in Space	848		
Cells Are Assigned Different Characters According to Their Positions	849		
Localized Determinants Are Sometimes Identifiable in the Cytoplasm of an Egg	850		
The Determinant of Germ-Cell Character Is Localized at One End of the <i>Drosophila</i> Egg	850		
Cell Character Is Controlled by Spatial Cues	851		
Sharp Differences of Character Emerge Gradually in an Initially Uniform Population	852		
Positional Information Is Refined by Installments	852		
Nonequivalence: Cells That Ultimately Differentiate in the Same Way Can Have Different Positional Information	853		
Cells in Separate Fields May Be Supplied with Positional Information in the Same Way But Interpret It Differently	854		
Since Embryonic Fields Are Small, Gross Features of the Adult Must Be Determined Early	855		
Summary	855		
Positional Information in Limb Development	856		
The Developing Chick Limb Can Be Analyzed in Terms of a Three-dimensional System of Coordinates	856		
The Cells of the Polarizing Region Control the Antero-Posterior Patterning of Adjacent Tissue	858		
Positional Information Along the Antero-Posterior Axis May Be Supplied by Gradations in the Magnitude of a Signal from the Polarizing-Region Cells	858		
The Polarizing Region of a Mammal or a Reptile Is Effective in the Chick Also	860		
The Parts of the Limb Are Laid Down in Succession Along the Proximo-Distal Axis	860		
The Apical Ectodermal Ridge Delimits the Special Region of Mesenchyme from Which Successive Distal Parts Develop, But It Does Not Instruct the Mesenchyme as to Which Parts It Should Form	860		
Positional Specification Along the Proximo-Distal Axis Depends on the Amount of Time Spent in the Progress Zone	861		
Neighboring Mesenchyme Cells in Early Chick Limb Buds May Interact so as to Smooth Out Discontinuities in the Pattern of Positional Values	863		
Some Limbs Can Regenerate	864		
Cockroach Legs Undergo Intercalary Regeneration	864		
The Same Pattern of Positional Values Is Repeated in Successive Segments of the Cockroach Leg	866		
Circumferential Intercalation Obeys the Same Rule as Proximo-Distal Intercalation	866		
Intercalation in the Epidermis Is a Two-dimensional Problem	867		
Regeneration of a Two-dimensional Patch Obeys the Rule of Intercalation	868		
The Rule of Intercalation May Apply to Many Different Systems	869		
Summary	869		
Inductive Interactions in the Development of Epithelia	870		
Mesoderm Induces Ectoderm to Form the Different Parts of the Neural Tube	870		
The Dermis Controls the Nature and Pattern of the Structures That Form from the Epidermis	870		
Epithelium Invades Mesenchyme to Form the Tubules of a Gland	872		
Summary	872		
Multicellular Development Studied Cell by Cell: The Nematode Worm	873		
<i>Caenorhabditis elegans</i> Is Anatomically and Genetically Simple	873		
Nematode Development Is Essentially Invariant	874		
Cytokinesis Is Not Required for Cell Differentiation	875		
The Influence of Local Cell Interactions on the Behavior of Cells During Development Can Be Studied by Laser Microsurgery	876		
An "Anchor Cell" Controls Vulva Development	876		
A "Distal Tip Cell" Causes Continued Proliferation of Nearby Germ Cells	877		
Cell Fate Can Be Controlled by Inhibitory Interactions Among Cells in an Equivalence Group	878		
Summary	879		
Migratory Cells	880		
Cells Wander: Selective Cohesion May Stop Them from Straying from Their Proper Place	880		

Germ Cells Leave the Yolk Sac and Settle in the Genital Ridges	880
Muscle Cells in the Chick Limb Originate by Migration from the Somites	881
Cells Disperse from the Neural Crest and Form Many Different Tissues	881
The Pathways of Migration Are Defined by the Host Connective Tissue	882
The Differentiation of Neural Crest Cells Is Decided by the Local Environment	884
The Development of the Nervous System Poses Special Problems	884
Summary	885
References	885

CHAPTER

16

Differentiated Cells and the Maintenance of Tissues

Maintenance of the Differentiated State	891
Differentiated Cells Commonly Remember Their Character Even in Isolation: Pigment Epithelium of the Retina	892
The Extracellular Matrix That a Cell Secretes Helps Maintain the Cell's Differentiated State	893
Cell-Cell Interactions Can Modulate the Differentiated State	894
Some Structures Are Maintained by a Continuing Interaction Between Their Parts: Taste Buds and Their Nerve Supply	895
An Agent That Causes Changes in DNA Methylation Can Cause Radical Alterations of Differentiated Character	896
Summary	897
Tissues With Permanent Cells	897
The Cells at the Center of an Adult Lens Are Remnants of the Embryo	897
Most Permanent Cells Renew Their Parts: Photoreceptor Cells of the Retina	899
Summary	902
Renewal by Simple Duplication	902
The Liver Is an Interface Between the Digestive Tract and the Blood	903
Liver Cell Loss Stimulates Liver Cell Proliferation	905
Regeneration May Be Hindered by Uncoordinated Growth of the Components of a Mixed Tissue	906
Endothelial Cells Constitute the Fundamental Component of All Blood Vessels	906
New Endothelial Cells Are Generated by Simple Duplication of Existing Endothelial Cells	908
New Capillaries Form by Sprouting	908
Growth of the Capillary Network Is Controlled by Factors Released by the Surrounding Tissues	909
Summary	910
Renewal by Stem Cells: Epidermis	910
Stem Cells Have the Ability to Divide Without Limit and to Give Rise to Differentiated Progeny	911

The Epidermis Is Organized into Proliferative Units	912
Differentiating Epidermal Cells Synthesize a Sequence of Different Keratins as They Mature	914
For Each Proliferative Unit, There Is an "Immortal" Stem Cell	915
Stem Cell Potential May Be Maintained by Contact with the Basal Lamina	915
Basal Cell Proliferation Is Regulated According to the Thickness of the Epidermis	916
Secretory Cells of the Skin Are Secluded in Glands and Have Different Population Kinetics	916
Summary	918
Renewal by Pluripotent Stem Cells: Blood Cell Formation	918
New Blood Cells Are Generated in the Bone Marrow	921
Bone Marrow Contains Pluripotent Stem Cells That Can Establish Hematopoietic Colonies	923
The Number of Specialized Blood Cells Is Amplified by Divisions That Follow Commitment	924
Production of Erythrocytes Is Controlled Through Hormonal Regulation of the Cell Divisions That Follow Commitment	925
Specific Glycoprotein Hormones Control the Survival and Fate of the Different Classes of Committed Hematopoietic Precursor Cells	927
Summary	927
Quiescent Stem Cells: Skeletal Muscle	928
Skeletal Muscle Cells Do Not Divide	929
New Skeletal Muscle Cells Form by Fusion of Myoblasts	929
Muscle Differentiation Requires Coordinated Changes in the Expression of Many Different Genes	929
Some Myoblasts Persist as Satellite Cells in the Adult	931
The State of Differentiation of Skeletal Muscle Fibers Can Be Modulated by Electrical Stimulation	931
Summary	933
Soft Cells and Tough Matrix: Growth, Turnover, and Repair in Skeletal Connective Tissue	933
Cartilage Grows by Swelling	933
Osteoblasts Secrete Bone Matrix While Osteoclasts Erode It	934
Cartilage Is Eroded by Osteoclasts to Make Way for Bone	937
Summary	938
Territorial Stability in the Adult Body	938
Epithelial Organization Helps to Keep Cells in Their Proper Territories	939
Normal Somatic Cells Are Destined to Die for the Sake of the Survival of the Germ Cell	939
Cancer Cells Break the Rules of Altruistic Social Behavior	940
Summary	941
Appendix	941
References	947

The Immune System**The Cellular Basis of Immunity****952**

The Immune System Is Composed of Billions of Lymphocytes 952

B Lymphocytes Make Humoral Antibody Responses; T Lymphocytes Make Cell-mediated Immune Responses 952

All Lymphocytes Develop from Pluripotent Hemopoietic Stem Cells 954

Cell-Surface Markers Make It Possible to Distinguish and Separate T and B Cells 955

Most Lymphocytes Continuously Recirculate Between the Blood and Lymph 956

The Immune System Works by Clonal Selection 956

Most Antigens Stimulate Many Different Lymphocyte Clones 958

Immunological Memory Is Due to Clonal Expansion and Lymphocyte Differentiation 959

The Failure to Respond to Self Antigens Is Due to Acquired Immunological Tolerance 960

Immunological Tolerance to Foreign Antigens Can Also Be Induced in Mature Animals 962

Summary 963**The Functional Properties of Antibodies****963**

The Antigen-specific Receptors on B Cells Are Antibody Molecules 964

B Cells Can Be Stimulated to Make Antibodies in a Culture Dish 964

Antibodies Have Two Identical Antigen-binding Sites 964

An Antibody Molecule Is Composed of Four Polypeptide Chains—Two Identical Light Chains and Two Identical Heavy Chains 965

There Are Five Different Classes of H Chains, Each with Different Biological Properties 966

Antibodies Can Have Either κ or λ Light Chains But Never Both 969

The Strength of an Antibody-Antigen Interaction Depends on Both the Affinity and the Number of Binding Sites 969

Antibody-Antigen Interactions Can Be Measured in Many Ways 971

The Size of the Antigen-Antibody Complexes Formed Depends on the Valence of the Antigen and on the Relative Concentrations of the Antigen and Antibody 972

Antibodies Recruit Complement and Various Cells to Fight Infection 973

Summary 974**The Fine Structure of Antibodies****974**

Myeloma Proteins Are Homogeneous Antibodies Made by Plasma-Cell Tumors 974

L and H Chains Consist of Constant and Variable Regions 975

The L and H Chains Each Contain Three Hypervariable Regions That Together Form the Antigen-binding Site 975

The L and H Chains Are Folded into Repeating Domains 976

X-ray Diffraction Studies Have Revealed the Structure of Immunoglobulin Domains and Antigen-binding Sites in Three Dimensions 977

Summary 979**The Generation of Antibody Diversity****979**

More than One Gene Segment Codes for Each L and H Chain 980

Two Gene Segments Code for the V Region of Each L Chain 981

Three Gene Segments Code for the V Region of Each H Chain 983

Antibody Diversity Is Increased by Somatic Recombination, by the Combinatorial Joining of Light and Heavy Chains, and by Somatic Mutation 983

The Mechanisms of Antibody Gene Expression Ensure That B Cells Are Monospecific 984

The Switch from a Membrane-bound to a Secreted Form of the Same Antibody Occurs Through a Change in the H-Chain RNA Transcripts 985

B Cells Can Switch the Class of Antibody They Make 986

Idiotypes on Antibody Molecules Form the Basis of an Immunological network 986

Summary 988**The Complement System****988**

Complement Activation Involves a Sequential Proteolytic Cascade 988

The Classical Pathway Is Activated by Antibody-Antigen Complexes 989

The Alternative Pathway Can Be Directly Activated by Microorganisms 990

The Assembly of the Late Complement Components Generates a Transmembrane Lytic Complex 991

The Complement Cascade Is Tightly Regulated and Designed to Attack a Nearby Membrane 992

Summary 993**T Lymphocytes and Cell-mediated Immunity****993**

The T-Cell-Receptor Enigma 993

Different T-Cell Responses Are Mediated by Different T-Cell Subpopulations 994

Cytotoxic T Cells Kill Virus-infected Cells 995

Helper T Cells Are Required for Most B Cells and T Cells to Respond to Antigen 995

Helper T Cells Activate Macrophages by Secreting Lymphokines 996

Suppressor T Cells Inhibit the Responses of Other Lymphocytes 997

Helper and Suppressor T Cells Can Recognize Foreign Antigens on the Target Lymphocyte Surface 997

Regulatory T Cells May Communicate with Their Target Lymphocytes by Secreting Soluble Helper or Suppressor Factors 998

Transplantation Reactions Are T-Cell-mediated Immune Responses 999

18

Cells of the Nervous System: A Preliminary Sketch

Nerve Cells Carry Electrical Signals	1015
Nerve Cells Communicate Chemically at Synapses	1016
Neural Tissue Consists of Neurons and Glial Cells	1016
<i>Summary</i>	1017

Voltage-gated Channels and the Action Potential

The Na ⁺ -K ⁺ Pump Charges the Battery That Powers the Action Potential	1018
The Membrane Potential Depends on Selective Membrane Permeability	1019
Ion Channels Are Characterized by Their Selectivity, Their Gating, and Their Sensitivity to Specific Toxins	1022
Depolarization Causes Na ⁺ Channels First to Open and Then to Become Inactivated	1023
Fluctuations in the Transmembrane Current Suggest That Individual Channels Are Opening and Closing Randomly	1023
Gated Channels Open and Close in an All-or-None Fashion	1024
The Membrane Electric Field Controls the Energies of the Different Channel Conformations	1025
Voltage-gated Na ⁺ Channels Are Responsible for the Action Potential	1027
Action Potentials Are All-or-None	1028
Voltage Changes Can Spread Passively Within a Neuron	1028

The Membrane Potential in the Cell Body Represents a Spatial Summation of Postsynaptic Potentials	1046
Temporal Summation Translates the Frequency of Presynaptic Signals into the Size of a PSP	1046
The Grand PSP Is Translated into Nerve Impulse Frequency for Long-Distance Transmission	1047
Encoding Requires a Combination of Different Voltage-sensitive Channels	1048
Early K ⁺ Channels Help to Make the Firing Rate Proportional to the Stimulus	1049
Adaptation Lessens the Response to an Unchanging Stimulus	1049
Not All Signals Are Delivered via the Axon	1050
<i>Summary</i>	1052

Channel Regulation and Memory 1052

The Distribution of Ion Channels in a Muscle Cell Changes in Response to Denervation	1053
The Receptivity of a Muscle Cell Can Be Controlled by Electrical Stimulation	1054
The Site of a Synaptic Contact Is Marked by a Persistent Specialization of the Basal Lamina	1054
Synaptic Plasticity Provides a Mechanism for Memory	1055
A Short-Term Memory Is Registered by Modification of Channel Proteins	1056
Memory Remains Mysterious	1058
<i>Summary</i>	1058

Sensory Input 1058

Stimulus Magnitude Is Reflected in the Receptor Potential	1059
Sense Receptors Are Tuned to Detect Specific Stimuli	1061
Rod Cells Can Detect a Single Photon	1063
The Visual World Is Mapped onto a Sequential Hierarchy of Arrays of Neurons	1064

Neurons at Higher Levels Detect More Complex Features of the Visual World
 Information Is Processed Through Local Computations Involving Convergence, Divergence, and Lateral Inhibition
Summary

Maintenance and Development of Nerve Cell Structure

Axonal Transport Maintains Chemical Communication Between the Cell Body and the Outlying Parts of a Neuron
 Axonal Transport Has Both Fast and Slow Components
 Developing Axons and Dendrites Terminate in a Growth Cone
 Cytoskeletal Proteins and Membrane for Outgrowth Are Inserted at Different Sites
 Without Microtubules, the Developing Neuron Retracts; Without Actin Filaments, It Cannot Advance
 Growth Cones Can Be Guided Along Chemically Defined Tracks in the Substratum
 The Growth Cones of Some Types of Neurons Are Also Guided by Chemotactic Molecules Such as Nerve Growth Factor
 The Growth Cone Must Recognize Its Target and Finally Halt
Summary

The Development of Neuromuscular Connections

The Motor Neurons That Control the Limbs Are Generated in the Neural Tube Epithelium
 Radial Glial Cells Form a Temporary Scaffold to Guide the Migrations of the Immature Neurons
 Axons Grow Out Along Precisely Defined Pathways to Specific Target Areas
 The Pattern of Nerve Connections Is Governed by the Nonequivalent Character of Cells, Not Merely by Their Positions
 Cells That Fail to Make Connections Die
 Normal Motor Neuron Death Is Prevented by Toxins That Block Neuromuscular Transmission
 Surplus Synapses Are Eliminated by a Competitive Process
 Denervated Muscle Cells Release a Factor That Stimulates Nerve Cells to Sprout
Summary

Neural Maps and the Development of the Visual System

Neuronal Specificity Appears to Control the Mapping of the Visual World from the Retina onto the Tectum
 Neuronal Specificity Does Not Exert Absolute Control over the Formation of the Neural Map
 Visual Connections in Young Mammals Are Adjustable and Sensitive to Visual Experience
 In the Visual Cortex of the Brain, the Projections from the Two Eyes are Mapped in Alternating Stripes
 Active Synapses Tend to Displace Inactive Synapses

Synchronously Active Synapses on the Same Cell Reinforce One Another; Asynchronously Active Synapses Compete for Survival
 The Pattern of Connections That Develops in the Brain Reflects the Regular Associations Between Events in the External World
Summary

References

1070

1072

1072

1074

1075

1076

1076

1077

1077

1078

1078

1079

1080

1081

1083

1083

1084

1085

1085

1086

1086

1088

1088

1089

1090

CHAPTER

19

Special Features of Plant Cells

The Central Significance of the Cell Wall 1099

The Cell Wall Is Composed of Cellulose Fibers Embedded in a Matrix of Polysaccharide and Protein 1100
 Fiber and Matrix Polysaccharides Form a Complex Cross-linked Structure 1101
 The Limited Porosity of Cell Walls Limits Molecular Exchange Between Plant Cells and Their Environment 1103
 The Tensile Strength of Cell Walls Allows Plant Cells to Generate an Internal Hydrostatic Pressure Called Turgor 1104
 Turgor Is Regulated by Feedback Mechanisms That Control the Concentration of Intracellular Solutes 1105
 The Cell Wall Is Modified During the Creation of Specialized Cell Types 1106
 The Wall of the Mature Cell Is Adapted to Its Function 1108
Summary 1110

The Interactions and Communication Between Cells 1110

Plant Cells Are Connected to Their Neighbors by Special Cytoplasmic Channels Called Plasmodesmata 1110
 Plasmodesmata Allow Molecules to Pass Directly from Cell to Cell 1111
 The Fluids in a Plant Are Segregated into One Large Intracellular Compartment and One Large Extracellular Compartment 1113
 Fluids Are Transported Throughout the Plant by Xylem and by Phloem 1113
 Chloroplasts Use Light Energy to Fix CO₂, Enabling the Photosynthetic Tissues to Export Sucrose 1116
 Symbiotic Bacteria Allow Some Plants to Use Atmospheric Nitrogen 1116
 Cell-Cell Recognition in Plants Involves Specific Sequences of Sugar Residues 1117
Summary 1118

The Internal Organization of Plant Cells 1119

The Chloroplast Is One Member of a Family of Organelles, the Plastids, Unique to Plants 1120
 The Plant Cell Vacuole Is a Remarkably Versatile Organelle 1122
 As Plants Grow, They Accumulate Water in Their Vacuoles 1123

Vacuoles Can Function as Storage Organelles	
Plant Cells Exocytose But Generally Seem Not to Endocytose Macromolecules	
Golgi Vesicles Deliver Cell-Wall Material to Specific Regions of Plasma Membrane	
Cellulose Synthesis Occurs at the Surface of Plant Cells	
Cortical Microtubules Orient the Extracellular Deposition of Cellulose Microfibrils	
The Movement of Materials in Large Plant Cells is Driven by Cytoplasmic Streaming	
The Interaction of Actin and Myosin Drives Cytoplasmic Streaming in Giant Algal Cells	
Regions of the Plant Cell Cytoskeleton Can Be Reorganized in Response to Local Stimuli	
Summary	

1124	Cell Growth and Division	1133
1124	Most New Plant Cells Arise in Special Areas Called Meristems	1134
1125	The Shape of a Growing Plant Cell Is Determined by the Organization of Cellulose Microfibrils	1135
1128	A Preprophase Band of Microtubules Marks the Future Plane of Cell Division	1136
1128	Hormones Help Control the Growth and Shape of Plants	1139
1130	Tissue Culture Facilitates Studies of Mechanisms of Cell Determination in Plants	1139
1131	Plant Cells Without Their walls Can Be Manipulated Much Like Animal Cells	1142
1132	Summary	1143
1133	References	1144