Ilia J. Leitch Genome size diversity and why it matters - insights from the Plant DNA C-values database Jiří Vondrášek ELIXIR CZ meets challenges of data management in Life Sciences Petr Novák

ELIXIR CZ supported tools for genome annotations and data sharing

## Tony Heitkam

Repetitive DNA in plants: a guide to genomic change

and regulatory innovation

Nicola Schmidt

Beet 'em up: Unifying 30 years of repeat research in beets and wild beets

Zdeněk Kubát

LTR retrotransposons preferentially spread through either the male or female plant germline

Pavel Jedlička

Reconstruction of LTR retrotransposons insertions in plant genomes

## Andre Marques

How to evolve without centromeres?

- Jiří Macas
- Weird but useful: what we can learn from atypical plant centromeres
- Yi-Tzu Kuo
- Plasticity in holocentromere organization: The holocentromere of Chionographis is comprised of a few megabase-sized CENH3 positive satellite arrays
- Pavel Neumann
- Identification of changes in structural and regulatory kinetochore protein genes in holocentric compared to monocentric Cuscuta species

- Andreas Houben
- Analysis and application of natural and artificially induced
- chromosome elimination
- Steven Dreissig
- Recombination landscape divergence between populations is marked
- by larger low-recombining regions in domesticated rye
- Václav Bačovský
- Analysis of synaptonemal complex in plants: dissection
- of SCs in Silene latifolia
- Jana Szecówka
- Both male and female meiosis contribute to non-Mendelian inheritance of parental chromosomes in interspecific plant hybrids (Lolium × Festuca)

Ingo Schubert DNA double-strand breaks as drivers of genome and karyotype evolution Martin Kováčik Developing an atlas of gene expression during barley grain development Yile Huang

Structure and evolution of the meso-octoploid genome of Heliophila variabilis (Brassicaceae)

Tomáš Janíček Back to the Future: Sex chromosomes and flower development Miroslava Karafiátová Mysterious biology of the B chromosome in wild Sorghum

## Petr Smýkal Towards the understanding of the seed dormancy as the key domestication trait in legumes