

Contents

About the Author	xi
Preface	xiii
Acknowledgements	xvii
1 General Introduction and Formulation of the Problem	1
1.1 A Closed Forest as Natural Vegetation	1
1.2 Formulation of the Problem	6
1.3 The Null Hypothesis and the Alternative Hypothesis	8
1.4 Outline of this Study	9
2 Succession, the Climax Forest and the Role of Large Herbivores	13
2.1 Introduction	13
2.2 History of the Concept of Succession	14
2.3 Clements's Theory of Succession	16
2.4 Tansley's Polyclimax Theory	18
2.5 The Lack of Regeneration in the Climax Forest	20
2.6 Secondary Succession from Grassland to Forest	21
2.7 Regeneration of the European Primeval Forest	28
2.8 The East of North America as an Analogy of Europe	30
2.9 Pioneer and Climax Species	35
2.10 Variations on the Succession Theme and the Gap Phase Model	37
2.11 Catastrophes as the Mechanism of Succession, Regeneration and Diversity	38

2.12	Establishment Factors in Relation to Succession and Diversity	44
2.13	Nutrients in Relation to Establishment and Succession	47
2.14	The Role of Large Mammals in Succession	51
2.15	Conclusions and Synthesis	57
3	Palynology, the Forest as Climax in Prehistoric Times and the Effects of Humans	61
3.1	Introduction	61
3.2	The Reconstruction of Prehistoric Vegetation	62
3.3	The Forest in Palynology	66
3.4	Palynology and Theories about Succession and the Forest as the Climax Vegetation	70
3.5	Drawing up Corrective Factors for Pollen Frequencies	76
3.6	The 'Landnam' Theory	78
3.7	The Closed Forest versus the Half-open Park Landscape	85
3.8	The Situation in the East of the United States	95
3.9	Conclusions and Synthesis	96
4	The Use of the Wilderness from the Middle Ages up to 1900	102
4.1	Introduction	102
4.2	The Wilderness and the Concept of 'Forestis'	103
4.3	The Meanings of the Term 'Wald'	109
4.4	The Meaning of the Terms 'Holt' and 'Bosch' in relation to 'Wald'	115
4.5	What was a 'Silva'?	119
4.6	Regulations on the Use of the Wilderness	123
4.7	Regulations on Grazing Livestock	129
4.8	Grazing Livestock in Relation to Coppices	132
4.9	Seedlings of Trees in Scrub	138
4.10	The Regeneration of Trees and the Grazing of Livestock	144
4.11	Coppices-with-standards (Standing Trees) Growing in Scrub	156
4.12	The 'Hage' or 'Haye'	159
4.13	The Vegetative and Generative Regeneration of Trees in Relation to the Grazing of Livestock	162
4.14	The Development of 'Natural' Regeneration	168
4.15	Grazing Livestock and the Destruction of the Forest	170
4.16	'Having but Little Wood, and that Oke like Stands left in our Pastures in England'	176
4.17	European Mental Models in North America	178
4.18	Conclusions and Synthesis	183
5	Spontaneous Succession in Forest Reserves in the Lowlands of Western and Central Europe	189
5.1	Introduction	189

5.2	La Tillaie and Le Gros-Fouteau in the Forêt de Fontainebleau, France	191
5.2.1	A brief description and history of the reserves	191
5.2.2	The present situation	196
5.3	The Neuenburger Urwald, Germany	206
5.3.1	A brief description and history of the reserve	206
5.3.2	The present situation	209
5.4	The Hasbrucher Urwald, Germany	212
5.4.1	A brief description and history of the reserve	212
5.4.2	The present situation	213
5.4.3	Gaps in the canopy in the Hasbrucher Urwald and Neuenburger Urwald	213
5.5	Sababurg in the Reinhardswald, Germany	215
5.6	Rohrberg in the Spessart, Germany	217
5.6.1	A brief description and history of the reserve	217
5.6.2	The present situation	218
5.7	Priorteich in the Southern Harz, Germany	220
5.7.1	A brief description and history of the reserve	220
5.7.2	The present situation	221
5.8	Forest Reserves in North Rhine-Westphalia, Germany	224
5.8.1	General characteristics	224
5.8.2	The present situation	224
5.9	The Oak Reserve Johannser Kogel in the Wienerwald, Austria	229
5.9.1	A brief outline and history of the reserve	229
5.9.2	The present situation	229
5.10	The So-called Primeval Forest Reserve of Krakovo, Slovenia	235
5.10.1	General characteristics	235
5.10.2	The present situation	236
5.11	The Unterhölzer at Donaueschingen in the Black Forest, Germany	237
5.12	Dalby Söderskog, Sweden	240
5.12.1	A brief description and history	240
5.12.2	The present situation	241
5.13	The Forest of Bialowieza, Poland	245
5.13.1	A brief description of the forest of Bialowieza	245
5.13.2	The history of the forest of Bialowieza	247
5.13.3	The present situation in the National Park of Bialowieza	255
5.13.4	Regeneration of oak in the thermophile oak forest (<i>Potentillo albae-Quercetum</i>)	267
5.13.5	Is the thermophile oak forest or the lime-hornbeam forest most similar to the original vegetation?	270
5.14	Other Forest Reserves in Europe	274

5.15	Spontaneous Succession in Forest Reserves in the East of the United States	276
5.16	Conclusions and Synthesis	281
6	Establishment of Trees and Shrubs in Relation to Light and Grazing	287
6.1	Introduction	287
6.2	Pedunculate Oak (<i>Q. robur</i>) and Sessile Oak (<i>Q. petraea</i>)	289
6.2.1	The reaction of pedunculate and sessile oak seedlings to reduced amounts of daylight	289
6.2.2	The establishment of both species of oak in relation to the alternative hypothesis	300
6.3	Beech (<i>F. sylvatica</i>)	308
6.3.1	The response of beech seedlings to a reduction in the amount of daylight	308
6.3.2	A comparison of beech and sessile oak seedlings	312
6.3.3	The establishment of beech in relation to the alternative hypothesis	319
6.4	Broad-leaved Lime (<i>Tilia platyphyllos</i>) and Small-leaved Lime (<i>T. cordata</i>)	322
6.4.1	The response of broad-leaved and small-leaved lime seedlings to reduced amounts of daylight	322
6.4.2	A comparison of broad-leaved and small-leaved lime seedlings and pedunculate and sessile oak seedlings	324
6.4.3	The establishment of both species of lime in relation to the alternative hypothesis	329
6.5	Hornbeam (<i>Carpinus betulus</i>)	330
6.5.1	The response of hornbeam seedlings to a reduction in the amount of daylight	330
6.5.2	The establishment of hornbeam in relation to the alternative hypothesis	332
6.6	Ash (<i>Fraxinus excelsior</i>), Field Maple (<i>Acer campestre</i>), Sycamore (<i>A. pseudoplatanus</i>) and Elm (<i>Ulmus</i> sp.)	332
6.6.1	The response of seedlings of ash, field maple, sycamore and elm to reduced amounts of daylight	332
6.6.2	Ash, field maple, sycamore and elm in relation to the alternative hypothesis	333
6.7	Hazel (<i>Corylus avellana</i>)	333
6.7.1	The response of hazel seedlings to reduced amounts of daylight	333
6.7.2	Hazel in relation to the alternative hypothesis	337

6.8	Wild Fruits, Wild Apple (<i>Malus Sylvestris</i>), Wild Pear (<i>Pyrus pyraster</i>), Wild Cherry (<i>Prunus avium</i>), and Members of the Rowan Family (<i>Sorbus</i> spp.)	339
6.8.1	The establishment of wild fruit trees	339
6.8.2	Wild fruit and members of the rowan family in relation to the alternative hypothesis	341
6.9	The Establishment of Blackthorn, Hawthorn, Juniper and other Thorny Shrubs in Relation to Grazing and Diversity	342
6.9.1	Germination conditions	342
6.10	The Establishment of Trees and Shrubs at the End of the Last Ice Age	345
6.10.1	Grazing and the establishment of trees and shrubs in prehistoric times	345
6.10.2	Continuity in diversity	356
6.11	A Park Landscape in the East of the United States?	359
6.12	Conclusions and Synthesis	363
7	Final Synthesis and Conclusions	369
7.1	The Null Hypothesis and the Alternative Hypothesis	369
7.2	The Findings	370
7.3	Final Conclusion	376
7.4	Epilogue	379
Appendix 1	Structuring the Use of the Wilderness	385
Appendix 2	The Use of the Wilderness	391
Appendix 3	The Destruction of the Wilderness	393
Appendix 4	The 'Hanau-Münzenbergse Forstordnung', the 'Mainzer Forstordnung' and the 'General-Verordnung'	397
Appendix 5	The Development of 'Natural' Regeneration	403
Appendix 6	The Selection System	407
Appendix 7	The Group Selection System	410
Appendix 8	Species of Trees and Shrubs in Landscapes Grazed by Large Herbivores	412
Appendix 9	Species of Herbs on the Fringes of Scrub and Groves	414